



## City Research Online

### City, University of London Institutional Repository

---

**Citation:** Barlow, S., Dixey, R., Todd, J., Taylor, V., Carney, S. & Newell, R. (2014). 'Abandoned by Medicine'? A qualitative study of women's experiences with lymphoedema secondary to cancer, and the implications for care. *Primary Health Care Research & Development*, 15(04), pp. 452-463. doi: 10.1017/s1463423613000406

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

---

**Permanent repository link:** <https://city-test.eprints-hosting.org/id/eprint/14992/>

**Link to published version:** <https://doi.org/10.1017/s1463423613000406>

**Copyright:** City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

**Reuse:** Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.



Primary Health Care Research and Development

**‘Abandoned by Medicine’? A Qualitative Study of Women’s Experiences with Lymphoedema Secondary to Cancer, and the Implications for Care.**

Sally Barlow<sup>1</sup>, Rachael Dixey<sup>2</sup>, Jacquelyne Todd<sup>3</sup>, Vanessa Taylor<sup>4</sup>, Sarah Carney<sup>5</sup>, and Rob Newell<sup>6</sup>

<sup>1</sup> Lecturer in Mental Health Nursing, School of Health Studies, University of Bradford, Bradford, UK

<sup>2</sup> Professor of Health Promotion, Leeds Metropolitan University, Leeds, UK

<sup>3</sup> Physiotherapist Consultant in Lymphoedema, Wharfedale Hospital, Leeds, UK

<sup>4</sup> Discipline Lead-Nursing: Higher Education Academy

<sup>5</sup> Co-Researcher: University of Bradford

<sup>6</sup> Emeritus Professor in Nursing Research, School of Health Studies, University of Bradford, Bradford, UK

Name and address of the corresponding author:

Dr Sally Barlow  
School of Health Studies  
University of Bradford  
Horton A Building,  
Richmond Road,  
Bradford,  
West Yorkshire  
BD7 1DP,  
UK  
Email: s.barlow1@bradford.ac.uk

## Abstract

**Background:** Lymphoedema secondary to cancer is a relatively neglected and under-researched condition. Few studies report people's experiences of care and treatment provision when living with the condition. Current practice focuses on the physical treatment yet psychosocial needs often remain unmet. A previous study examining the patient perspective identified the theme of being 'abandoned by medicine'. Perceived lack of support may result in a delayed adaptation and acceptance of this long term condition and can significantly impact on psychological wellbeing. We explore this emerging theme alongside others in order to provide a guide to action for improvements for patient benefit.

**Aim:** The central aim is to explore women's views of their care and treatment following a diagnosis with lymphoedema secondary to cancer. This forms part of a larger study aimed at assessing appropriate screening tools to measure psychosocial distress.

**Methods:** A mixed-methods approach was used for the main study. Here we report the qualitative component, derived from in-depth semi-structured interviews conducted in the homes of the participants (n=14) and focus group discussions (n=15). In addition, qualitative comments from questionnaire data from a large scale postal survey are included (n=104).

**Findings:** Participants identified considerable deficiencies in health care workers' knowledge and awareness of lymphoedema, which subsequently impacted on the patients' needs for information, support and understanding. Access to appropriate treatment was patchy and problems were identified with the process of obtaining compression garments, massage and other sources of help. Although lymphoedema is a long term condition and a disfiguring one, and much is known about how these impact on patients' emotional well-being, little attention was paid by health professionals to potential psychosocial consequences. In essence women had to become experts of their own condition and cope as best as they could. We provide recommendations to improve service delivery and address these unmet needs.

**Key words:** lymphoedema, psychosocial, body image, supported self-care, cancer, treatment.

## Introduction

Lymphoedema is a chronic, incurable and progressive condition (Browse, 2003, Framework, 2006). It is characterised by a disfiguring swelling resulting from the accumulation of lymph fluid in the interstitial spaces of a limb or affected area (Fu and Rosedale, 2009). Secondary lymphoedema (SL) may occur as a consequence of disruption to the lymphatics from cancer or its related therapy (Keeley, 2000). Approximately 20% of breast cancer patients develop swelling of the upper limb (Clark et al., 2005, Todd et al., 2008) and between 28% and 47% develop swelling of the lower limb after gynaecological surgery (Moffatt et al., 2003, Ryan et al., 2003). Although body image concerns in breast cancer patients have been well-researched (Fobair et al., 2006) focus on lymphoedema is often lacking or dismissed. This is despite it being an “all too common occurrence” (Rockson, 2000:2814) after breast cancer surgery. Overall this is an under-researched area in the UK. Insights into how women adjust to the diagnosis of this long-term condition, consideration of their psychosocial needs and qualitative research that gives them a voice is minimal.

The literature exploring people’s adjustment to a diagnosis of lymphoedema and subsequent consequences for their mental health and wellbeing is scanty. In particular the psychosocial ramifications of the condition require specific attention in addition to a focus on physical function (Ridner et al., 2012). Early studies emphasised this unmet need. Carter (1997), for example, researching women’s experiences of lymphoedema in the USA, called for more research into the psychosocial impact on body image, self-esteem, and social support. In addition the women reported that physicians did not know enough about lymphoedema and patients felt they needed more advice. Three themes emerged: abandonment by medicine, concealing the imperfect body, and living the interrupted life. Earlier, Woods (1993) described lymphoedema as a ‘complex’ experience for patients and also reported a lack of social support and support from healthcare professionals. This provision remains missing despite research revealing that women with lymphoedema following breast cancer, who perceived less social support, experienced greater overall levels of distress (Passik et al., 1995).

Lymphoedema affects in excess of 80,000 patients in the UK, and is associated with greater anxiety and distress and poorer mental wellbeing and quality of life (Morgan et al., 2005). Passik and McDonald (1998:2818) demonstrate a further complication in women with cancer-related SL; they describe lymphoedema as a “secondary blow to their physical and emotional wellbeing”. Moreover, women with post-breast cancer lymphoedema report more psychological distress than matched controls (Tobin et al., 1993). Another study confirms cancer-related lymphoedema has a significant impact on quality of life (Beaulac et al., 2002).

## **Aims**

The rationale for this research arose from two main insights. The first was recognition among clinicians of insufficient evidence to facilitate effective psycho-social support of people with lymphoedema. Secondly, there is still a perceived need to develop appropriate resources and clinics to treat and support patients with this condition (Williams et al., 2004).

The objective was to explore the journey of the service user through the pathway from diagnosis, to treatment, and to maintenance of condition. We present qualitative data from interviews, focus group discussions alongside questionnaire data on the themes of health care workers' knowledge of the condition, patients' views of their treatment, management of the condition, social support and other issues related to health care. Insight into barriers and facilitators for improving the experience of the care pathway are presented.

## **Methodology**

This study was approved by the University of Bradford Ethics committee.

### **Study Design**

This research is part of a larger scale mixed-methods study exploring appropriate screening tools for psychosocial difficulties resulting from SL and has been reported elsewhere (working papers, Dixey et al., 2012; Newell et al., 2012). The research presented here is from the qualitative component of the study, which included semi-structured interviews (phase 1), focus groups (phase 2) and free responses to postal questionnaires (phase 3). We utilised a participatory model (Cornwall and Jewkes, 1995) with service users as key members of the study steering group committee and who were, therefore central to the design process. Two co-researchers who were also service-users were recruited to the team to facilitate the focus group discussions. User groups, comprising members of the national and local patient support groups, Lymphoedema Support Network (LSN) and the Leeds Lymphoedema Link (LLL) agreed to discuss their experiences of lymphoedema and help pilot the focus groups. This informed the content of recruitment letters, information sheets, interview questions and questionnaire development. For added expert user engagement, the proposal received feedback from the Bradford Institute for Health Research Patient Panel.

### **Sample selection**

Purposive sampling was used in order to explore varied experiences from a breadth of age, gender, clinical histories and services accessed. The study drew a national sample. Inclusion criteria stated that the participant had to have lymphoedema secondary to cancer or its relevant treatment.

### **Participant Recruitment**

Initially, participants were recruited following an advertisement placed in the service-user newsletter of the Lymphoedema Support Network (LSN), *lymph line*. Word of mouth was also used, together with liaison with a local patient support group. A total of 172 responses were generated, and 142 took part in the study. Participants were selected to each phase of the study by using a screening question, 'Have you experienced difficulty socialising since you developed lymphoedema?' Those who answered 'yes' to the question were self-identified with psychosocial problems related to their SL and were invited to take part in a one-to-one interview, whereas those who did not were classified as 'reasonably adjusted to the condition' and were invited to attend focus group

discussions. The remaining volunteers were invited to complete a postal questionnaire.

### **Data Collection**

Fourteen women from around the UK were interviewed a further 15 took part in focus group discussions. A total of 104 responded to a postal questionnaire which involved completing validated questionnaires and two free response questions. At the beginning of the interviews and focus groups consent to participate and audiotape the sessions was sought. Any queries that arose were clarified at this stage. The interviews took approximately one hour each and were carried out in a private setting convenient to the respondent. Two focus groups were held (one in Leeds and one in London), facilitated by the research fellow and co-researcher/service user. Lunch was provided at both meetings and participants' travel expenses were paid. The focus groups were carried out using the same interview schedule from the interviews. The focus groups yielded information from 14 women and one man, we only report on the female perspective in this paper. A questionnaire pack was sent out to participants contributing to the postal survey, including a cover letter outlining the study and a stamped addressed return envelope. Non-respondents were sent a follow-up letter two weeks after the initial mailing. As standard, completion and return of the questionnaire was constituted consent to the study.

### **Data Analysis**

All interviews were transcribed verbatim using a contents analysis approach (Burnard (1991) pragmatic approach) to generate themes. This is a particularly well suited approach for exploring personal experiences where information is already known about elements of the phenomenon. Themes were extracted from the transcripts of the semi-structured interviews, focus group meetings and qualitative content of the free text responses of the postal survey. The interviewer generated codes and at least one other researcher carried out independent coding. Meetings were held to examine discrepancies and commonalities in coding and to organize low level codes into more general themes.



## Results

### Characteristics of Participants

The characteristics of the participants are summarised in table 1.

INCLUDE TABLE 1 HERE

### Study Findings

Transcripts yielded six themes: impact of lymphoedema; psychological implications and self-perceptions; other people's perceptions; clothing restrictions; physical limitations and the management of treatment. The themes are not presented in their entirety; rather, emphasis is placed on the management of lymphoedema, and those that relate to potential barriers and facilitators to service provision. Some reference to the impact of lymphoedema on the lived world of the person with SL is provided to contextualize the management of treatment from the initial diagnosis.

#### *Reaction to a diagnosis*

Women varied in their reaction to the diagnosis of SL. There was, however, a general consensus that it had a devastating impact on their lives with references made to loss of control or feelings of helplessness. The impact was overarching and affected many psychosocial domains: relation to a cancer diagnosis, self-esteem, relationships, work and treatment. Responses reflected major biographical disruption, this theme has been explored elsewhere (working paper, Dixey et al., 2012).

It affects every part of your life ....  
(SS: interview)

It's always something that I'm aware of ....  
(SJ: interview)

the first realisation that it was going to be permanent, just emotionally I just didn't want to go anywhere, I didn't want to do anything and I felt as though the person that I was before the breast cancer was gone ....  
(AW: interview)

These accounts highlight the profound impact that SL had on women and appeared universally presented even where a positive outlook was adopted. In addition to this, functional limitations that impacted on continuation of employment were highlighted.

### *Risk of acquiring lymphoedema*

Overall, there was a strong sense of anger towards the failure of healthcare professionals to explain the risk of SL. Alarming, some patients were completely unaware of the potential risk of SL whilst others refer to the inadequacy of information provided pre- and post-operatively. Consequently, women were often ill-prepared in identifying signs of developing lymphoedema and the diagnosis came as a shock.

I really think that if I'd been warned that it was a possibility... it was such a terrible shock.

(RP: interview)

Also I had not been warned of the possibility, I still feel great anger.  
(Questionnaire)

Some individuals referred to underestimating the significance of the condition, either when lymphoedema was listed on the consent form, or when information was provided in leaflet format post-operatively. One woman who was well-informed of the condition pre-operatively recounts how she was falsely reassured regarding her concerns of acquiring lymphoedema.

(...) I didn't know how it would affect me, so when I filled the form in consenting for my operation and they just go through the different things, lymphoedema, nobody actually explains to you what lymphoedema is...it's just lymphoedema...and sign here'.

(GF: interview)

I mentioned the day before my operation when I was told that they'd be removing my lymph nodes would I get lymphoedema? And the consultant said oh no, no, no that's very, very rare. You, you're I'm almost certain you won't get lymphoedema. Four days after my hysterectomy I was very poorly with an upset tummy and the nurse came to help me back into bed because I'd got drains and things attached. And she lifted the sheet and saw my legs and they're literally really swollen and said oh I think you've got lymphoedema...

(GB: interview)

Further, one woman with midline lymphoedema was warned about arm lymphoedema but not about the potential of lymphoedema elsewhere, '*the breast cancer nurses genuinely didn't know*' (PS: interview). This inevitably led to a delay in diagnosis. Women were often left with the impression that the SL could have been avoided if health care professionals were more aware of the risk, resulting in feelings of anger and distrust towards health care professionals.

I mean the doctor did say the reason I got it was the fact that they took blood from my arm...from this arm after I had cancer...and it was through that trauma that caused the lymphoedema...so yeah going into hospital

for a lumpectomy having an operation for something to...then coming out with something else that they've done wrong!

(S): interview)

### *Healthcare professionals' knowledge of lymphoedema*

Participants reported the medical profession's lack of knowledge, often referring to GPs as '*understanding nothing about lymphoedema*'. The majority of those who had suffered difficulties adapting to their condition referred to the dismissal of their concerns and lack of information and support given by their GPs, suggesting that there is an inequity in support for people living with SL.

...what I have found odd is from doctor's points of view...They don't know what it is...they know if you've got something else wrong with you. They don't know what lymphoedema is...

(SB: interview)

(...) even your GP said 'oh I don't know anything about lymphoedema you'd better ask your nurse'...

(RP: interview)

One respondent said that she was not told that the swelling was permanent, and had to probe for information in order to know the extent of her condition. The consensus was that there was a need for individuals to seek out their own information as the information provided by healthcare professionals was often lacking. Service users stated that they have to become experts in their own right, but this had resulted in barriers in their therapeutic relationship.

nobody, not the surgeons or the cancer nurse said this is something that you're going to have for life it was only when I went for my check-up that I said to him am I going to have this always?, can you cure it? He said I'm afraid so.

(RP: interview)

It's like most people with lymphoedema become experts because you go on the internet, because GPs know nothing... Even now, so you make sure that you're an expert...

(GB: interview)

### *Perceived Support*

Emphasis was placed on the importance of support both from healthcare professionals and socially. Many women referred to a lack of support among healthcare professionals and were often left with a sense of isolation and distress. This could arise from the lack of sensitivity over timing and language used when communicating an SL diagnosis. Women often felt dismissed or a nuisance and that their concerns were trivialised.

I rang the doctor on the Saturday morning and I said 'oh I think I've got lymphoedema' and he said 'well if you have then there's nothing you can do about it' and that was it...

(DN: interview)

(...) and I think the way that I was told when I was following the nurse down the corridor and she just sort of said it over her shoulder...

(SB: interview)

because when I was first diagnosed I was given my stockings, were prescribed for me and I had twenty minutes with somebody who showed me how to put it on...and they gave me the two stockings and that was the support I had for eleven years.

(Questionnaire)

Several people mentioned the isolating nature of lymphoedema, and that they had not been informed of support groups. The acceptability of support groups was mixed; some found them helpful and held them in positive regard. Conversely, one woman replied that she would not find a support group beneficial, but emphasised the importance of her social network. Women who mentioned the national support groups such as LSN spoke of them favourably.

I do find it's a very lonely thing to have lymphoedema...ermm I haven't, I don't know anybody else in the whole of [place name] and I have quite a lot of connections in [place name], I don't know anybody else who's got it but there must be people..

(RP: interview)

(... ) Very helpful...yes because they put me in touch with the lymphoedema nurse, yes I get the newsletters and a few times I've rung them for advice... But if they don't know they perhaps have the number of someone that you can contact... so they're really good

(GB: interview)

Where support was provided by healthcare professionals, it was seen as pivotal, able to '*change the outlook*' of a person after diagnosis, improve confidence and facilitate regaining of control.

'(...) and they [lymphoedema practitioner] were amazing, absolutely amazing, they explained things to me, they told me what the treatment was going to be. They said they can't cure me but they will keep it under control and that is precisely what they did...I think I began to have some

confidence when I could see that what they were doing was working and I became more confident again.'

(JE: interview)

### *Access to treatment and follow-up care*

Most participants referred to difficulties accessing initial treatment, or to follow-up treatment upon diagnosis, leading to '*frustration from lack of care*' and anger.

'But I felt very angry that I that I had to, that I had to sort of fight for treatment... And that shouldn't be...It should be, everybody should be entitled to the same treatment and education....

(DN: interview)

Further, available treatment focused on physical implications, ignoring psychological adjustment, which made women feel they were being troublesome or overly emotional:

'Because all they're looking at is the measurement of the arm...fitting you with the right sleeve, the right treatment? It's all very well and good but if you say I really hate this and it makes you feel so and so... it tends to be put to one side, this is my experience I'm not saying everybody's like that... so that does make you feel like you're making a fuss about nothing...'

(AW - interview)

The lack of funding for treatment, such as manual lymphatic drainage (MLD) and bandaging was continually raised. Many funded their own treatment and felt that they have to fight in order to be offered NHS treatment. One had self-funded after being told that her condition was only 'mild' whereas if it had been 'moderate' the NHS would have funded it, suggesting that preventative treatment is not a priority. However, many reported that active compression and massage treatment provided huge benefits, resulting in increased mobility and independence.

'the MLD is really, really good...my arm was much more flexible..... it just makes you feel as though you can do things.....Instead of having somebody else to do them for you...and it just makes you feel more useful to yourself...'

(BM - interview)

Most reported difficulties obtaining their compression garments, often with long delays from measurement to receiving the garment. This led to great frustration, with women chasing up their garment orders often to find that the order was incorrect or no longer appropriate.

'Yeah but now you can't even get your sleeve at the hospital...you have to get it on prescription and it takes you two months to get it...he writes the prescription ...I went, it actually took be six or seven weeks to get a sleeve this year... my local chemist wouldn't even order one...'

(SB: interview)

Often lymphoedema treatment was provided at a local hospice, and this was reported negatively. References were made to the connotations of the venue and the imagery this conjured up after surviving a former cancer diagnosis.

'...I go to the hospice and I think will I end up in here in a few years? When I'm seeing the consultant he was amazed I was still alive. I mean how long it will be before I'm in this hospice thank you very much...you know I'd rather go somewhere else...

(Focus group)

### *Use of compression garments*

Many of those that had been issued with compression garments had to acclimatise to managing them, including the effect it had on clothes-wearing, self-esteem, confidence and visibility. Garments were often covered up for fear of attracting attention, suggesting disability and inviting questions from others.

....if I'm dressed up it looks horrific wearing a sleeve, right it's not really nice. You can imagine yourself in a nice evening dress and a sleeve like this, it's not lovely is it... and you know so you decide right...well I do, if I'm going somewhere really posh I won't wear my sleeve. But then my arm's massive! So I have to compensate one thing for another thing ...'

(S): interview)

Despite discomfort about the garments' appearance, women were fully aware of their benefits; one woman said she '*feels unsafe without it*'. Interestingly, a woman with truncal lymphoedema said that the compression garment benefitted her body image:

so in a way... It's actually done my body image good... because I have these control vests on all the time...and now when I can so it does pull me in a bit... So it's a bit of a reverse body image....

(PC: interview)

### *Patient ideas for improved services*

Better training for healthcare workers on the condition itself, on more timely treatment and on the psychosocial needs of those affected was definitely a high priority:

‘I would like to think that both nurses and doctors and any staff dealing with people that are likely to have Lymphoedema are going to be educated, I feel extremely strongly about that.’

(JE, interview)

Timeliness was deemed a crucial variable for supportive interventions, particularly around the time of diagnosis. This has been mentioned in tandem with other themes and the importance of communication of a formal diagnosis.

‘Yeah so I think if you go right back to the beginning and look at how people are told about lymphoedema and how they’re treated when they do develop it I think you can help people a lot to come to terms with it.’

(AW: interview)

A strong theme emerging highlighted the importance of control in treatment and empowerment. Respondents referred to improvements in wellbeing and progression when accurate information was provided relating to their capacity to complete certain activities. Engagement with sporting activities was mentioned in the focus group attendees who were more accepting of the condition. This is contradictory to some of the responses where service users were apprehensive in engaging with certain activities and the fear of worsening symptomology.

(....) getting control about it has helped me because some time ago I didn’t know as much as I do now... and when it was out of control I found it difficult to cope with...because you do if something’s happening out of your control whether it’s the initial cancer or whatever...the more you get to know the more you’re in control...I feel in control of it now and I exercise a lot specifically for my lymphoedema different exercises to what I might have done at one time but I know that all of this if it’s cycling or Pilates classes or swimming every movement is doing my leg good and that makes me feel in control...and makes me feel you know good about things...and it has a spin off in doing my whole body good anyway but that helps me cope with it...

(Focus Group)

Reference to the benefit of supported self-care was mentioned and was derived as a positive element in mental wellbeing. Not surprisingly, a dedicated lymphoedema service was high on people’s priorities and when it was available was often transformative to patient benefit.

(.....) You can ring them at any time and they will get back to you immediately and they ring you to check-up, has everything settled down is everything alright? And you know I’ve said to them that makes a huge difference if you know that if you have any worries you can just ring them up the lymphoedema nurses and they’re there for you all the time. Whereas a few years ago they just weren’t there...

## Discussion

### Summary of the main findings

The study provides new insights into the experiences that women with SL have in accessing appropriate health care. It appears that there are a number of deficits: women are let down on their information needs, both about risk and management, staff involved through the cancer treatment pathway were often insensitive to the enormity of the diagnosis, professionals' knowledge of lymphoedema was insufficient, and treatment was patchy. More positively, dedicated lymphoedema services provided essential support and empowered women in their own treatment which promoted wellbeing. However, the majority of respondents reported the negative impact of a diagnosis on their mental wellbeing, and that psychosocial support was lacking in all or some of the stages of their journey with lymphoedema. It is clear that there are inequities in access to treatment with dedicated services only available in certain parts of the country. Thus, we have to conclude that our sample was reflective of the American study (Carter, 1997), "abandoned by medicine".

Despite the emphasis on better care for people with long term conditions, there are still some shortcomings. There is no explicit mention of psychosocial needs, for example in the Department of Health's list of goals in supporting people with long term conditions (DoH, 2005:7). Studies of other long term conditions, such as chronic bronchitis (Nicolson and Anderson, 2003) show that patients face psychological distress, negative self-esteem and disruption to relationships. Lymphoedema is not only a long term condition, but also a disfiguring one. Rumsey et al. (2004) in their studies of people with disfiguring body changes conclude that psychosocial needs and body image concerns are inadequately met, whilst the severity of the visible change is not necessarily associated with the level of distress (Rumsey et al., 2002). Existing literature on long term conditions and on disfiguring conditions affecting body image clearly shows that they impact negatively on mental wellbeing. It is therefore alarming that our data show such a failure to address women's potential distress. Not only was the diagnosis often communicated in an insensitive manner, but insufficiencies in knowledge of lymphoedema could lead to increased risk of it occurring.

Our data concur with McWayne and Heiney (2005) who found a continuing lack of understanding from hospital staff about lymphoedema. Our data show how devastating and life-changing lymphoedema is, something that healthcare professions do not appear to appreciate fully despite acknowledgements in other studies (Posner and Momenzadeh, 2004, Pyszel et al., 2006). Further, in the USA at least, Gordon and Mortimer (2007:741) assert, "Many patients currently receive inadequate care as lymphedema fails to receive appropriate recognition by the medical community." Our data would indicate that this occurs in the UK too. A recent study highlights an unmet education need for healthcare professions and provides insight into the extent of the perceived competency from the perspective of the health care professionals. Alarming, only 13% of



generalist healthcare practitioners felt that their education needs were adequately met and 69% of GPs identified difficulties in assessment of lymphoedema (Davies, 2011).

Rockson (2000:2816) argues that there is “a growing body of knowledge surrounding the important precipitating factors in the lymphedema that follows breast carcinoma therapy.” There is thus no excuse for not applying this knowledge in treatment regimes. Instead, women in our study felt that they had to become the experts, and that knowledge among GPs in particular was sadly deficient. Rockson (2000:2816) points to the failure to apply what is known: “Patients at risk and those with established disease await informed and sophisticated advice from their medical practitioners.” Evidence suggests that supported self-care empowers the service user to maintain their own wellbeing but relies heavily on expert guidance and support from healthcare professionals (Williams, 2011). An international consensus on best practice in lymphoedema management does exist (Framework, 2006) yet our work suggests that it is not applied uniformly.

There is no excuse for not knowing about lymphoedema, and also no excuse for poor communication skills. It was argued years ago that effective communication skills provide the key to good cancer care (Fallowfield and Jenkins, 1999, Moore et al., 2004). Indeed, they are key to good practice in all aspects of health care. It was clear and unsurprising that patients felt much happier and in control when they were taught about lymphoedema, and had supportive staff, usually in specialist lymphoedema services, providing MLD and compression garments. Programmes developed in other areas such as to help patients with disfigurement, including social interaction skills training, (e.g. (Robinson et al., 1996, Newell and Clarke, 2000, Rumsey and Harcourt, 2010) may also be effective for people with lymphoedema. Again, there is a need to apply what is already known.

### **Limitations of the study**

The study has a number of limitations. First, these reports are provided by service users and often relate to retrospective accounts, some from many years' experiences of healthcare which since then may have improved. Second, the data are reflective of a predominantly female sample and is not culturally diverse. The nature of the main study investigating social difficulties and appearance concerns may not be conducive to persuading men to volunteer. In addition, the majority of the participants had SL secondary to breast cancer, and it is natural that the sample is predominantly female. Caution is therefore required when interpreting how these data relate to present day service provisions and when generalising to the whole population.

### **Implications for policy, practice and future research**

Our research leads to a number of recommendations, which are outlined in table 2. They all relate to supported self-care in guiding the service user from a place of dependence to one of independence using a patient-centred approach.

INSERT TABLE 2 HERE

Our research has revealed a failure to translate existing evidence into practice and has also added patients' perspectives on how services could be improved. A subtheme arising predominantly in the focus groups detailed possible coping mechanisms which acted as facilitators for adjustment. These may be utilised to inform service improvement and relate to timeliness, communication and control through supported self-care. Recently, Todd et al (2010) developed a matrix designed to facilitate lymphoedema service development. This may prove a useful model in tailoring service provision to the patient's specific needs.

## **Conclusion**

Data from our sample of women with lymphoedema shows that their lives were dramatically changed by lymphoedema, but their information needs were not met and the risks of lymphoedema as a side effect of cancer treatment was not explained to them. They experienced healthcare professionals who were unaware of the condition or who expressed lack of knowledge. Their experiences with trying to access treatment were frustrating except where they had managed to find appropriate dedicated and specialist services. From the reports provided it appears that these are not available countrywide. Despite existing knowledge about the mental health impact of diagnosis with such a condition, appropriate psychosocial support is not forthcoming. Urgent attention to our recommendations would result in improved outcomes for this important group of patients.

## **Acknowledgements**

The authors would like to express gratitude to the people who responded to our advertisements and participated in the study, giving their time and insights with such care and detail. We would also thank Dr Gabrielle Tracy McClelland for help with research training and Julie Cottam, Ros Lam, the Lymphoedema Support Network (LSN) and Leeds Lymphoedema Link (LLL) for help with recruitment.

## **Funding**

This study was generously funded and supported by Macmillan Cancer Support.

## References

- Beaulac, S. M., McNair, L. A., Scott, T. E., Lamorte, W. W. & Kavanah, M. T. 2002. Lymphedema and quality of life in survivors of early-stage breast cancer. *Arch Surg*, 137, 1253-7.
- Browse, N. B., K.; Mortimer, P. 2003. *Diseases of the Lymphatics*, London, Arnold.
- Burnard, P. 1991. A method of analysing interview transcripts in qualitative research. *Nurse Educ Today*, 11, 461-6.
- Carter, B. J. 1997. Women's experiences of lymphedema. *Oncol Nurs Forum*, 24, 875-82.
- Clark, B., Sitzia, J. & Harlow, W. 2005. Incidence and risk of arm oedema following treatment for breast cancer: a three-year follow-up study. *QJM*, 98, 343-348.
- Cornwall, A. & Jewkes, R. 1995. What is participatory research? *Social science & medicine*, 41, 1667-1676.
- Davies, R. 2011. Education needs regarding lymphoedema: the specialists, the generalists and the differences of opinion. *Journal of Lymphoedema*, 6, 105.
- Dixey, R., Barlow, S., Todd, J., Taylor, V., Carney, S., Newell, R., "... emotionally I just didn't want to go anywhere...": a qualitative study of the impact of lymphoedema secondary to cancer on women's mental wellbeing. working paper, University of Bradford, Bradford. 17 October 2012.
- DOH: Department of Health 2005. The national service framework for long-term conditions. Department of Health London.
- Fallowfield, L. & Jenkins, V. 1999. Effective communication skills are the key to good cancer care. *Eur J Cancer*, 35, 1592-7.
- Fobair, P., Stewart, S. L., Chang, S., D'Onofrio, C., Banks, P. J. & Bloom, J. R. 2006. Body image and sexual problems in young women with breast cancer. *Psychooncology*, 15, 579-94.
- Framework, L. 2006. Best practice for the management of lymphoedema. *International consensus. London: MEP Ltd.*
- Fu, M. R. & Rosedale, M. 2009. Breast cancer survivors' experiences of lymphedema-related symptoms. *J Pain Symptom Manage*, 38, 849-59.
- Gordon, K. D. & Mortimer, P. S. 2007. A guide to lymphedema. *Expert Review of Dermatology*, 2, 741-752.
- Keeley, V. 2000. *Classification of lymphoedema*, Abingdon, Radcliffe medical Press Ltd

- McWayne, J. & Heiney, S. P. 2005. Psychologic and social sequelae of secondary lymphedema: a review. *Cancer*, 104, 457-66.
- Moffatt, C., Franks, P., Doherty, D., Williams, A., Badger, C., Jeffs, E., Bosanquet, N. & Mortimer, P. S. 2003. Lymphoedema: an underestimated health problem. *QJM*, 96, 731-738.
- Moore, P., Wilkinson, S. & Rivera Mercado, S. 2004. Communication skills training for health care professionals working with cancer patients, their families and/or carers. *Cochrane Database Syst Rev*, 2.
- Morgan, P. A., Franks, P. J. & Moffatt, C. J. 2005. Health-related quality of life with lymphoedema: a review of the literature. *Int Wound J*, 2, 47-62.
- Newell, R. & Clarke, M. 2000. Evaluation of a self-help leaflet in treatment of social difficulties following facial disfigurement. *Int J Nurs Stud*, 37, 381-8.
- Newell, R., Barlow, S., Dixey, R., Todd, J., Taylor, V., Carney, S., Validation of a lymphoedema specific scale for the assessment of Psychosocial difficulties, working paper, University of Bradford, Bradford, 17 October 2012.
- Nicholson, P. & Anderson, P. 2003. Quality of life, distress and self-esteem: a focus group study of people with chronic bronchitis. *Br J Health Psychol*, 8, 251-70.
- Passik, S. D. & McDonald, M. V. 1998. Psychosocial aspects of upper extremity lymphedema in women treated for breast carcinoma. *Cancer*, 83, 2817-20.
- Passick, S. D., Newman, M. L., Brennan, M. & Tunkel, R. 1995. Predictors of psychological distress, sexual dysfunction and physical functioning among women with upper extremity lymphedema related to breast cancer. *Psycho - Oncology*, 4, 255-263.
- Posner, T. N. & Momenzadeh, S. 2004. Swollen limbs: The impact on quality of life. *Australian Journal of Primary Health*, 10, 90-96.
- Pyszel, A., Malyszczak, K., Pyszel, K., Andrzejak, R. & Szuba, A. 2006. Disability, psychological distress and quality of life in breast cancer survivors with arm lymphedema. *Lymphology*, 39, 185-92.
- Ridner, S. H., Bonner, C. M., Deng, J. & Sinclair, V. G. 2012. Voices from the shadows: living with lymphedema. *Cancer Nurs*, 35, E18-26.
- Robinson, E., Rumsey, N. & Partridge, J. 1996. An evaluation of the impact of social interaction skills training for facially disfigured people. *Br J Plast Surg*, 49, 281-9.
- Rockson, S. G. 2000. Precipitating factors in lymphedema: myths and realities. *Cancer*, 83, 2814-2816.

- Rumsey, N., Clarke, A. & Musa, M. 2002. Altered body image: the psychosocial needs of patients. *Br J Community Nurs*, 7, 563-6.
- Rumsey, N., Clarke, A., White, P., Wyn-Williams, M. & Garlick, W. 2004. Altered body image: appearance-related concerns of people with visible disfigurement. *J Adv Nurs*, 48, 443-53.
- Rumsey, N. & Harcourt, D. 2010. Body image and congenital conditions resulting in visible difference.
- Ryan, M., Stainton, M. C., Jaconelli, C., Watts, S., Mackenzie, P. & Mansberg, T. 2003. The experience of lower limb lymphedema for women after treatment for gynecologic cancer. *Oncol Nurs Forum*, 30, 417-23.
- Tobin, M. B., Lacey, H. J., Meyer, L. & Mortimer, P. S. 1993. The psychological morbidity of breast cancer-related arm swelling. Psychological morbidity of lymphoedema. *Cancer*, 72, 3248-52.
- Todd, J. 2010. Achieving independence through self-management.
- Todd, J., Scally, A., Dodwell, D., Horgan, K. & Topping, A. 2008. A randomised controlled trial of two programmes of shoulder exercise following axillary node dissection for invasive breast cancer. *Physiotherapy*, 94, 265-273.
- Williams, A. 2011. *A qualitative study of supported self-care in women with lymphoedema associated with breast cancer*. Edinburgh Napier University.
- Williams, A. F., Moffatt, C. J. & Franks, P. J. 2004. A phenomenological study of the lived experiences of people with lymphoedema. *Int J Palliat Nurs*, 10, 279-86.
- Woods, M. 1993. Patients' perceptions of breast-cancer-related lymphoedema. *Eur J Cancer Care (Engl)*, 2, 125-8.

## **Figures**

**Table 1. Participant Characteristics**

Study Phase	Semi-Structured Interviews	Focus Groups	Postal Questionnaires
Participant Number	n = 14	n = 15	N = 104
Age Range	43-80 years	52-82 years	-
Mean age	62 years	65 years	-
Lymphoedema Location			
Upper Limb	8	10	64
Lower Limb	5	4	31
Truncal	1	-	3
Mixed	-	1	6
Time of Onset of Lymphoedema (months)			
Range	0-86	0-204	0-216
Median	5.5	12	6
Years since diagnosis (years)			
Range	2-30 years	2-30 years	1-47 years
Median	9 years	6.5 years	6 years
Marital Status			
Married/Living with Partner	9	14	-
Single and living alone:	2	1	-
Single and living with relatives:	3	-	-

**Table 2. Recommendations for Improving Service Provision**

- Comprehensive measures to raise awareness of lymphoedema and inform patients pre-operatively of the related risk. Awareness of the risk may aid in care strategies for prevention, although such strategies remain inconclusive.
- Raise awareness of lymphoedema amongst health care professionals.
- Improve access to manual lymphatic drainage services.
- Support coordinated care between multidisciplinary teams (i.e. oncology, general practice and lymphoedema departments). Include lymphoedema in individual care plans.
- Improvements in communication skills for health care professionals around diagnosis. The use of an empathic patient-centred approach is recommended in order to increase sensitivity to the emotional consequences of lymphoedema.
- Implement a stepped care model approach to psychosocial interventions for lymphoedema patients who present with psychosocial difficulties.
- Patients need to be guided to appropriate local and national support groups.
- Recognise the importance of compression garments. Improve availability and pathway to receiving compression garments. Efforts should be made in providing adequate supplies dependent on the individual's needs and in a timely manner.
- Access garments that may improve appearance. Provide choice for the colour of garment received. This may help in supporting women to feel more attractive.