



City Research Online

City, University of London Institutional Repository

Citation: Dimitrova, D. S., Ignatov, Z. G. & Kaishev, V. K. (2017). On the First Crossing of Two Boundaries by an Order Statistics Risk Process. *Risks*, 5(3), 43. doi: 10.3390/risks5030043

This is the published version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/18049/>

Link to published version: <https://doi.org/10.3390/risks5030043>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

Article

On the First Crossing of Two Boundaries by an Order Statistics Risk Process

Dimitrina S. Dimitrova ^{1,*}, Zvetan G. Ignatov ² and Vladimir K. Kaishev ¹

¹ Faculty of Actuarial Science and Insurance, Cass Business School, City, University of London, 106 Bunhill Row, London EC1Y 8TZ, UK; v.kaishev@city.ac.uk

² Faculty of Economics and Business Administration, Sofia University “St Kliment Ohridski”, 125 Tsarigradsko Shosse Blv., bl.3, Sofia 1113, Bulgaria; ignatov@feb.uni-sofia.bg

* Correspondence: D.Dimitrova@city.ac.uk; Tel.: +44-207-040-8470

Academic Editor: Mogens Steffensen

Received: 14 July 2017; Accepted: 15 August 2017; Published: 18 August 2017

Abstract: We derive a closed form expression for the probability that a non-decreasing, pure jump stochastic risk process with the order statistics (OS) property will not exit the strip between two non-decreasing, possibly discontinuous, time-dependent boundaries, within a finite time interval. The result yields new expressions for the ruin probability in the insurance and the dual risk models with dependence between the claim severities or capital gains respectively.

Keywords: double boundary non-crossing probability; point process; risk process; ruin probability; Appell polynomials

1. Introduction

The probability that a stochastic process stays between two boundaries is of significant importance in many areas among which: statistics, for constructing confidence intervals for distribution functions, (see [Wald and Wolfowitz 1939](#); [Steck 1971](#)) or in sequential analysis; finance, for pricing double-barrier options, (see [Borovkov and Novikov 2005](#)); actuarial science, in modelling the surplus of an insurance company (see [Teunen and Goovaerts 1994](#)), and also in probability (see e.g., [Potzelberger and Wang 2001](#); [Lotov 1996](#); [Buonocore et al. 1990](#)) and other areas.

The double-boundary crossing problem we are concerned with in this paper is as follows. We consider a stochastic process S_t , constructed by the two independent sequences of random variables $0 \leq T_1 \leq T_2 \leq \dots$ and $0 < Y_1 < Y_2 < \dots$, with the assumption that at each instant T_i the process jumps at the level Y_i , $i = 1, 2, \dots$. It is assumed that the instants T_1, T_2, \dots , model the arrival of certain events, e.g., insurance claims, or capital gains or some other events in a particular application. The jumps, $0 < Y_1 < Y_2 < \dots$, could be interpreted as reflecting the current level of some process of interest, e.g., the aggregate claims to an insurance company, or the accumulation of reserves of a bank. It is further assumed that the instants, T_1, T_2, \dots form a point process ξ , on $[0, \infty)$, with a cumulative intensity function $\Lambda([0, z]) = \Lambda(z) = E\xi[0, z] < \infty$, such that, $\Lambda(z) > 0, \forall z \in (0, \infty)$, $\Lambda(0) \geq 0$, where $\xi[0, z]$ denotes the number of arrivals in $[0, z]$.

In summary, we consider a pure-jump stochastic process $S_t = Y_{\xi[0,t]}$ with right-continuous, non-decreasing trajectories and are interested in the probability, $P(g(t) \leq S_t \leq h(t), 0 \leq t \leq z)$ that within a fixed time interval, $[0, z]$, the process S_t stays between an upper and a lower deterministic, non-decreasing time-dependent boundaries, $h(t)$ and $g(t)$. The interpretation of the boundaries is also dependent on the application. For example, $h(t)$ and $g(t)$ could model the cumulative insurance premium income and expense outgo, respectively in the insurance and dual risk models considered in the context of ruin theory. Or these could be interpreted as correspondingly, the maximum and minimum recommended bounds for a bank's loan-to-deposit ratio so that the bank maximizes its

returns while at the same time remains liquid. Formal specification of the boundaries, $h(t)$ and $g(t)$, is provided in Section 2.

We shall make the following assumptions regarding the (risk) process $S_t = Y_{\xi[0,t]}$. It will be assumed that the point process, ξ , is a process with the so called *order statistics* (OS) property, or simply OS point process. The latter is characterized by the property that, given j arrivals in a finite interval $[0, z]$, $z > 0$, the successive arrival times coincide in distribution with the order statistics of j independent and identically distributed random variables with a cumulative distribution function, $F_z(t)$, $0 \leq t \leq z$, $F_z(0) = 0$ and $F_z(z) = 1$.

OS processes with unit jumps at the arrival times $0 < T_1 < T_2 < \dots < T_j$, i.e., for which $F_z(t)$ is assumed continuous, have been thoroughly studied (see e.g., Nawrotzki 1962; Crump 1975; Feigin 1979; Puri 1982) and have also been applied in risk and ruin theory by (Willmot 1989; De Vylder and Goovaerts 1999, 2000; Lefèvre and Picard 2011, 2014; Sendova and Zitikis 2012), to model insurance claim arrivals. Such OS processes are Markovian (see e.g., Thompson 1988) and include the (mixed) Poisson process, the Pólya-Lundberg process, and the linear birth process with immigration, equivalent to negative binomial processes and the linear death process (see e.g., Crump (1975), and more recently Goffard and Lefèvre (2017) for details).

As noted by Lefèvre and Picard (2011), OS processes are particularly suitable for modelling claim arrivals with dependence as they capture situations where the arrival of a claim increases the likelihood of more claim-arrivals leading to non-overlapping clustering, see e.g., the birth process considered in Bühlmann (1970). In contrast, in a death point process the arrival of a claim decreases the likelihood of arrival of more claims, as is the case in modelling the arrival of claims from a group life policy covering a closed group of individuals. It should also be mentioned that OS processes are appealing in modelling population growth where (OS) birth and death processes are a key modelling tool (see e.g., Kendall. 1949; Haccou et al. 2005).

Recently, Dimitrova et al. (2014) considered more general OS point processes by allowing $\Lambda(t)$ to be discontinuous at some fixed instants, i.e, $F_z(t)$ to be discontinuous at these instants, since as shown by Crump (1975), for an OS process, $F_z(t) = \Lambda(t) / \Lambda(z)$. This extends further the flexibility of the OS risk process, S_t , allowing risk events to arrive not only at random moments but also at fixed instants with non-zero probability, possibly forming clusters. This is an appealing feature, e.g., in life and non-life insurance applications, see Section 4 and Dimitrova et al. (2014) for particular examples in the special case of ruin by exceeding an upper boundary.

Our aim in the present paper is to derive, under this more general definition of an OS risk process, S_t , closed-form expressions for the non-exit probability $P(g(t) \leq S_t \leq h(t), 0 \leq t \leq z)$, under reasonably general assumptions on the boundaries, $g(t)$ and $h(t)$, allowing them to have jump discontinuities (see Theorem 1). Its proof is based on extending two results of Stek (1971), which give the probability that the order statistics from a sample of j uniform r.v.s all lie in a j -dimensional rectangle. We show also how Theorem 1 directly yields new ruin probability formulas in the insurance and the dual risk models under reasonably general assumptions, which have recently attracted considerable amount of research (see e.g., Ignatov and Kaishev 2000, 2004; Lefèvre and Picard 2011, 2014; Dimitrova et al. (2015)). We further note that our paper (see also an earlier version Dimitrova et al. (2015)) covers and extends the non-crossing probability results for parallel boundaries of Xu (2012), Goffard and Lefèvre (2017), Goffard (2017).

The paper is organized as follows. In Section 2, we prove our main result given by Theorem 1 and its discrete version, (see Corollary 2). In Section 3 we give Corollaries 3, 4, which establish new ruin probability formulas for the two important special cases, the insurance and the dual risk models with dependence between the claim severities or capital gains respectively. In Section 4, we consider the numerical implementation of the results of Theorem 1 and Corollary 2 and illustrate these results for some special cases of OS claim arrival processes, (see 1 and 2).

2. A Formula for the Non-Exit Probability $P(g(t) \leq S_t \leq h(t), 0 \leq t \leq z)$

We assume that the two functions $h(t)$ and $g(t)$, $t \in [0, +\infty)$, have the following properties: $h(t)$ and $g(t)$ are non-decreasing functions, such that $h(0) \geq 0$ and $g(0) \leq 0$; $h(t) \geq g(t)$, $\forall t \in [0, +\infty)$; $h(t)$ and $g(t)$ may have jump-discontinuities, and we assume that $h(t)$ is right-continuous and $g(t)$ is left-continuous. We will define the inverse functions, $h^{-1}(y) = \inf\{t : h(t) \geq y\}$, $g^{-1}(y) = \sup\{t : g(t) \leq y\}$, $y \in [0, +\infty)$, so that $h^{-1}(y)$ is left-continuous and $g^{-1}(y)$ is right-continuous. We will further consider restrictions of $h(t)$ and $g(t)$ on $[0, z]$, $z > 0$, denoted by $h_z(t)$ and $g_z(t)$, and define the corresponding restrictions of the inverse functions as $h_z^{-1}(y) = \min(z, h^{-1}(y))$ and $g_z^{-1}(y) = \min(z, g^{-1}(y))$. In view of these definitions, we will assume that the process S_t does not exit if its trajectory touches either one of the two boundaries.

As noted, here we will adopt the more general definition of an OS arrival process, recently considered by [Dimitrova et al. \(2014\)](#). We assume that, ξ has an arbitrary cumulative intensity function $\Lambda(z)$, such that $\Lambda(z) > 0, z > 0, \Lambda(0) \geq 0$, and cumulative distribution function

$$F_z(t) = \begin{cases} 0 & \text{if } t < 0 \\ \Lambda(t)/\Lambda(z) & \text{if } 0 \leq t < z \\ 1 & \text{if } t \geq z, \end{cases} \tag{1}$$

where $z > 0$. If X_1, \dots, X_j are independent and identically distributed random variables with distribution function $F_z(t)$ and if, $X_{1,j}, \dots, X_{j,j}$ denote their order statistics, then the process ξ is said to have the OS property if, given j arrivals in a finite interval $[0, z]$, (i.e., given $\xi[0, z] = j$), $j = 1, 2, \dots$, the successive points, $0 \leq T_1 \leq T_2 \leq \dots \leq T_j$ of the process ξ , have a conditional joint distribution which coincides with the joint distribution of the order statistics $X_{1,j}, \dots, X_{j,j}$.

From the definition of $F_z(t)$ it can be seen that it is right-continuous and hence, $F_z(t) = F_z(t+)$, with possible jumps, $F_z(t) - F_z(t-) \neq 0$, where $F_z(t+) = \lim_{s \downarrow t} F_z(s)$ and $F_z(t-) = \lim_{s \uparrow t} F_z(s)$. Let $0 \equiv y_0 < y_1 < \dots < y_j < \infty$ be a realization of the sequence $0 < Y_1 < Y_2 < \dots < Y_j$, with joint cdf, $F(y_1, \dots, y_j)$. We can now state our main result.

Theorem 1. *The non-exit probability $P(g_z(t) \leq S_t \leq h_z(t), 0 \leq t \leq z)$, $0 < z < \infty$, for an OS arrival process, ξ with an arbitrary distribution function, $F_z(t)$ defined as in (1), is given by:*

$$P(g_z(t) \leq S_t \leq h_z(t), 0 \leq t \leq z) = \sum_{j=0}^{\infty} j! P(\xi[0, z] = j) \int \dots \int_{C_j} \mathbf{1}_{g_z(z)}(y_j) \det \left(\left(\delta_{m,l}^{(j)} \right)_{1 \leq m,l \leq j} \right) dF(y_1, \dots, y_j), \tag{2}$$

where $C_j = \{(y_1, \dots, y_j) : 0 < y_1 < \dots < y_j < h_z(z)\}$, $\mathbf{1}_{g_z(z)}(y_j) = \begin{cases} 1 & \text{if } y_j \geq g_z(z) \\ 0 & \text{if } y_j < g_z(z) \end{cases}$,

and $\det \left(\left(\delta_{m,l}^{(j)} \right)_{1 \leq m,l \leq j} \right)$ is the determinant of an $j \times j$ matrix with elements:

$$\delta_{m,l}^{(j)} = \frac{[F_z(g_z^{-1}(y_{m-1})) - F_z(h_z^{-1}(y_l) -)]_+^{l-m+1}}{(l-m+1)!}, \tag{3}$$

$1 \leq m, l \leq j$, where $[x]_+ = \max(0, x)$, $\delta_{m,l}^{(j)} = \begin{cases} 0 & \text{if } l - m + 1 < 0 \\ 1 & \text{if } l - m + 1 = 0 \end{cases}$, and where for $j = 0$ we assume that the multiple integral in (2) is equal to 0 if $g_z^{-1}(0) < z$ or is equal to 1 if $g_z^{-1}(0) = z$. Alternatively,

$$\begin{aligned}
 &P(g_z(t) \leq S_t \leq h_z(t), 0 \leq t \leq z) \\
 &= \sum_{j=0}^{\infty} P(\xi[0, z] = j) \int \dots \int_{C_j} \mathbf{1}_{g_z(z)}(y_j) \det \left(\left(\rho_{m,l}^{(j)} \right)_{1 \leq m, l \leq j} \right) dF(y_1, \dots, y_j), \tag{4}
 \end{aligned}$$

where

$$\rho_{m,l}^{(j)} = \binom{l}{l-m+1} \left[F_z \left(g_z^{-1}(y_{m-1}) \right) - F_z \left(h_z^{-1}(y_l) \right) \right]_+^{l-m+1}, \tag{5}$$

for $1 \leq m, l \leq j$ and $\rho_{m,l}^{(j)} = \begin{cases} 0 & \text{if } l - m + 1 < 0 \\ 1 & \text{if } l - m + 1 = 0 \end{cases}$, and where for $j = 0$ we assume that the multiple integral in (4) is equal to 0 if $g_z^{-1}(0) < z$ or is equal to 1 if $g_z^{-1}(0) = z$.

Remark 1. From (3) and (5), it follows that $\left(\delta_{m,l}^{(j)} \right)_{1 \leq m, l \leq j}$ and $\left(\rho_{m,l}^{(j)} \right)_{1 \leq m, l \leq j}$ are lower Hessenberg matrix. A matrix whose elements above or below the first subdiagonal are equal to zero (i.e., all elements, $a_{ij} = 0$ if $j - i > 1$ or if $i - j > 1$) are called Hessenberg matrixes. The determinants of the latter are computed using a straightforward recurrence relation established by Lemma 2. For properties of Hessenberg matrixes and their determinants we refer to, e.g., [Vein and Dale \(1999\)](#).

Since Appell polynomials and other types of Appell structures are also expressed as Hessenberg determinants (see e.g., [Vein and Dale \(1999\)](#), [Ignatov and Kaishev 2000, 2016](#), and [Dimitrova et al. \(2014\)](#)), here we will interchangeably refer to $\det \left(\left(\delta_{m,l}^{(j)} \right)_{1 \leq m, l \leq j} \right)$ and $\det \left(\left(\rho_{m,l}^{(j)} \right)_{1 \leq m, l \leq j} \right)$ as Appell-Hessenberg functions.

In order to prove Theorem 1, we will need an auxiliary result, stated as Lemma 1 bellow. Let X_1, \dots, X_j be independent and identically distributed random variables with distribution function $F_z(t)$, $X_{1,j}, \dots, X_{j,j}$ denote their order statistics and let, U_1, \dots, U_j be independent uniformly on $[0, 1]$ distributed random variables with order statistics, $U_{1,j}, \dots, U_{j,j}$. Let us also define

$$F_z^{-1}(u) = \inf\{t : F_z(t) \geq u\}, \quad 0 \leq u \leq 1. \tag{6}$$

Lemma 1. If $F_z(t)$ is a right continuous cdf then, for $\alpha_1, \dots, \alpha_j$ and β_1, \dots, β_j such that $0 \leq \alpha_i < \beta_i \leq z$, $i = 1, \dots, j$ and j , a natural number, we have

$$\begin{aligned}
 &P \left(\bigcap_{i=1}^j (\alpha_i \leq X_{i,j} \leq \beta_i) \right) \\
 &= P \left(\bigcap_{i=1}^j (\alpha_i \leq F_z^{-1}(U_{i,j}) \leq \beta_i) \right) = P \left(\bigcap_{i=1}^j (F_z(\alpha_i-) \leq U_{i,j} \leq F_z(\beta_i)) \right). \tag{7}
 \end{aligned}$$

For a proof of Lemma 1, see Appendix A.

Let us introduce the shorter notation $d_j = P \left(\bigcap_{i=1}^j (\alpha_i \leq X_{i,j} \leq \beta_i) \right)$. The following Lemma gives a recurrence expression for the probability d_j , obtained by [Epanechnikov \(1968\)](#), which directly follows, expanding the determinant in (11), with respect to the last column, with $u_i = F_z(\alpha_i-)$ and $v_i = F_z(\beta_i)$.

Lemma 2. For $\alpha_1, \dots, \alpha_j$ and β_1, \dots, β_j such that $0 \leq \alpha_i < \beta_i \leq z, i = 1, \dots, j$ and $\alpha_1 \leq \dots \leq \alpha_j, \beta_1 \leq \dots \leq \beta_j$, we have

$$d_k = - \sum_{i=0}^{k-1} \binom{k}{i} \left(F_z(\beta_{i+1}) - F_z(\alpha_{k-i}) \right)_+^{k-i} (-1)^{k-i} d_i, \quad k = 1, \dots, j, \tag{8}$$

where $d_0 \equiv 1$.

We will also need the following two results, due to [Steck \(1971\)](#).

Theorem 2. (G.P. Steck). Let u_1, \dots, u_j and v_1, \dots, v_j be such that $0 \leq u_i < v_i \leq 1, i = 1, \dots, j, u_1 \leq \dots \leq u_j, v_1 \leq \dots \leq v_j$. Then,

$$P \left(\bigcap_{i=1}^j (u_i \leq U_{i,j} \leq v_i) \right) = j! \det \left(\left(\tilde{\delta}_{m,l}^{(j)} \right)_{1 \leq m,l \leq j} \right), \tag{9}$$

where $\det \left(\left(\tilde{\delta}_{m,l}^{(j)} \right)_{1 \leq m,l \leq j} \right)$ is the determinant of an $j \times j$ matrix with element on the m -th row and the l -th column defined as:

$$\tilde{\delta}_{m,l}^{(j)} = \begin{cases} (v_m - u_l)_+^{l-m+1} / (l - m + 1)! & \text{if } l - m + 1 \geq 0, \\ 0 & \text{if } l - m + 1 < 0. \end{cases} \tag{10}$$

Corollary 1. (G.P. Steck). Under the conditions of [Theorem 2](#), we have

$$P \left(\bigcap_{i=1}^j (u_i \leq U_{i,j} \leq v_i) \right) = \det \left(\left(\tilde{\rho}_{m,l}^{(j)} \right)_{1 \leq m,l \leq j} \right), \tag{11}$$

where

$$\tilde{\rho}_{m,l}^{(j)} = \binom{l}{l - m + 1} (v_m - u_l)_+^{l-m+1}. \tag{12}$$

We are now ready to prove [Theorem 1](#).

Proof of Theorem 1. Recall that, since $g_z(t)$ is restricted to the interval $[0, z]$ and $g_z(0) \leq 0$, two cases need to be considered, $g_z^{-1}(0) < z$ and $g_z^{-1}(0) = z$. In the case of $g_z^{-1}(0) < z$, applying the formula of total probability, we have

$$\begin{aligned} & P(g_z(t) \leq S_t \leq h_z(t), 0 \leq t \leq z) \\ &= \sum_{j=1}^{\infty} P(\xi[0, z] = j) P(g_z(t) \leq S_t \leq h_z(t), 0 \leq t \leq z \mid \xi[0, z] = j), \end{aligned} \tag{13}$$

where the summation starts from, $j = 1$, since for $j = 0$ the conditional probability is 0 when $g_z^{-1}(0) < z$. For the conditional probability on the right-hand side of (13), we have

$$\begin{aligned}
 &P(g_z(t) \leq S_t \leq h_z(t), 0 \leq t \leq z \mid \xi[0, z] = j) \\
 &= P\left((h_z(z) \geq Y_j \geq g_z(z)) \bigcap_{i=1}^j \left((h_z^{-1}(Y_i) -) \leq T_i \leq g_z^{-1}(Y_{i-1}) \right) \right) \\
 &= \int \dots \int_{C_j} \mathbb{1}_{g_z(z)}(y_j) P\left(\bigcap_{i=1}^j \left((h_z^{-1}(y_i) -) \leq T_i \leq g_z^{-1}(y_{i-1}) \right) \right) dF(y_1, \dots, y_j) \\
 &= \int \dots \int_{C_j} \mathbb{1}_{g_z(z)}(y_j) P\left(\bigcap_{i=1}^j \left(F_z(h_z^{-1}(y_i) -) \leq U_{i,j} \leq F_z(g_z^{-1}(y_{i-1})) \right) \right) dF(y_1, \dots, y_j),
 \end{aligned} \tag{14}$$

where the last equality follows from Lemma 1, with $\alpha_i = (h_z^{-1}(y_i) -)$, $\beta_i = g_z^{-1}(y_{i-1})$, noting that, T_1, \dots, T_j coincide in distribution with the order statistics, X_{1j}, \dots, X_{jj} . The asserted equalities (2) and (4) now follow substituting, $P(g_z(t) \leq S_t \leq h_z(t), 0 \leq t \leq z \mid \xi[0, z] = j)$, expressed by the last equality in (14), back in (13) and applying Theorem 2 and Corollary 1 to express $P\left(\bigcap_{i=1}^j (F_z(h_z^{-1}(y_i) -) \leq U_{i,j} \leq F_z(g_z^{-1}(y_{i-1}))) \right)$, with $F_z(h_z^{-1}(y_i) -) = u_i$ and $F_z(g_z^{-1}(y_{i-1})) = v_i$.

In the case of $g_z^{-1}(0) = z$, for $j = 0$, the conditional probability is 1 and the term $P(\xi[0, z] = 0)$ should be added to the sum on the right-hand side of (13). This completes the proof of the assertion of Theorem 1. \square

It can directly be seen that Theorem 1 is also valid for discrete Y_1, Y_2, \dots in which case multiple integration is replaced by appropriate multiple summation, and (2) and (4) become exact, closed form expressions more appealing for numerical implementation. This is illustrated by Corollary 2 given next, where formula (15) is a discrete version of (4).

Corollary 2. *The non-exit probability $P(g_z(t) \leq S_t \leq h_z(t), 0 \leq t \leq z)$, $0 < z < \infty$, for an OS arrival process, ξ , with an arbitrary distribution function, $F_z(t)$ defined as in (1), and arbitrarily distributed (possibly dependent) discrete integer-valued Y_1, Y_2, \dots with joint probability mass function $P(Y_1 = y_1, \dots, Y_j = y_j) = p(y_1, \dots, y_j)$, is given by:*

$$\begin{aligned}
 &P(g_z(t) \leq S_t \leq h_z(t), 0 \leq t \leq z) \\
 &= \sum_{j=0}^{\lfloor h_z(z) \rfloor} P(\xi[0, z] = j) \sum_{\substack{1 \leq y_1 < y_2 < \dots < y_j \leq \lfloor h_z(z) \rfloor \\ \lceil g_z(z) \rceil \leq y_j}} \det\left(\left(\rho_{m,l}^{(j)} \right)_{1 \leq m, l \leq j} \right) p(y_1, \dots, y_j), \\
 &= \sum_{j=0}^{\lfloor h_z(z) \rfloor} P(\xi[0, z] = j) \sum_{y_1=1}^{\lfloor h_z(z) \rfloor - (j-1)} \sum_{y_2=y_1+1}^{\lfloor h_z(z) \rfloor - (j-2)} \dots \sum_{y_{j-1}=y_{j-2}+1}^{\lfloor h_z(z) \rfloor - 1} \sum_{y_j=\max(y_{j-1}, \lceil g_z(z) \rceil)}^{\lfloor h_z(z) \rfloor} \\
 &\det\left(\left(\rho_{m,l}^{(j)} \right)_{1 \leq m, l \leq j} \right) \times p(y_1, \dots, y_j),
 \end{aligned} \tag{15}$$

where $\lfloor h_z(\cdot) \rfloor$ is the integer part of $h_z(\cdot)$, $\lceil g_z(\cdot) \rceil$ is the least integer $\geq g_z(\cdot)$, $\det\left(\left(\rho_{m,l}^{(j)} \right)_{1 \leq m, l \leq j} \right)$ is the determinant of an $j \times j$ matrix with elements, $\rho_{m,l}^{(j)}$ defined as in (5) and where for $j = 0$ we assume that the multiple summation in (15) is equal to 0 if $g_z^{-1}(0) < z$ or is equal to 1 if $g_z^{-1}(0) = z$.

Proof of Corollary 2. The proof directly follows the lines of the proof of Theorem 1, replacing multiple integration with the corresponding multiple summation. \square

3. Application to Ruin Probability

In this section we will consider two corollaries of Theorem 1, which yield new explicit formulas for the probability of non-ruin in the insurance and dual risk models.

3.1. Ruin in the Insurance Risk Model

Ruin under the insurance risk model has been thoroughly investigated, but very few closed form expressions have been established. See e.g., Picard and Lefèvre (1997), Ignatov and Kaishev 2000, 2004; Ignatov et al. (2001); Lefèvre and Loisel (2009) and Dimitrova et al. (2016) for finite-time ruin formulas under Poisson claim arrivals, involving Appell polynomials. Ruin formulas under more general arrival processes have been recently established: by Ignatov and Kaishev (2016) for the case when the claim arrival process ξ is a process with independent increments; and by Lefèvre and Picard 2011, 2014 and Dimitrova et al. (2014) for the case when ξ is a process with the OS property.

In the insurance risk model, the upper boundary, $h(t)$ models the cumulative premium income of an insurance company up to time t , and $S_t = Y_{\xi[0,t]}$ is interpreted as its aggregate claim amount process with partial sums of the individual claims, Y_1, Y_2, \dots , arriving at the instants, T_1, T_2, \dots . Denote by $R_t = h(t) - S_t$ the insurance company's risk process and define the instant of ruin, T as $T := \inf \{t : 0 \leq t \leq z, R_t < 0\}$ or $T = \infty$ if $R_t \geq 0$ for all $0 \leq t \leq z$. It is not difficult to see that the probability of non-ruin in $[0, z]$, $P(T > z) = P(0 \leq R_t, 0 \leq t \leq z) = P(S_t \leq h(t), 0 \leq t \leq z)$ coincides with the non-exit probability $P(g_z(t) \leq S_t \leq h_z(t), 0 \leq t \leq z)$, where the lower boundary, $g_z(t) = 0$ for every $t \in [0, z]$, i.e., is such that, $g_z^{-1}(y) \equiv z, 0 \leq y < \infty$. We can now give the following.

Corollary 3. The finite-time non-ruin probability $P(T > z) = P(g_z(t) \leq S_t \leq h_z(t), 0 \leq t \leq z) = P(S_t \leq h_z(t), 0 \leq t \leq z)$, $0 < z < \infty$, for an OS claim arrival process, ξ with an arbitrary distribution function, $F_z(t)$, defined as in (1), is given by (2) with (3) replaced by:

$$\delta_{m,l}^{(j)} = \frac{[1 - F_z(h_z^{-1}(y_l) -)]^{l-m+1}}{(l - m + 1)!}, \quad 1 \leq m, l \leq j, \tag{16}$$

where $(l - m + 1)! = 0$ if $l - m + 1 < 0$; or as (4) with (5) replaced by

$$\rho_{m,l}^{(j)} = \binom{l}{l - m + 1} [1 - F_z(h_z^{-1}(y_l) -)]^{l-m+1}, \quad 1 \leq m, l \leq j, \tag{17}$$

where $\binom{l}{l - m + 1} \equiv 0$ if $l - m + 1 < 0$.

Proof of Corollary 3. The asserted equalities (16) and (17) follow from (3) and (5) noting that since, $g_z^{-1}(y) \equiv z, 0 \leq y < \infty$, then $F_z(g_z^{-1}(y_{m-1})) = F_z(z) = 1, 1 \leq m \leq j$. \square

3.2. Ruin in the Dual Risk Model

The dual risk model has been intensively studied in the ruin probability literature which is summarized in Dimitrova et al. (2015), where an instructive link between the insurance and the dual risk models is established and exploited to obtain new ruin formulas, for the dual model.

In the dual model, the lower boundary, $g(t)$ models the cumulative expense outgo of a company up to time t , and $S_t = Y_{\xi[0,t]}$ is interpreted as its aggregate capital gain process with partial sums of the individual capital gains, Y_1, Y_2, \dots , arriving at the instants, T_1, T_2, \dots . Such companies could typically, be mineral exploration companies or companies specializing in research and development, where capital gains occur at random instants of e.g., mineral finds or technological breakthroughs.

It is not difficult to see that the probability of non-ruin of the company, $P(g(t) \leq S_t, 0 \leq t \leq z)$ coincides with the non-exit probability $P(g_z(t) \leq S_t \leq h_z(t), 0 \leq t \leq z)$, where the upper boundary, $h_z(t) = +\infty$ for every $t \in [0, z]$, i.e., is such that, $h_z^{-1}(y) \equiv 0, 0 \leq y \leq \infty$. We can now give the following.

Corollary 4. *The finite-time non-ruin probability $P(g_z(t) \leq S_t \leq h_z(t), 0 \leq t \leq z) = P(g_z(t) \leq S_t, 0 \leq t \leq z), 0 < z < \infty$, for an OS claim arrival process, ξ with an arbitrary distribution function, $F_z(t)$, defined as in (1), is given by (2) with (3) replaced by:*

$$\delta_{m,l}^{(j)} = \frac{[F_z(g_z^{-1}(y_{m-1}))]^{l-m+1}}{(l-m+1)!}, \quad 1 \leq m, l \leq j, \tag{18}$$

where $(l-m+1)! = 0$ if $l-m+1 < 0$; or as (4) with (5) replaced by

$$\rho_{m,l}^{(j)} = \binom{l}{l-m+1} [F_z(g_z^{-1}(y_{m-1}))]^{l-m+1}, \quad 1 \leq m, l \leq j, \tag{19}$$

where $\binom{l}{l-m+1} \equiv 0$ if $l-m+1 < 0$.

Proof of Corollary 4. The asserted equalities (18) and (19) follow from (3) and (5) noting that since, $h_z^{-1}(y) \equiv 0, 0 \leq y \leq \infty$, then $F_z(h_z^{-1}(y_l) -) = F_z(0-) = 0, 1 \leq l \leq j$. \square

4. Numerical Considerations and Examples

In this section, we give details on how our main result established by Theorem 1 can be implemented numerically and illustrate it based on two particular examples of OS point processes modelling insurance claim arrivals.

It should be noted that for the efficient evaluation of $P(g_z(t) \leq S_t \leq h_z(t), 0 \leq t \leq z)$, following (2), or (4), it is essential to be able to:

1. appropriately truncate the infinite summation;
2. compute the underlying (multiple) j -dimensional integrals, for possibly large values of j ;
3. efficiently compute the integrand functions $\det\left(\left(\delta_{m,l}^{(j)}\right)_{1 \leq m, l \leq j}\right)$, or equivalently $\det\left(\left(\rho_{m,l}^{(j)}\right)_{1 \leq m, l \leq j}\right)$.

We will address items 1–3 as follows.

1. It should be mentioned that explicit expressions for the probability $P(\xi[0, z] = j)$ exist for the well-established OS point processes in the literature (see e.g., the four cases discussed in Goffard and Lefèvre (2017)), and it can directly be substituted in either (2), or (4). This allows for truncating the infinite sum with respect to j up to a maximum value say j_ϵ^* above which the probability $\sum_{j_\epsilon^*+1}^\infty P(\xi[0, z] = j) < \epsilon$ is negligibly small leading to negligibly small sum of the product terms in (13)), hence in (2), or (4). Similar approach has been exploited in Section 4 of Dimitrova et al. (2016) for the Poisson special case. If an expression for $P(\xi[0, z] = j)$ is not available for an OS point process, one could view (2), or (4), as an expectation with respect to $P(\xi[0, z] = j)$ and apply Monte Carlo simulation, generating $\xi[0, z]$, calculating the multiple integral of dimension $\xi[0, z]$ and then averaging over the number of simulations.
2. In order to compute the j -dimensional integral in (2), or (4), one can view it as an appropriate expectation with respect to the order statistics $Y_{1,j}, Y_{2,j}, \dots, Y_{j,j}$ of j independent uniformly distributed on $[0, h_z(z)]$ random variables and apply a Monte Carlo approach (employing only order statistics of uniforms) to compute the integral. This approach has been applied and

thoroughly investigated in Section 5 of [Dimitrova et al. \(2016\)](#) for computing similar multiple integrals. More precisely, following [Dimitrova et al. \(2016\)](#), one can rewrite (4) as

$$\begin{aligned}
 P(g_z(t) \leq S_t \leq h_z(t), 0 \leq t \leq z) = & \\
 \sum_{j=0}^{\infty} P(\xi[0, z] = j) V_j \int_0^{h_z(z)} \int_{y_1}^{h_z(z)} \cdots \int_{y_{j-1}}^{h_z(z)} & \mathbb{1}_{g_z(z)}(y_j) \det \left(\left(\rho_{m,l}^{(j)} \right)_{1 \leq m, l \leq j} \right) \\
 \times f(y_1, \dots, y_j) \times \frac{1}{V_j} dy_j \cdots dy_1, &
 \end{aligned}$$

where

$$V_j = \int_0^{h_z(z)} \int_{y_1}^{h_z(z)} \cdots \int_{y_{j-1}}^{h_z(z)} 1 dy_j \cdots dy_1 = \frac{h_z(z)^j}{j!}.$$

As is well known, $1/V_j$ can be viewed as the joint probability density function of the order statistics, $Y_{1,j}, Y_{2,j}, \dots, Y_{j,j}$, of j independent uniformly distributed on $[0, h_z(z)]$ random variables. Therefore, the multiple integrals in (2), or (4), can be interpreted as expectations with respect to the order statistics $Y_{1,j}, Y_{2,j}, \dots, Y_{j,j}$ and so, a Monte-Carlo simulation approach can be used to evaluate these. Thus, one may simulate, say $N > 0$, samples from the j -tuples $Y_{1,j}, Y_{2,j}, \dots, Y_{j,j}$ and utilizing the method for truncating the infinite sum so that a prescribed accuracy is achieved, as outlined in item 1 above (see also section 4 of [Dimitrova et al. \(2016\)](#)), we arrive at

$$\begin{aligned}
 P(g_z(t) \leq S_t \leq h_z(t), 0 \leq t \leq z) & \\
 \approx \left(P(\xi[0, z] = 0) + \sum_{k=1}^N \sum_{j=1}^{j_k^*} P(\xi[0, z] = j) V_j^c \mathbb{1}_{g_z(z)}(y_{j,j}) \right) & \quad (20) \\
 \times \det \left(\left(\rho_{m,l}^{(j)} \right)_{1 \leq m, l \leq j} \right) \times f(y_{1,j}^k, \dots, y_{j,j}^k) / N &
 \end{aligned}$$

It should be noted that this Monte Carlo approach can be applied also when the Y_i 's are discrete, i.e., with respect to (15), since although it is exact, its evaluation may still be prohibitively time consuming for very large values of $\lfloor h_z(z) \rfloor$. See [Dimitrova et al. \(2016\)](#) for numerical details and comparison with the direct Monte Carlo approach.

3. Finally, as noted in Remark 1, the determinants in (2) and (4) are of a Appell-Hessenberg type, they can be efficiently evaluated using recurrence formula (8), noting that

$$\begin{aligned}
 j! \det \left(\left(\delta_{m,l}^{(j)} \right)_{1 \leq m, l \leq j} \right) &= \det \left(\left(\rho_{m,l}^{(j)} \right)_{1 \leq m, l \leq j} \right) \\
 = P \left(\bigcap_{i=1}^j \left(F_z \left(h_z^{-1}(y_i) - \right) \leq U_{i,j} \leq F_z \left(g_z^{-1}(y_{i-1}) \right) \right) \right) &= d_j, \quad (21)
 \end{aligned}$$

with $\alpha_i = F_z(h_z^{-1}(y_i) -)$ and $\beta_i = F_z(g_z^{-1}(y_{i-1}))$.

Next for illustrative purposes we specify formula (4), for the case of two particular OS arrival processes, the non-homogeneous linear birth process, known also as the Pólya-Lundberg process for which $F_z(t)$ is continuous, and an (OS) Poisson process with Poisson clusters at fixed instants, where $F_z(t)$ has jump discontinuities.

First we consider a double boundary non-crossing problem equivalent to a non-ruin probability model, as described in Section 3.1. We assume the arrival of insurance claims is modelled by a Pólya-

Lundberg process with parameters $\lambda > 0$ and $b > 0$. The latter is a non-homogeneous linear birth process for which $\xi[0, z]$ has a negative binomial distribution, i.e.,

$$P(\xi[0, z] = j) = \binom{j-1+1/b}{j} \left(\frac{\lambda bz}{1+\lambda bz}\right)^j \left(\frac{1}{1+\lambda bz}\right)^{1/b}, j \geq 0, \tag{22}$$

As known, the Pólya-Lundberg process has cdf $F_z(t) = t/z$ for $0 \leq t \leq z$ with $F_z(t) = 1$ for $t \geq z$. It admits representation as a mixed poisson Process, i.e., $\xi(t) = N(Wt)$, where $N(t)$ is a homogeneous Poisson process with unit rate and W is Gamma distributed with shape parameter $1/b$ and scale parameter $1/\lambda b$, i.e., $W \sim \text{Gamma}(1/b, 1/\lambda b)$. Assuming such an OS process for the claim arrivals, we specify the remaining parameters of the underlying OS risk model as follows.

Example 1. Consider a double boundary non-crossing problem equivalent to a non-ruin probability model as described in Section 3.1, where the arrival of claim amounts is modelled by a Pólya-Lundberg process with parameters $\lambda = 2$ and $b = 1$. Assume the upper boundary equivalent to the premium income function is $h_z(t) = t^2 + 1.5$, the time horizon $z = 2$, the lower boundary function $g_z(t) \equiv 0, \forall t \geq 0$ and the partial claim sums, Y_1, Y_2, \dots to be the integers $1, 2, \dots$, respectively. Utilizing expression (4) of Theorem 1, and more precisely Corollary 2, for the non-ruin probability we have

$$\begin{aligned} P(0 \leq S_t \leq h_z(t), 0 \leq t \leq z) &= P(T > z) \\ &= \sum_{j=0}^5 \binom{j-1+1/b}{j} \left(\frac{\lambda bz}{1+\lambda bz}\right)^j \left(\frac{1}{1+\lambda bz}\right)^{1/b} \det \left(\left(\rho_{m,l}^{(j)}\right)_{1 \leq m,l \leq j} \right) \\ &= 1/2 + \sum_{j=1}^5 1 \times \left(\frac{4}{5}\right)^j \left(\frac{1}{5}\right)^1 \det \left(\left(\rho_{m,l}^{(j)}\right)_{1 \leq m,l \leq j} \right) \\ &= 0.568265 \end{aligned} \tag{23}$$

where we have used that $\lfloor h_z(z) \rfloor = 5, g_z^{-1}(0) = z$ and where the number 0.568265 is computed using the recurrence formula (8) to evaluate the determinants, following (21) and using (17) substituting in it $F_z(h_z^{-1}(y_l) -) = \min(z, \sqrt{y_l - 1.5})/z$ for $y_l \geq 1.5$ or zero otherwise.

Let us note that this example has been considered by Goffard and Lefevre (2017) and our result of $P(T > z) = 0.568265$ coincides with theirs.

Second, to demonstrate the generality and flexibility of our model, consider an OS point process, ξ , with a continuous component and a pure jump component in the underlying cdf, $F_z(x)$, i.e., $\xi(0, x] = \eta(0, x] + \xi_1 \mathbb{1}_{(0,x]}(t_1) + \dots + \xi_j \mathbb{1}_{(0,x]}(t_j)$, where $\eta(0, x]$ is a Poisson process with unit rate, defined on $(0, \infty]$ and independent of the Poisson random variables, $\xi_1, \dots, \xi_j, \xi_i \in \mathcal{P}(\mu_i), i = 1, \dots, j$, assumed also mutually independent. By construction, ξ is an OS process with independent increments. It could be particularly suitable for applications, especially when data comes from two (or more) independent insurance portfolios (lines of business), among which one with claim frequency data at fixed instants $0 < t_1 < t_2 < \dots < t_j$ (e.g., annual observations) and a second one with data at policy level of the instants of claiming. In view of the Solvency II requirements, it would be instructive to be able to evaluate the probability of non-ruin in a finite time interval, $(0, z]$, due to claims coming from all lines of business. Without loss of generality, we will illustrate this, based on the following example.

Example 2. Let $0 < t_1 < t_2 \leq z$. Then, the cumulative intensity function, $\Lambda(x)$ is

$$\Lambda(x) = \begin{cases} x & \text{for } 0 < x < t_1, \\ x + \mu_1 & \text{for } t_1 \leq x < t_2, \\ x + \mu_1 + \mu_2 & \text{for } t_2 \leq x \leq z, \end{cases}$$

and the related cdf

$$F_z(x) = \begin{cases} 0 & \text{for } -\infty < x < 0, \\ x/(z + \mu_1 + \mu_2) & \text{for } 0 \leq x < t_1, \\ (x + \mu_1)/(z + \mu_1 + \mu_2) & \text{for } t_1 \leq x < t_2, \\ (x + \mu_1 + \mu_2)/(z + \mu_1 + \mu_2) & \text{for } t_2 \leq x \leq z, \\ 1 & \text{for } z < x < +\infty. \end{cases}$$

Take the upper boundary (premium income) function $h_z(x)$ to be,

$$h_z(x) = \begin{cases} 0.5 & \text{for } x < t_1, \\ 1.5 & \text{for } t_1 \leq x < t_2, \\ 2.5 & \text{for } t_2 \leq x, \end{cases} \tag{24}$$

the lower boundary function to be $g_z(x) \equiv 0, \forall x \geq 0$ and the partial claim sums, Y_1, Y_2, \dots to be the integers $1, 2, \dots$, respectively. Applying Equation (4) of Theorem 1, and more precisely Corollary 3, for the finite-time non-ruin probability we have

$$\begin{aligned} P(0 \leq S_t \leq h_z(t), 0 \leq t \leq z) &= P(T > z) \\ &= P(\zeta(0, z] = 0) + \sum_{j=1}^{\lfloor h_z(z) \rfloor} P(\zeta(0, z] = j) \det \left(\left(\rho_{m,l}^{(j)} \right)_{1 \leq m,l \leq j} \right) \\ &= \exp(-z + \mu_1 + \mu_2) + (z + \mu_1 + \mu_2) \exp(-z + \mu_1 + \mu_2) \rho_{1,1}^{(1)} \\ &+ \frac{(z + \mu_1 + \mu_2)^2}{2!} \exp(-z + \mu_1 + \mu_2) \det \left(\left(\rho_{m,l}^{(2)} \right)_{1 \leq m,l \leq 2} \right) \\ &= \exp(-z + \mu_1 + \mu_2) \left(1 + z(1 + \mu_1 + \mu_2 + \frac{z}{2} - t_1) + t_1 t_2 - \frac{t_2^2}{2} - \mu_1 t_2 \right. \\ &\left. + \mu_1 \mu_2 + \frac{\mu_2^2}{2} - \mu_2 t_1 + \mu_1 + \mu_2 - t_1 \right), \end{aligned} \tag{25}$$

where we have used that $\lfloor h_z(z) \rfloor = 2$ and where the last equality follows expanding the determinants, following (17) and performing some straightforward algebra.

Remark 2. The last expression in (25), can be verified through direct but tedious calculations, see [Dimitrova et al. \(2014\)](#).

Author Contributions: The theoretical part of the work was conducted jointly by D.D., Z.I. and V.K. The paper was written by V.K. and D.D.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A. Proof of Lemma 1

From [David and Nagaraja \(2003\)](#), (see p. 15 therein) we have

$$(X_{1,j}, \dots, X_{j,j}) \stackrel{d}{=} \left(F_z^{-1}(U_{1,j}), \dots, F_z^{-1}(U_{j,j}) \right), \tag{A1}$$

where $\stackrel{d}{=}$ stands for equality in distribution.

From (A1), we have

$$\begin{aligned}
 P\left(\bigcap_{i=1}^j (\alpha_i < X_{i,j} \leq \beta_i)\right) &= P\left(\bigcap_{i=1}^j (\alpha_i < F_z^{-1}(U_{i,j}) \leq \beta_i)\right) \\
 &= P\left(\bigcap_{i=1}^j [\{F_z^{-1}(U_{i,j}) \leq \beta_i\} \setminus \{F_z^{-1}(U_{i,j}) \leq \alpha_i\}]\right) \\
 &= P\left(\bigcap_{i=1}^j [\{U_{i,j} \leq F_z(\beta_i)\} \setminus \{U_{i,j} \leq F_z(\alpha_i)\}]\right) = P\left(\bigcap_{i=1}^j (F_z(\alpha_i) < U_{i,j} \leq F_z(\beta_i))\right).
 \end{aligned} \tag{A2}$$

The third equality in (A2) follows since, for a right-continuous cdf $F_z(t)$ the inequality $u \leq F_z(t)$ holds iff $F_z^{-1}(u) \leq t$ (see e.g., David and Nagaraja (2003)), where one can replace u with $U_{i,j}$. Consider now the probability on the left-hand side of (7). We have

$$\begin{aligned}
 P\left(\bigcap_{i=1}^j (\alpha_i \leq X_{i,j} \leq \beta_i)\right) &= \lim_{\epsilon \rightarrow 0} P\left(\bigcap_{i=1}^j ((\alpha_i - \epsilon) < X_{i,j} \leq \beta_i)\right) \\
 &= \lim_{\epsilon \rightarrow 0} P\left(\bigcap_{i=1}^j (F_z(\alpha_i - \epsilon) < U_{i,j} \leq F_z(\beta_i))\right) = P\left(\bigcap_{i=1}^j (F_z(\alpha_{i-}) < U_{i,j} \leq F_z(\beta_i))\right) \\
 &= P\left(\bigcap_{i=1}^j (F_z(\alpha_{i-}) \leq U_{i,j} \leq F_z(\beta_i))\right),
 \end{aligned} \tag{A3}$$

where in the second equality of (A3) we have used (A2) and the last equality follows since $U_{i,j}$, $i = 1, \dots, j$ are absolutely continuous. \square

References

- Borovkov, K., and Alexander Novikov. 2005. Explicit Bounds for approximation rates of boundary crossing probabilities for the Wiener process. *Journal of Applied Probability* 42: 82–92.
- Bühlmann, Hans 1970. *Mathematical Methods in Risk Theory*. Heidelberg: Springer.
- Buonocore, A., V. Giorno, A. G. Nobile, and L. M. Ricciardi. 1990. On the two-boundary first-crossing-time problem for diffusion processes. *Journal of Applied Probability* 27: 102–14.
- Crump, Kenny S. 1975. On point processes having an order statistics structure. *Sankhyā: The Indian Journal of Statistics, Series A* 37: 396–404.
- David, Herbert A., and Haikady N. Nagaraja. 2003. *Order Statistics*, 3rd ed. New York: John Wiley.
- De Vylder, Floriaen, and Marc Goovaerts. 1999. Inequality extensions of Prabhu's formula in ruin theory. *Insurance: Mathematics and Economics* 24: 249–71.
- De Vylder, Floriaen, and Marc Goovaerts. 2000. Homogeneous risk models with equalized claim amounts. *Insurance: Mathematics and Economics* 26: 223–38.
- Dimitrova, Dimitrina S., Zvetan G. Ignatov, and Vladimir K. Kaishev. 2015. On the first crossing of two boundaries by an order statistics risk process. Available online: https://www.researchgate.net/publication/306364994_On_the_first_crossing_of_two_boundaries_by_an_order_statistics_risk_process (accessed on 22 August 2016).
- Dimitrova, Dimitrina S., Zvetan G. Ignatov, and Vladimir K. Kaishev. 2014. Ruin and deficit under claim arrivals with the order statistics property. Available online: https://www.researchgate.net/publication/306364689_Ruin_and_deficit_under_claim_arrivals_with_the_order_statistics_property (accessed on 22 August 2016).
- Dimitrova, Dimitrina S., Vladimir K. Kaishev, and Shouqi Zhao. 2016. On the evaluation of finite-time ruin probabilities in a dependent risk model. *Applied Mathematics and Computation* 275: 268–86.
- Dimitrova, Dimitrina S., Vladimir K. Kaishev, and Shouqi Zhao. 2015. On finite-time ruin probabilities in a generalized dual risk model with dependence. *European Journal of Operational Research* 242: 134–48.

- Epanechnikov, V. A. 1968. The significance level and power of two-sided Kolmogorov's test in case of small sample sizes. *Teoriya Veroyatnostei i ee Primeneniya* 13: 725–30.
- Feigin, Paul D. 1979. On the characterization of point processes with the order statistic property. *Journal of Applied Probability* 16: 297–304.
- Goffard, Pierre-Olivier. 2017. Two-sided exit problems in the ordered risk model. Available online: <https://hal.archives-ouvertes.fr/hal-01528204> (accessed on 6 June 2017).
- Goffard, Pierre-Olivier, and Claude Lefèvre. 2017. Boundary crossing of order statistics point processes. *Journal of Mathematical Analysis and Applications* 447: 890–907.
- Haccou, Patsy, Peter Jagers, and Vladimir A. Vatutin. 2005. *Branching Processes: Variation, Growth, and Extinction of Populations*. Cambridge: Cambridge University Press.
- Ignatov, Zvetan G., and Vladimir K. Kaishev. 2000. Two-sided bounds for the finite-time probability of ruin. *Scandinavian Actuarial Journal* 2000: 46–62.
- Ignatov, Zvetan G., and Vladimir K. Kaishev. 2004. A Finite-time ruin probability formula for continuous claim severities. *Journal of Applied Probability* 41: 570–78.
- Ignatov, Zvetan G., and Vladimir K. Kaishev. 2016. First crossing time, overshoot and Appell-Hessenberg type functions. *Stochastics* 88: 1240–60.
- Ignatov, Zvetan G., Vladimir K. Kaishev, and Rossen S. Krachunov. 2001. An Improved Finite-time Ruin Probability Formula and its "Mathematica" Implementation. *Insurance: Mathematics and Economics* 29: 375–86.
- Kendall, David G. 1949. Stochastic Processes and Population Growth. *Journal of the Royal Statistical Society. Series B* 11: 230–82.
- Lefèvre, Claude, and Stéphane Loisel. 2009. Finite-time ruin probabilities for discrete, possibly dependent, claim severities. *Methodology and Computing in Applied Probability* 11: 425–41.
- Lefèvre, Claude, and Philippe Picard. 2011. A new look at the homogeneous risk model. *Insurance: Mathematics and Economics* 49: 512–19.
- Lefèvre, Claude, and Philippe Picard. 2014. Ruin probabilities for risk models with ordered claim arrivals. *Methodol Comput Appl Probab* 16: 885–905.
- Lotov, V. I. 1996. On some boundary crossing problems for Gaussian random walks. *The Annals of Probability* 24: 2154–71.
- Nawrotzki, Kurt. 1962. Ein Grenzwertsatz für homogene zufällige Punktfolgen (Verallgemeinerung eines Satzes von A. Rnyi). *Mathematische Nachrichten* 24: 201–17.
- Picard, Philippe, and Claude Lefèvre. 1997. The probability of ruin in finite time with discrete claim size distribution. *Insurance Mathematics and Economics* 20: 260–61.
- Pötzelberger, Klaus, and Liqun Wang. 2001. Boundary Crossing Probability For Brownian Motion. *Journal of Applied Probability* 38: 152–64.
- Puri, Prem S. 1982. On the characterization of point processes with the order statistic property without the moment condition. *Journal of Applied Probability* 19: 39–51.
- Sendova, Kristina P., and Ričardas Zitikis. 2012. The order statistic claim process with dependent claim frequencies and severities. *Journal of Statistical Theory and Practice* 6: 597–620.
- Steck, G. P. 1971. Rectangle probabilities for uniform order statistics and the probability that the empirical distribution function lies between two distribution functions, *The Annals of Mathematical Statistics* 42: 1–11.
- Teunen, Marleen, and Marc Goovaerts. 1994. Double boundary crossing result for the brownian motion. *Scandinavian Actuarial Journal* 2: 139–50.
- Thompson, W.A. 1988. *Point Process Models with Applications to Safety and Reliability*. Boston: Springer.
- Vein, Robert, and Paul Dale. 1999. *Determinants and Their Applications in Mathematical Physics*. New York: Springer-Verlag.
- Wald, Abraham, and Jacob Wolfowitz. 1939. Confidence limits for continuous distribution functions. *The Annals of Mathematical Statistics* 10: 105–18.

Willmot, Gordon E. 1989. The total claims distribution under inflationary conditions. *Scandinavian Actuarial Journal* 1: 1–12.

Xu, Yifan. 2012. First exit times of compound Poisson processes with parallel boundaries. *Sequential Analysis* 31: 135–44.



© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).