



City Research Online

City, University of London Institutional Repository

Citation: Coletti, E., Forini, V., Nardelli, G., Grignani, G. & Orselli, M. (2004). Exact potential and scattering amplitudes from the tachyon non-linear β -function. *Journal of High Energy Physics*, 2004(03), 030. doi: 10.1088/1126-6708/2004/03/030

This is the published version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/19747/>

Link to published version: <https://doi.org/10.1088/1126-6708/2004/03/030>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

Exact potential and scattering amplitudes from the tachyon non-linear β -function

To cite this article: Erasmo Coletti *et al* JHEP03(2004)030

View the [article online](#) for updates and enhancements.

Related content

- [Boundary string field theory as a field theory—mass spectrum and interaction](#)
Koji Hashimoto and Seiji Terashima
- [Spectral flow and boundary string field theory for angled D-branes](#)
Nicholas T. Jones and S.-H. Henry Tye
- [Abelian and nonabelian vector field effective actions from string field theory](#)
Erasmo Coletti, Ilya Sigalov and Washington Taylor

Recent citations

- [Tachyon solutions in boundary and open string field theory](#)
Gianluca Calcagni
- [Nonlocal instantons and solitons in string models](#)
Gianluca Calcagni and Giuseppe Nardelli
- [Superstring disk amplitudes in a rolling tachyon background](#)
Esko Keski-Vakkuri *et al*

Exact potential and scattering amplitudes from the tachyon non-linear β -function

Erasmus Coletti

*Center for Theoretical Physics, M.I.T.
Cambridge, MA 02139, U.S.A.
E-mail: colettie@mit.edu*

Valentina Forini and Giuseppe Nardelli

*Dipartimento di Fisica and I.N.F.N. Gruppo Collegato di Trento, Università di Trento
38050 Povo (Trento), Italia
E-mail: forini@science.unitn.it, nardelli@science.unitn.it*

Gianluca Grignani and Marta Orselli

*Dipartimento di Fisica and Sezione I.N.F.N., Università di Perugia
Via A. Pascoli I-06123, Perugia, Italia
E-mail: gianluca.grignani@pg.infn.it, marta.orselli@pg.infn.it*

ABSTRACT: We compute, on the disk, the non-linear tachyon β -function, β^T , of the open bosonic string theory. β^T is determined both in an expansion to the third power of the field and to all orders in derivatives and in an expansion to any power of the tachyon field in the leading order in derivatives. We construct the Witten-Shatashvili (WS) space-time effective action S and prove that it has a very simple universal form in terms of the renormalized tachyon field and β^T . The expression for S is well suited to studying both processes that are far off-shell, such as tachyon condensation, and close to the mass-shell, such as perturbative on-shell amplitudes. We evaluate S in a small derivative expansion, providing the exact tachyon potential. The normalization of S is fixed by requiring that the field redefinition that maps S into the tachyon effective action derived from the cubic string field theory is regular on-shell. The normalization factor is in precise agreement with the one required for verifying all the conjectures on tachyon condensation. The coordinates in the space of couplings in which the tachyon β -function is non linear are the most appropriate to study RG fixed points that can be interpreted as solitons of S , i.e. D-branes.

KEYWORDS: Tachyon Condensation, String Field Theory.

Contents

1. Introduction	1
2. Boundary string field theory	4
3. Integration over the bulk variables	6
4. Partition function on the disk and the renormalized tachyon field	7
5. Background-field method	11
6. β-function	13
7. Witten-Shatashvili action	17
8. Cubic vs. Witten-Shatashvili tachyon effective actions	22
9. Conclusions	24
A. Computation of $I(k_1, k_2, k_3)$	25

1. Introduction

One of the most interesting problems in string theory is to understand how the background space-time on which the string propagates arises in a self-consistent way. For open strings, there are two main approaches to this problem, cubic string field theory [1] and background independent string field theory [2]–[5].

Background independent open string field theory has been useful for finding the classical tachyon potential energy functional and the leading derivative terms in the tachyon effective action [6]–[9]. It is formulated as a problem in boundary conformal field theory. One begins with the partition function of open-string theory where the world-sheet is a disk. The strings in the bulk are considered to be on-shell and a boundary interaction with arbitrary operators is added. The configuration space of open string field theory is then taken to be the space of all possible boundary operators modulo gauge transformations and field redefinitions. Renormalization fixed points, which correspond to conformal field theories, are solutions of classical equations of motion and should be viewed as the solutions of classical string field theory.

Due to the existence of a tachyon in the bosonic string theory, the 26-dimensional Minkowski space background about which the string is quantized is unstable. An unstable state should decay to something and the nature of both the decay process and the endpoint

of the decay are crucial questions. Some understanding of this process has been achieved for the open bosonic string. The key idea is that of Sen [10]. The open bosonic string tachyon reflects the instability of the D-25 brane. This unstable D-brane should decay by condensation of the open string tachyon field. The energy per unit volume released in the decay should be the D-25 brane tension and the end-point of the decay is the closed string vacuum [10, 11, 12, 7]. There are also intermediate unstable states which are the D-branes of all dimensions between zero and 25.

The decay of unstable systems of D-branes, pictured as a tachyon field rolling down a potential toward a stable minimum, can also be addressed in the context of the boundary string field theory. It involves deforming the world sheet conformal field theory of the unstable D-brane by an exact marginal, time dependent tachyon profile. The quantitative study of rolling tachyons was initiated by A. Sen [13]–[17] and has been recently used to obtain other forms of tachyon effective actions [18]–[22]. The construction of the space-time tachyon effective action in background independent open string field theory is the subject of the present paper.

By classical power-counting the tachyon field $T(X)$ has dimension one and is a relevant operator. If $T(X)$ is the only interaction, the field theory is perturbatively super-renormalizable. If $T(X)$ and the other fields are adjusted so that the sigma model that they define is at an infrared fixed point of the renormalization group (RG), these background fields are a solution of the classical equations of motion of string theory. Witten and Shatashvili [2, 4] have argued that these equations of motion come from an action which can be derived from the disk partition function Z by a prescription which we shall make use of below. According to this prescription the effective action for a generic coupling constant g^i (which can be identified with the tachyon, the gauge or any other field that correspond to excitations of the open bosonic string) is related to the renormalized partition function of open string theory on the disk, $Z(g^i)$, through

$$S = \left(1 - \beta^i \frac{\delta}{\delta g^i} \right) Z(g^i), \tag{1.1}$$

where β^i is the beta-function¹ of the coupling g^i . Note that (1.1) fixes the additive ambiguity in S by requiring that at RG fixed points g^* , in which $\beta^i(g^*) = 0$,

$$S(g^*) = Z(g^*). \tag{1.2}$$

The derivative of the action S with respect to the coupling constant g^i must be related to the β -function through a metric according to

$$\frac{\partial S}{\partial g^i} = -\beta^j G_{ij}(g). \tag{1.3}$$

G_{ij} should be a non-degenerate metric, otherwise there would be an extra zero which could not be interpreted as a conformal field theory on the world sheet. Eq. (1.3) indicates that the RG flow is actually a gradient flow. The prescription (1.1) provides a definition of the metric G_{ij} in the space of couplings.

¹In this paper the β function is positive for relevant perturbations. In some other papers on the subject, e.g. [7], the opposite conventions are used.

The β -functions appearing in (1.1) are in general non-linear functions of the couplings g^i . When the linear parts of the β^i (i.e. the anomalous dimensions λ_i of the corresponding coupling) satisfy a so called “resonant condition”, the non linear parts of the β -function cannot be removed by a coordinate redefinition in the space of couplings [5]. Such resonant condition is nothing but the mass-shell condition so that, near the mass-shell, the β -functions are necessarily non-linear.

However, when the resonant condition does not hold, a possible choice of coordinates on the space of string fields is one in which the β -functions are exactly linear. This choice can always be made locally [7] and is well suited to studying processes which are far off-shell, such as tachyon condensation. These coordinates, however, become singular when the components of the string field (e.g. $T(X)$, $A_\mu(X)$ etc.) go on-shell. These coordinates can be used to construct, for example, the tachyon effective potential, but become singular when one tries to derive an effective action which reproduces the on-shell amplitudes. In particular, if the Veneziano amplitude needs to emerge from the tachyon effective action it is necessary to consider the whole non-linear β -function in (1.1). A complete renormalization of the theory in fact makes the β -function non-linear in $T(X)$ [23] so that, since the vanishing of the β -function is the field equation for T , these nonlinear terms describe tachyon scattering. One of the goal of this paper is to construct non-linear expressions for the β -functions which are valid away from the RG fixed point. With these expressions for the non-linear tachyon β -function we shall construct the Witten-Shatashvili (WS) space-time action (1.1). We shall prove that (1.1) has the following very simple form in the coupling space coordinates in which the tachyon β -function is non-linear

$$S = K \int d^{26}X [1 - T_R(X) + \beta^T(X)], \tag{1.4}$$

where T_R is the renormalized tachyon field and K is a constant related to the D25-brane tension. This formula is universal as it does not depend on how many couplings are switched on. Eq. (1.4) arises from the expression that links the renormalized tachyon field to the partition function that appears in (1.1), namely $Z = K \int d^{26}X (1 - T_R)$. T_R is then a non-linear function of the bare coupling T and in these coordinates the β -function is non-linear. When couplings other than the tachyon are introduced in Z , β^T will depend on them so that S will provide the space-time effective action also for these couplings.

With this prescription we shall compute the non-linear β -function β^T for the tachyon field up to the third order in powers of the field and to any order in derivatives of the field. From this we shall show that the solutions of the RG fixed point equations generate the three and four-point open bosonic string scattering amplitudes involving tachyons. Then, with the same renormalization prescription, we shall compute β^T to the leading orders in derivatives but to any power of the tachyon field and we shall show that S coincides with the one-found in [6, 7, 8]. Obviously, S up to the first three powers of T and expanded to the leading order in powers of derivatives can be obtained from both calculations and the results coincide.

In the case of profiles $T_R(k)$ that have support near the on-shell momentum $k^2 \simeq 1$ the equation $\beta^T(k) = 0$ can be derived as the equation of motion of an action. We shall show

that this action coincides with the tachyon effective action computed, for the almost on-shell profiles, from the cubic string field theory up to the fourth power of the tachyon field.

The knowledge of the non-linear tachyon β -function is very important also for another reason. The solutions of the equation $\beta^T = 0$ give the conformal fixed points, the backgrounds that are consistent with the string dynamics. In the case of slowly varying tachyon profiles, we shall show that the equations of motion for the WS action can be made identical to the RG fixed point equation $\beta^T = 0$. We shall find solutions of this equation to which correspond a finite value of the WS action. Being solutions of the RG equations, these solitons are lower dimensional D-branes for which the finite value of S provides a quite accurate prediction of the D-brane tension.

We shall also show that the WS action constructed in terms of a linear β -function [24] is related to the action (1.4) by a field redefinition, and that this field redefinition becomes singular on-shell. This is in agreement with the Poincaré-Dulac theorem [25] used in [5] to prove that when the resonant condition holds, namely near the on-shellness, the β -function has to be non-linear.

The tachyon effective action up to the third power in the fields is known exactly also from the cubic string field theory [1]. This raises the interesting question of how the action S obtained in this paper is related to the cubic SFT. It seems clear that the cubic SFT must correspond to (1.1), (1.3) for a particular choice of coordinates on the space of string fields (or worldsheet couplings). The two sets of coordinates are related by a complicated transformation which we shall derive in this paper. The cubic SFT parametrization of worldsheet RG is regular close to the mass shell. It very well reproduces tachyon scattering [26], to it must correspond a non-linear beta-function. Thus a coordinate transformation that relates the two effective actions needs a non-linear beta function in the definition (1.1). We shall show that this field redefinition exists and that it is non-singular on-shell only when K in (1.4) coincides with the tension of the D25-brane, in agreement with all the conjectures involving tachyon condensation [10, 27, 28].

2. Boundary string field theory

In Witten's construction of open boundary string field theory [2] the space of all two dimensional worldsheet field theories on the unit disk, which are conformal in the interior of the disk but have arbitrary boundary interactions, is described by the world-sheet action

$$S = S_0 + \int_0^{2\pi} \frac{d\tau}{2\pi} \mathcal{V} \tag{2.1}$$

where S_0 is a free action describing an open plus closed conformal background and \mathcal{V} is a general perturbation defined on the disk boundary. We will discuss the twenty six dimensional bosonic string, for which (2.1) can be expressed in terms of a derivative expansion (or level expansion) of the form

$$\mathcal{V} = T(X) + iA_\mu(X)\partial_\tau X^\mu + B_{\mu\nu}(X)\partial_\tau X^\mu\partial_\tau X^\nu + C_\mu(X)\partial_\tau^2 X^\mu + \dots \tag{2.2}$$

Without the perturbation \mathcal{V} the boundary conditions on X are $\partial_r X^\mu|_{r=1} = 0$, where r is the radial variable on the disk.

\mathcal{V} is a ghost number zero operator and it is useful to introduce a ghost number one operator \mathcal{O} via

$$\mathcal{V} = b_{-1}\mathcal{O}. \tag{2.3}$$

We shall consider the simplest case in which ghosts decouple from matter so that, as in (2.2), \mathcal{V} is constructed out of matter fields alone

$$\mathcal{O} = c\mathcal{V}. \tag{2.4}$$

The space-time string field theory action S is defined through its derivative dS which is a two point function computed with the worldsheet action (2.1). More generally one can introduce some basis elements \mathcal{V}_i for operators of ghost number 0 so that the space of boundary perturbations \mathcal{V} can be parametrized as

$$\mathcal{V} = \sum_i g^i \mathcal{V}_i \tag{2.5}$$

where the coefficients g^i are couplings on the world-sheet theory, which are regarded as fields from the space-time point of view, and $\mathcal{O} = \sum_i g^i \mathcal{O}_i$. In this parametrization the space-time action is defined through its derivatives with respect to the couplings and has the form

$$\frac{\partial S}{\partial g^i} = \frac{K}{2} \int_0^{2\pi} \frac{d\tau}{2\pi} \int_0^{2\pi} \frac{d\tau'}{2\pi} \langle \mathcal{O}_i(\tau) \{Q, \mathcal{O}(\tau')\} \rangle_g, \tag{2.6}$$

where Q is the BRST charge and the correlator is evaluated with the full perturbed worldsheet action \mathcal{S} .

If \mathcal{V}_i is a conformal primary field of dimension Δ_i , for \mathcal{O} 's of the form (2.4), one has

$$\{Q, c\mathcal{V}_i\} = (1 - \Delta_i)c\partial_\tau c\mathcal{V}_i, \tag{2.7}$$

so that from (2.6) one gets

$$\frac{\partial S}{\partial g^i} = -(1 - \Delta_j)g^j G_{ij}(g), \tag{2.8}$$

where

$$G_{ij} = 2K \int_0^{2\pi} \frac{d\tau}{2\pi} \int_0^{2\pi} \frac{d\tau'}{2\pi} \sin^2\left(\frac{\tau - \tau'}{2}\right) \langle \mathcal{V}_i(\tau)\mathcal{V}_j(\tau') \rangle_g. \tag{2.9}$$

Eq. (2.8) cannot be true in general, since it does not transform covariantly under reparametrizations of the space of theories, $g^j \rightarrow f^j(g^i)$. Indeed, $\partial_i S$ and G_{ij} transform as tensors, (the latter is the metric on the space of worldsheet theories), but g^i does not.

The correct covariant generalization of (2.8) was given in [4, 5]. The worldsheet RG defines a natural vector field on the space of theories: the β -function $\beta^i(g)$, which transforms as a covariant vector under reparametrizations of g^i . The covariant form of (2.8) is thus (1.3). If we assume that total derivatives inside the correlation function decouple and that there are no contact terms, it turns out that the β -function in (1.1) is the linear β -function. This implies that the equations of motion derived from the action (1.1) are just linear. However, as shown by Shatashvili [4, 5], contact terms show up in the computation

on the world-sheet and cannot be ignored. The point is that the operator Q , which is constructed out of the BRST operator in the bulk and should be independent on the couplings because the perturbation is on the boundary, actually depends on the couplings when the contour integral approaches the boundary of the disk. A way to fix the structure of the contact terms is to consider that, since dS is a one-form, the derivative of dS should be zero independently of the choice of the contact terms that one makes in the computation. This leads to the following formula for the vector field in equation (1.1)

$$\beta^i = (1 - \Delta_i)g^i + \alpha_{jk}^i g^j g^k + \gamma_{jkl}^i g^j g^k g^l + \dots \quad (2.10)$$

This is an expression for the β -function with all the non-linear terms. According to the Poincare-Dulac Theorem about vector fields (whose relevance to the β -function related issues was stressed many times by Zamolodchikov [25]) every vector field can be linearized by an appropriate redefinition of the coordinates up to the resonant term. In the second order of equation (2.10) the resonance condition is given by

$$\Delta_j + \Delta_k - \Delta_i = 1. \quad (2.11)$$

The resonance condition means that the β -function cannot be linearized by a coordinate transformation and that all the non-linear terms cannot be removed from the β -function equation (2.10). When g^i is the tachyon field $T(k)$, the resonant condition (2.11) corresponds to the mass-shell conditions for three tachyons. We shall prove in what follows that the WS action S up to the third order in the tachyon fields, constructed in terms of the linear β -function [24], is related to the S made of a non-linear β -function by a field redefinition, but that this field redefinition becomes singular on-shell.

3. Integration over the bulk variables

Let us now restrict ourselves to the specific example of open strings propagating in a tachyon background. The partition function reads

$$Z = \int [dX^\mu(\sigma, \tau)] \exp(-S[X]), \quad (3.1)$$

where the action is

$$S[X] = \int d\sigma d\tau \frac{1}{4\pi} \partial_a X(\sigma, \tau) \cdot \partial_a X(\sigma, \tau) + \int_0^{2\pi} \frac{d\tau}{2\pi} T(X(\tau)). \quad (3.2)$$

Here, the first term in (3.2) is the bulk action and is integrated over the volume of the unit disk. The second term in (3.2) is integrated on the circle which is the boundary of the unit disk and describes the interactions. The scalar fields X^μ have D components with $\mu = 1, \dots, D$ and we shall assume $D = 26$ in what follows for a critical string. We are working in a system of units where $\alpha' = 1$.

We begin with the observation that the bulk excitations can be integrated out of (3.1) to get an effective non-local field theory which lives on the boundary [29]. To do this we write the field in the bulk as [9]

$$X = X_{cl} + X_{qu},$$

where

$$\partial^2 X_{\text{cl}} = 0$$

and X_{cl} approaches the fixed (for now) boundary value of X ,

$$X_{\text{cl}} \rightarrow X_{\text{bdry}} \text{ and } X_{\text{qu}} \rightarrow 0.$$

Then, in the bulk, the functional measure is $dX = dX_{\text{qu}}$ and

$$S = \int \frac{d^2\sigma}{4\pi} \partial X_{\text{qu}} \cdot \partial X_{\text{qu}} + \int \frac{d\tau}{2\pi} \left\{ \frac{1}{2} X^\mu |i\partial_\tau| X^\mu + T(X) \right\}, \quad (3.3)$$

where we omitted the cl index in the last integral. Then, the integration of X_{qu} produces a multiplicative constant in the partition function - the partition function of the Dirichlet string, which we shall denote K . The kinetic term in the boundary action is non-local. The absolute value of the derivative operator is defined by the Fourier transform,

$$|i\partial_\tau| \delta(\tau - \tau') = \sum_n \frac{|n|}{2\pi} e^{in(\tau - \tau')}.$$

The partition function of the boundary theory is then

$$Z(J) = K \int [dX_\mu] e^{-\int_0^{2\pi} \frac{d\tau}{2\pi} (\frac{1}{2} X^\mu |i\partial| X^\mu + T(X) - J \cdot X)}, \quad (3.4)$$

where we have added a source $J^\mu(\tau)$ so that the path integral can be used as a generating functional for correlators of the fields X^μ restricted to the boundary. In particular, this source will allow us to compute the correlation functions of vertex operators of open string degrees of freedom. The remaining path integral over the boundary $X^\mu(\tau)$ defines a one-dimensional field theory with non-local kinetic term. If the tachyon field were absent ($T = 0$), the further integration over $X^\mu(\tau)$ would give a factor which converts the Dirichlet string partition function to the Neumann string partition function.

4. Partition function on the disk and the renormalized tachyon field

When only the tachyon field is considered as a boundary perturbation, the Witten-Shatashvili action is given by

$$S = \left(1 - \int \beta^T \frac{\delta}{\delta T} \right) Z, \quad (4.1)$$

where Z is the partition function of the boundary theory on the disk and β^T is the tachyon β -function. It is useful to introduce a constant source term k for the zero mode of the X field, the integral over the zero mode variable will just provide the energy-momentum conservation δ -function. The partition function (3.4) in the presence of this constant source reads

$$Z(k) = K \int [dX_\mu] e^{-\int_0^{2\pi} \frac{d\tau}{2\pi} (\frac{1}{2} X^\mu |i\partial_\tau| X^\mu + T(X) - ik \cdot \hat{X})}, \quad (4.2)$$

where \hat{X} is the zero mode which is defined by

$$\hat{X}^\mu = \int_0^{2\pi} \frac{d\tau}{2\pi} X^\mu(\tau). \quad (4.3)$$

In this section we shall expand the exponential in eq. (4.2) in powers of $T(X)$. The first non-trivial term is

$$Z^{(1)}(k) = -K \int [dX_\mu] \int dk_1 \int_0^{2\pi} \frac{d\tau_1}{2\pi} T(k_1) e^{-\int_0^{2\pi} \frac{d\tau}{2\pi} (\frac{1}{2} X^\mu |i\partial_\tau| X^\mu) - ik\hat{X} + ik_1 X(\tau_1)}. \quad (4.4)$$

The functional integral over the non-zero modes of $X(\tau)$ gives

$$Z^{(1)}(k) = -K \int d\hat{X}_\mu \int dk_1 T(k_1) e^{-\frac{k^2}{2} G(0) + i(k_1 - k)\hat{X}}, \quad (4.5)$$

where $G(\tau)$ is the Green function of the operator $|i\partial_\tau|$

$$G(\tau_1 - \tau_2) = 2 \sum_{n=1}^{\infty} e^{-\epsilon n} \frac{\cos n(\tau_1 - \tau_2)}{n} = -\log [1 - 2e^{-\epsilon} \cos(\tau_1 - \tau_2) + e^{-2\epsilon}] \quad (4.6)$$

and ϵ is a cut-off. In all the calculations we shall use the following prescription for $G(\tau)$

$$G(\tau) = \begin{cases} -\log \left[c \sin^2 \left(\frac{\tau}{2} \right) \right] & \tau \neq 0 \\ -2 \log \epsilon & \tau = 0 \end{cases}. \quad (4.7)$$

The coefficient c reflects the ambiguity involved in subtracting the divergent terms. Its value is scheme dependent and should be fixed by some renormalization prescription. We choose the value $c = 4$ for later convenience. This arbitrariness was discussed in [8, 9]. The integrals over the zero-modes in eq. (4.5) give a 26-dimensional δ -function so that

$$-Z^{(1)}(k) = KT(k)\epsilon^{k^2-1} \quad (4.8)$$

and we can identify

$$T_R(k) \equiv T(k)\epsilon^{k^2-1} = -\frac{Z^{(1)}(k)}{K}. \quad (4.9)$$

This equation provides the renormalized coupling T_R in terms of the bare coupling T to the lowest order in perturbation theory. $1 - k^2$ is the anomalous dimension of the tachyon field. The second order term in T is given by

$$Z^{(2)}(k) = K \int_0^{2\pi} \frac{d\tau_1}{4\pi} \frac{d\tau_2}{2\pi} \int dk_1 dk_2 T(k_1) T(k_2) \left\langle e^{ik_1 X(\tau_1)} e^{ik_2 X(\tau_2)} e^{-ik\hat{X}} \right\rangle. \quad (4.10)$$

Again in (4.10) the integral over the zero modes \hat{X}^μ gives just a 26-dimensional δ -function, $\delta(k - k_1 - k_2)$, and we can perform the integral over the non-zero modes of $X(\tau)$ to get

$$Z^{(2)}(k) = K \int_0^{2\pi} \frac{d\tau_1}{4\pi} \frac{d\tau_2}{2\pi} \int dk_1 dk_2 (2\pi)^D \delta(k - k_1 - k_2) T(k_1) T(k_2) \times \\ \times \exp \left[-\frac{1}{2} (k_1^2 + k_2^2) G(0) - k_1 k_2 G(\tau_1 - \tau_2) \right]. \quad (4.11)$$

The integral in (4.11) becomes

$$\begin{aligned}
 Z^{(2)}(k) &= K \int dk_1 dk_2 (2\pi)^D \delta(k - k_1 - k_2) \epsilon^{k_1^2 + k_2^2 - 2} T(k_1) T(k_2) \times \\
 &\quad \times \int_0^{2\pi} \frac{d\tau_1}{4\pi} \frac{d\tau_2}{2\pi} \left[4 \sin^2 \left(\frac{\tau_1 - \tau_2}{2} \right) \right]^{k_1 k_2}. \quad (4.12)
 \end{aligned}$$

The integral over the relative variable $x = (\tau_1 - \tau_2)/2$ does not need regularization, it converges when $1 + 2k_1 k_2 > 0$, providing the result

$$Z^{(2)}(k) = \frac{K}{2} \int dk_1 dk_2 (2\pi)^D \delta(k - k_1 - k_2) \epsilon^{k_1^2 + k_2^2 - 2} T(k_1) T(k_2) \frac{\Gamma(1 + 2k_1 k_2)}{\Gamma^2(1 + k_1 k_2)}. \quad (4.13)$$

The integrand in (4.13) can be analytically continued also to the region where $1 + 2k_1 k_2 < 0$, so that the integral can be performed.

To the second order in perturbation theory the renormalized coupling in terms of the bare coupling reads

$$\begin{aligned}
 T_R(k) &= - \frac{Z^{(1)}(k) + Z^{(2)}(k)}{K} \\
 &= \epsilon^{k^2 - 1} \left[T(k) - \frac{1}{2} \int dk_1 dk_2 (2\pi)^D \delta(k - k_1 - k_2) \times \right. \\
 &\quad \left. \times T(k_1) T(k_2) \epsilon^{-(1 + 2k_1 k_2)} \frac{\Gamma(1 + 2k_1 k_2)}{\Gamma^2(1 + k_1 k_2)} \right]. \quad (4.14)
 \end{aligned}$$

The third order contribution to the partition function is given by

$$Z^{(3)}(k) = - \frac{K}{3!} \int dk_1 dk_2 dk_3 (2\pi)^D \delta \left(k - \sum_{i=1}^3 k_i \right) \epsilon^{\sum_{i=1}^3 k_i^2 - 3} T(k_1) T(k_2) T(k_3) I(k_1, k_2, k_3), \quad (4.15)$$

where $I(k_1, k_2, k_3)$ is the integral

$$\begin{aligned}
 I(k_1, k_2, k_3) &= \frac{2^{2k_1 k_2 + 2k_2 k_3 + 2k_1 k_3}}{(2\pi)^3} \int_0^{2\pi} d\tau_1 d\tau_2 d\tau_3 \left[\sin^2 \left(\frac{\tau_1 - \tau_2}{2} \right) \right]^{k_1 k_2} \times \\
 &\quad \times \left[\sin^2 \left(\frac{\tau_2 - \tau_3}{2} \right) \right]^{k_2 k_3} \left[\sin^2 \left(\frac{\tau_1 - \tau_3}{2} \right) \right]^{k_1 k_3}. \quad (4.16)
 \end{aligned}$$

The complete computation of $I(k_1, k_2, k_3)$ will be given in appendix A. The result is given by the completely symmetric formula

$$I(a_1, a_2, a_3) = \frac{\Gamma(1 + a_1 + a_2 + a_3) \Gamma(1 + 2a_1) \Gamma(1 + 2a_2) \Gamma(1 + 2a_3)}{\Gamma(1 + a_1) \Gamma(1 + a_2) \Gamma(1 + a_3) \Gamma(1 + a_1 + a_2) \Gamma(1 + a_2 + a_3) \Gamma(1 + a_1 + a_3)}, \quad (4.17)$$

where we have set $a_1 = k_1 k_2$, $a_2 = k_2 k_3$ and $a_3 = k_1 k_3$. The integral (4.16) converges when $1 + a_1 + a_2 + a_3 > 0$, but its result (4.17) can be analytically continued also outside this convergence region. The result (4.17) is in agreement with the one obtained, with a different procedure, in [24] but does not coincide with the one provided in the appendix of

ref. [7]. Up to the third order in powers of T and to all orders in k_i the relation between the bare and the renormalized couplings reads

$$\begin{aligned}
 T_R(k) &= -\frac{Z^{(1)}(k) + Z^{(2)}(k) + Z^{(3)}(k)}{K} \\
 &= \epsilon^{k^2-1} \left[T(k) - \frac{1}{2} \int dk_1 dk_2 (2\pi)^D \delta \left(k - \sum_{i=1}^3 k_i \right) T(k_1) T(k_2) \epsilon^{-(1+2k_1 k_2)} \times \right. \\
 &\quad \times \frac{\Gamma(1+2k_1 k_2)}{\Gamma^2(1+k_1 k_2)} + \int dk_1 dk_2 dk_3 \frac{(2\pi)^D}{3!} \delta \left(k - \sum_{i=1}^3 k_i \right) \times \\
 &\quad \left. \times T(k_1) T(k_2) T(k_3) \epsilon^{-2(1+\sum_{i<j} k_i k_j)} I(k_1, k_2, k_3) \right]. \tag{4.18}
 \end{aligned}$$

In section 6 we shall use this expression to construct the non-linear β -function.

The renormalized tachyon field can be constructed to all powers of the bare tachyon field in the case in which the tachyon profile appearing in (4.2) is a slowly varying function of X^μ . In this case one can consider an expansion of (4.2) in powers of derivatives of T . To this purpose consider the n -th term in the expansion of (4.2) in powers of $T(X(\tau))$, $Z^{(n)}(k)$. Taking the Fourier transform of the tachyon field and performing all the contractions of the $X(\tau_i)$ fields, for $Z^{(n)}(k)$ we get

$$\begin{aligned}
 Z^{(n)}(k) &= K \frac{(-1)^n}{n!} \epsilon^{-n} \int \prod_{i=1}^n dk_i T(k_i) \int_0^{2\pi} \prod_{i=1}^n \left(\frac{d\tau_i}{2\pi} \right) \times \\
 &\quad \times e^{-\sum_{i=1}^n \frac{k_i^2}{2} G(0) - \sum_{i<j} k_i k_j G(\tau_i - \tau_j)} \delta \left(k - \sum_{i=1}^n k_i \right). \tag{4.19}
 \end{aligned}$$

Note that with our regularization prescription the dependence on the cut-off in (4.19) comes only from the zero distance propagator $G(0)$ and from the explicit scale dependence of the tachyon field. If the tachyon profile is a slowly varying function of X^μ we can expand inside the integrand of (4.19) in powers of the momenta k_i . The leading and next to leading terms in this expansion read

$$\begin{aligned}
 Z^{(n)}(k) &= K \frac{(-1)^n}{n!} \prod_{i=1}^n \int dk_i \delta \left(k - \sum_{i=1}^n k_i \right) \epsilon^{-n} \prod_{i=1}^n T(k_i) \times \\
 &\quad \times \left(1 + \sum_{i=1}^n k_i^2 \log \epsilon + \sum_{i<j} k_i k_j \log \frac{c}{4} \right), \tag{4.20}
 \end{aligned}$$

where the last term comes from the integral over a couple of τ variables of the propagator $G(\tau_i - \tau_j)$, the other integrations over τ_k $k \neq i, j$ being trivial. Here we have kept explicit the ambiguity c appearing in the propagator (4.7) to show that the result greatly simplifies with the choice $c = 4$. Unless otherwise stated, we shall adopt this choice throughout the paper. As before, the renormalized tachyon field $T_R(k)$ can be obtained from (4.20) by summing over n from 1 to ∞ , changing sign and dividing by K . Taking the Fourier

transform of $T_R(k)$ with $c = 4$, to all orders in the bare tachyon field and to the leading order in derivatives, we get the renormalized tachyon field $T_R(X)$

$$T_R(X) = 1 - \exp \left\{ -\frac{1}{\epsilon} [T(X) - \Delta T(X) \log \epsilon] \right\} \quad (4.21)$$

where Δ is the laplacian. Again in section 6 we shall use this expression to compute the non-linear tachyon β -function.

From eqs. (4.9), (4.14), (4.18), (4.21) it is clear that the general relation between the renormalized tachyon field $T_R(X)$ and the partition function $Z \equiv Z(k = 0)$ is simply

$$Z = K \int d^{26} X [1 - T_R(X)]. \quad (4.22)$$

This expression is true also when other couplings are present. T_R in this case would be a non linear function also of the other bare couplings but its relation with the partition function of the theory would always be given by (4.22). We shall prove eq. (4.22) in the next section.

5. Background-field method

The partition function of the boundary theory on the disk in general is given by

$$Z = K \int [dX_\mu] e^{-(S_0[X] + \int_0^{2\pi} \frac{d\tau}{2\pi} \mathcal{V}[X(\tau)]),} \quad (5.1)$$

where $S_0 = \int d\tau X^\mu |i\partial_\tau| X^\mu$ and $\mathcal{V}[X(\tau)]$ is given in (2.2). Our goal is to determine the relationship between the renormalized and the bare couplings of the one-dimensional field theory. To this purpose we shall make use of the background field method [23]. We expand the fields X^μ around a classical background X_0^μ which satisfies the equations of motion and which varies slowly compared to the cut-off scale,

$$X^\mu = X_0^\mu + Y^\mu.$$

The effective action is $S_{\text{eff}}[X_0] = -\log Z[X_0]$ and the aim of the renormalization process is to rewrite the local terms of $S_{\text{eff}}[X_0]$ in terms of renormalized couplings in such a way that $S_{\text{eff}}[X_0]$ has the same form of the original action

$$S_{\text{eff}}[X_0] \Big|_{\text{local}} = S_0[X_0] + \int_0^{2\pi} \frac{d\tau}{2\pi} \mathcal{V}_R[X_0(\tau)]. \quad (5.2)$$

$Z[X_0]$ can be conveniently calculated in powers of the boundary interaction \mathcal{V} . The first order for example reads, up to the multiplicative constant K ,

$$- \int_0^{2\pi} \frac{d\tau}{2\pi} \int dk e^{ikX_0} \langle [T(k) + iA_\mu(k) \partial_\tau (X_0^\mu + Y^\mu) + B_{\mu\nu}(k) \partial_\tau (X_0^\mu + Y^\mu) \partial_\tau (X_0^\nu + Y^\nu) + C_\mu(k) \partial_\tau^2 (X_0^\mu + Y^\mu) + \dots] e^{ikY} \rangle. \quad (5.3)$$

The renormalized couplings $T_R(k)$ will be given by the opposite of the coefficient of the term in (5.3) that does not contain X_0 derivatives. Analogously, the renormalized $A_\mu^R(k)$ will be determined by the coefficient of $\partial_\tau X_0^\mu$, $B_{\mu\nu}^R(k)$ by the coefficient of $\partial_\tau X_0^\mu \partial_\tau X_0^\nu$ and so on. The second order term in the expansion of $Z[X_0]$ is

$$\int_0^{2\pi} \frac{d\tau_1}{2\pi} \int_0^{2\pi} \frac{d\tau_2}{4\pi} \int dk_1 dk_2 e^{ik_1 X_0(\tau_1) + ik_2 X_0(\tau_2)} \times \\ \times \langle e^{ik_1 Y(\tau_1) + ik_2 Y(\tau_2)} [T(k_1) + iA_\mu(k_1) \partial_{\tau_1} (X_0^\mu + Y^\mu) + \dots] \times \\ \times [T(k_2) + iA_\nu(k_2) \partial_{\tau_2} (X_0^\nu + Y^\nu) + \dots] \rangle. \quad (5.4)$$

An expansion of the background field X_0 in powers of its derivatives is required to determine the coefficients of $1, \partial_\tau X_0^\mu, \partial_\tau X_0^\mu \partial_\tau X_0^\nu, \dots$,

$$X_0(\tau_2) = X_0(\tau_1) + (\tau_2 - \tau_1) \partial_{\tau_1} X_0(\tau_1) + \dots \quad (5.5)$$

If we are interested in renormalization of couplings of the form $\exp[ikX_0]$, namely in the renormalized tachyon field $T_R(k)$, we can disregard the terms in (5.5), (5.4) involving derivatives acting on X_0 . For example, at the second order, the only non-vanishing terms in T and A_μ contributing to T_R are

$$T_R(k) = - \int dk_1 \int dk_2 \delta(k - k_1 - k_2) \int_0^{2\pi} \frac{d\tau_2}{4\pi} \times \\ \times \langle e^{ik_1 Y(\tau_1) + ik_2 Y(\tau_2)} [T(k_1)T(k_2) + A_\mu(k_1)A_\nu(k_2) \partial_{\tau_1} Y^\mu \partial_{\tau_2} Y^\nu + \dots] \rangle, \quad (5.6)$$

where the correlator does not depend on τ_1 since the propagator (4.7) of $X(\tau)$ and its derivatives are periodic functions on the unit circle. It is not difficult to see that $T_R(k)$ in (5.6) coincides with the opposite of the second order term in the expansion of the partition function

$$Z(k) = \int [dY_\mu] e^{-(S_0[Y] + \int_0^{2\pi} \frac{d\tau}{2\pi} \mathcal{V}[Y(\tau)] - ik\hat{Y}} \quad (5.7)$$

in powers of the couplings. Here k is a constant source for the zero mode of the Y^μ field, \hat{Y}^μ (4.3). Such a constant source will just provide the δ -function in (5.6) that imposes the energy-momentum conservation. This will be true at any order in the expansion in powers of the coupling fields. Therefore, to all orders in whatever coupling, the expression for the renormalized tachyon field $T_R(X)$ is related to the partition function $Z = Z(k=0)$ precisely by (4.22), which is the relation that we wanted to prove. Note that T_R depends not only on the bare tachyon field but also on the other coupling fields (in particular T_R will exist also if one starts from a boundary interaction that does not contain the bare tachyon). As a consequence, the tachyon β function will contain for example also the gauge field [30], and this is as it should be, since the solution of the equation $\beta^T = 0$ will then describe the scattering of a tachyon by other excitations (e.g. from (5.6) by two vector fields).

6. β -function

In this section we shall perform a calculation of the non-linear tachyon β -function. The resulting expression will then be used to derive the Witten-Shatashvili action (1.4), (4.1). Following [23], the most general RG equations for a set of couplings g^i can be written as

$$\beta^i \equiv \frac{dg^i}{dt} = \lambda_i g^i + \alpha_{jk}^i g^j g^k + \gamma_{jkl}^i g^j g^k g^l + \dots, \quad (6.1)$$

where the scale t is $t = -\log \epsilon$, λ_i are the anomalous dimensions corresponding to the couplings g^i and there is no summation in the first term on the right-hand side. This equation has the solution

$$g^i(t) = e^{\lambda_i t} g^i(0) + \left[e^{(\lambda_j + \lambda_k)t} - e^{\lambda_i t} \right] \frac{\alpha_{jk}^i}{\lambda_j + \lambda_k - \lambda_i} g^j(0) g^k(0) + b_{jkl}^i(t) g^j(0) g^k(0) g^l(0) + \dots, \quad (6.2)$$

where $g^i(0)$ are the bare couplings and

$$b_{jkl}^i(t) g^j(0) g^k(0) g^l(0) = \left[\left(\frac{2\alpha_{jm}^i \alpha_{kl}^m}{\lambda_j + \lambda_m - \lambda_i} - \gamma_{jkl}^i \right) \frac{e^{\lambda_i t}}{\lambda_j + \lambda_k + \lambda_l - \lambda_i} + \left(\frac{2\alpha_{jm}^i \alpha_{kl}^m}{\lambda_k + \lambda_l - \lambda_m} + \gamma_{jkl}^i \right) \frac{e^{(\lambda_j + \lambda_k + \lambda_l)t}}{\lambda_j + \lambda_k + \lambda_l - \lambda_i} - \frac{2\alpha_{jm}^i \alpha_{kl}^m}{(\lambda_j + \lambda_m - \lambda_i)(\lambda_k + \lambda_l - \lambda_m)} e^{(\lambda_j + \lambda_k)t} \right] g^j(0) g^k(0) g^l(0). \quad (6.3)$$

Let us now consider the case of interest for this paper: open strings propagating in a tachyon background. In this case the coupling g^i is the tachyon field $T(k)$. Then $\lambda_i = 1 - k^2$ and $\lambda_j = 1 - k_j^2$. Comparing the general solution (6.2) with eq. (4.18) derived in the previous section, we will be able to identify the renormalized tachyon field in terms of the bare field up to the third order in powers of the field and to all orders in its derivatives. In the second order term of (4.18) the coefficient proportional to $e^{\lambda_i t} = \epsilon^{1-k^2}$ appearing in (6.1) is absent. This is due to the fact that the convergence condition for the integral (4.12), $1 + 2k_1 k_2 > 0$, implies that $\lambda_j + \lambda_k > \lambda_i$ so that in the limit $t \rightarrow \infty$ the dominant contribution comes from $e^{(\lambda_j + \lambda_k)t}$. From similar arguments, the first and the second terms of the right-hand side of (6.3) are negligible compared to the second term, due to the convergence conditions for the integral $I(k_1, k_2, k_3)$ computed in the previous section. This is a general feature of our renormalization procedure. At the n -th order in the bare coupling in the expansion (6.2), the renormalized coupling will contain only the term of the form

$$e^{t \sum_{k=1}^n \lambda_k}. \quad (6.4)$$

This is due to the fact that the integrals over the τ 's do not need an explicit regulator, rather they can be evaluated in a specific region of the k_i variables and then analytically continued. Therefore the only dependence on the cut-off does not come from such integrals but from the propagators (4.7) evaluated at zero distance.

Comparing our result for the renormalized tachyon field (4.18) with the general expressions (6.2), (6.3), for the coefficients in the expansion (6.1) we find

$$\begin{aligned} \alpha_{jk}^i &= -\frac{1}{2} \frac{\Gamma(2 + 2k_j k_k)}{\Gamma^2(1 + k_j k_k)} \delta(k - k_j - k_k) \\ \gamma_{jkl}^i &= \frac{1}{3!} \int dk_j dk_k dk_l \delta(k - k_j - k_k - k_l) \\ &\quad \times \left[2(1 + k_j k_k + k_j k_l + k_k k_l) I(k_j, k_k, k_l) - \right. \\ &\quad \left. - \left(\frac{\Gamma(2 + 2k_j k_k + 2k_j k_l) \Gamma(1 + 2k_k k_l)}{\Gamma^2(1 + k_j k_k + k_j k_l) \Gamma^2(1 + k_k k_l)} + \text{cycl.} \right) \right], \end{aligned} \quad (6.5)$$

where $I(k_j, k_k, k_l)$ is given in equation (4.17). The perturbative expression for the β -function up to the third order in the tachyon field obtained using this procedure therefore is

$$\begin{aligned} \beta^T(k) &= (1 - k^2) T_R(k) - \frac{1}{2} \int dk_1 dk_2 (2\pi)^D \delta(k - k_1 - k_2) T_R(k_1) T_R(k_2) \frac{\Gamma(2 + 2k_1 k_2)}{\Gamma^2(1 + k_1 k_2)} + \\ &\quad + \frac{1}{3!} \int dk_1 dk_2 dk_3 (2\pi)^D \delta(k - k_1 - k_2 - k_3) T_R(k_1) T_R(k_2) T_R(k_3) \times \\ &\quad \times \left[2(1 + k_1 k_2 + k_1 k_3 + k_2 k_3) I(k_1, k_2, k_3) - \right. \\ &\quad \left. - \left(\frac{\Gamma(2 + 2k_1 k_2 + 2k_1 k_3) \Gamma(1 + 2k_2 k_3)}{\Gamma^2(1 + k_1 k_2 + k_1 k_3) \Gamma^2(1 + k_2 k_3)} + \text{cycl.} \right) \right]. \end{aligned} \quad (6.6)$$

We have thus succeeded in deriving a β -function for tachyon backgrounds which do not satisfy the linearized on-shell condition. Exactly the same result can be obtained by taking the derivative of (4.18) (or of the opposite of $Z(k)$) with respect to the logarithm of the cut-off $-\log \epsilon$. The result obtained in this way must then be expressed in terms of the renormalized field by inverting (4.18) and it coincides with (6.6).

It is interesting to note that all the known conformal tachyon profiles, like e^{iX^0} or $\cos X^i$ where i is a space index, are solutions of the equation $\beta^T(X) = 0$, where $\beta^T(X)$ is the Fourier transform of (6.6). These solutions and perturbations around them have been recently used to construct tachyon effective actions around the on-shellness [31, 18, 19, 20, 22] and to study the problem of the rolling tachyon [13, 14, 15, 16, 17, 32, 21].

That the non-linear β -function (6.6) is the correct one can be shown by solving the $\beta^T(k) = 0$ equation perturbatively. The solution of this equation will generate the correct scattering amplitudes of open string theory [23]. This in turn will show the validity of the general formula (4.22). To the lowest order the equation is $(1 - k^2) T_0(k) = 0$, so that the solution $T_0(k)$ satisfies the linearized on-shell condition. By writing $T(k) = T_0(k) + T_1(k)$ and substituting into the equation $\beta_T(k) = 0$, to the next order we find

$$T_1(k) = \frac{1}{2} \int dk_1 dk_2 (2\pi)^D \delta(k - k_1 - k_2) T_0(k_1) T_0(k_2) \frac{\Gamma(k^2)}{(1 - k^2) \Gamma^2(k^2/2)}. \quad (6.7)$$

The presence of the couplings T_0 in (6.7) sets two of the three k_i on-shell. To pick out the propagator pole corresponding to the third k we set it on-shell too. The scattering amplitude for three on-shell tachyons is given by the residue of the pole and is $1/2\pi$ with our normalization.

The calculation at the next order proceed in a similar fashion. One sets $T(k) = T_0(k) + T_1(k) + T_2(k)$ and finds

$$\begin{aligned}
 T_2(k) = & -\frac{(2\pi)^D}{3!(1-k^2)} \int dk_1 dk_2 dk_3 \delta(k - k_1 - k_2 - k_3) T_0(k_1) T_0(k_2) T_0(k_3) I(k_1, k_2, k_3) \times \\
 & \times \left\{ 2 \left(1 + \sum_{i < j} k_i k_j \right) I(k_1, k_2, k_3) - \left[\frac{\Gamma(2 + 2k_1 k_2 + 2k_1 k_3) \Gamma(1 + 2k_2 k_3)}{\Gamma^2(1 + k_1 k_2 + k_1 k_3) \Gamma^2(1 + k_2 k_3)} + \text{cycl.} \right] - \right. \\
 & \left. - \left[\frac{\Gamma(2 + 2k_1 k_2 + 2k_1 k_3) \Gamma(2 + 2k_2 k_3)}{\Gamma^2(1 + k_1 k_2 + k_1 k_3) \Gamma^2(1 + k_2 k_3) [1 - (k_2 + k_3)^2]} + \text{cycl.} \right] \right\}. \quad (6.8)
 \end{aligned}$$

When all the tachyons are on-shell, the last two terms on eq. (6.8) cancel and, as it should be for consistency, the residue of the pole in k is the scattering amplitude of four on-shell tachyons. It is given by

$$\frac{\Gamma(1 + 2k_1 k_2) \Gamma(1 + 2k_2 k_3) \Gamma(1 + 2k_1 k_3)}{\Gamma(1 + k_1 k_2) \Gamma(1 + k_2 k_3) \Gamma(1 + k_1 k_3) \Gamma(1 + k_1 k_2 + k_2 k_3) \Gamma(1 + k_2 k_3 + k_1 k_3) \Gamma(1 + k_1 k_2 + k_1 k_3)}, \quad (6.9)$$

where the on-shell condition is $1 + k_1 k_2 + k_2 k_3 + k_1 k_3 = 0$. By means of the on-shell condition, from the above expression, we recover, up to a normalization constant, the Veneziano amplitude, the scattering amplitude of four on-shell tachyons. Eq. (6.9) in fact becomes

$$\begin{aligned}
 \frac{1}{\pi^3} \Gamma(1 + 2k_1 k_2) \Gamma(1 + 2k_2 k_3) \Gamma(1 + 2k_1 k_3) \sin(\pi k_1 k_2) \sin(\pi k_2 k_3) \sin(\pi k_1 k_3) = \\
 = \frac{1}{(2\pi)^2} [B(1 + 2k_1 k_2, 1 + 2k_2 k_3) + \text{cycl.}], \quad (6.10)
 \end{aligned}$$

where $B(x, y)$ is the Euler beta function. The expression between square brackets is just the Veneziano amplitude. The ambiguity c appearing in the propagator (4.7) could be kept undetermined throughout the calculations of the scattering amplitudes. It is not difficult to see that this would just consistently change the normalization of the on-shell amplitudes.

For tachyon profiles $T_R(k)$ supported over near on-shell momentum $k^2 \simeq 1$, the equation of motion $\beta^T = 0$ with β^T given in (6.6) becomes

$$\begin{aligned}
 \beta^T(k) = & (1 - k^2) T_R(k) - \frac{(2\pi)^D}{2\pi} \int dk_1 dk_2 \delta(k - k_1 - k_2) T_R(k_1) T_R(k_2) + \\
 & + \frac{(2\pi)^D}{3!(2\pi)^2} \int dk_1 dk_2 dk_3 \delta(k - k_1 - k_2 - k_3) T_R(k_1) T_R(k_2) T_R(k_3) \times \\
 & \times \{ [B(1 + 2k_1 k_2, 1 + 2k_2 k_3) + \text{cycl.}] + 2\pi \tan(\pi k_1 k_2) \tan(\pi k_1 k_3) \tan(\pi k_2 k_3) \} = 0. \quad (6.11)
 \end{aligned}$$

The coefficients of the quadratic and cubic terms in (6.11) are symmetric with respect to all the k_i and k when these are on the mass-shell. Thus (6.11) can be derived as the equation of motion of an effective action. Such effective action for near on-shell tachyons up to the fourth order in powers of the tachyon fields can be derived from the results of the cubic string field theory. In [33] it was shown that the cubic SFT reproduces the Veneziano

amplitude with great accuracy already at level $L = 50$. The tachyon effective action arising from the cubic string field theory for near on shell tachyon profiles $\Phi(k)$ therefore reads²

$$S_C = 2\pi^2 T_{25} (2\pi)^D \left\{ -\frac{1}{2} \int dk \Phi(k) \Phi(-k) (1 - k^2) + \frac{1}{3} \int \prod_{i=1}^3 dk_i \Phi(k_i) \delta\left(\sum_{i=1}^3 k_i\right) + \frac{1}{4!} \int \prod_{i=1}^4 dk_i \Phi(k_i) \delta\left(\sum_{i=1}^4 k_i\right) [B(1+2k_1k_2, 1+2k_2k_3) + \text{cycl.}] \right\}, \quad (6.12)$$

where the tachyon momenta in the fourth order term satisfy

$$k_1 = (0, 1, 0, 0, \dots, 0) \quad k_2 = (0, \sin \theta, \cos \theta, 0, \dots, 0) \quad (6.13)$$

$$k_3 = (0, -1, 0, 0, \dots, 0) \quad k_4 = (0, -\sin \theta, -\cos \theta, 0, \dots, 0). \quad (6.14)$$

Since the Veneziano amplitude is completely symmetric in the four momenta k_i , it is not difficult to see that the equation of motion deriving from (6.12) becomes precisely (6.11) once the simple field rescaling $T = 2\pi\Phi$ is performed. Thus the cubic string field theory for almost on-shell tachyons reproduces the non-linear $\beta^T = 0$ equation of motion.

In section 4 we also derived the renormalized tachyon field for the case of a slowly varying tachyon profile, to all orders in the bare field and to the leading order in derivatives, eq. (4.21). From this we can easily compute the corresponding β function. The task in this case is much simpler, as we just need to take the derivative of (4.21) with respect to $-\log \epsilon$

$$\beta(X) = \frac{\partial T_R(X)}{\partial(-\log \epsilon)} = \frac{1}{\epsilon} \exp\left(-\frac{T(X)}{\epsilon}\right) \left\{ T(X) + \Delta T(X) \left[1 - \left(1 - \frac{T(X)}{\epsilon}\right) \log \epsilon \right] \right\}. \quad (6.15)$$

Then we have to invert the relation (4.21) between T_R and T . To the leading order in derivatives one has

$$T(X) = -\epsilon \{ [1 + (\log \epsilon)\Delta] \log(1 - T_R(X)) \}, \quad (6.16)$$

from which it is clear that the admissible range for T_R is $-\infty \leq T_R \leq 1$. Plugging (6.16) into (6.15) we get the non-linear tachyon β -function to all powers of the renormalized tachyon and to the leading order in its derivatives

$$\beta^T(X) = (1 - T_R(X)) [-\log(1 - T_R(X)) - \Delta \log(1 - T_R(X))]. \quad (6.17)$$

$\beta^T(X) = 0$ is the tachyon equation of motion for a slowly varying tachyon profile.

Since in our calculations of the non-linear β -function we have always used the same coordinates in the space of string fields, the two results (6.17) and (6.6) should coincide when expanded up to the third power of the field and to the leading order in derivatives, respectively. This is indeed the case and the result in both cases reads

$$\beta^T(X) = \Delta T_R + \partial_\mu T_R \partial_\mu T_R + T_R \partial_\mu T_R \partial_\mu T_R. \quad (6.18)$$

²Here we include also the contribution due to the tachyon field in the quartic term so that this on-shell effective action reproduces exactly the Veneziano amplitude and the correct equation of motion.

It is interesting to compute the β -function also in the case in which the ambiguity constant c appearing in (4.7) is kept undetermined. $T_R(k)$ can be easily obtained as before from (4.20) without fixing $c = 4$. By taking the Fourier transform and by differentiating with respect to $-\log \epsilon$, the β -function expressed in terms of the renormalized tachyon field $T_R(X)$ turns out to be

$$\beta^T(X) = (1 - T_R) \left[-\log(1 - T_R) + \frac{\Delta T_R}{1 - T_R} + \left(1 + \frac{1}{2} \log \frac{c}{4} \right) \frac{\partial_\mu T_R \partial_\mu T_R}{(1 - T_R)^2} \right]. \quad (6.19)$$

In the next section we shall use also this form of the β -function to construct the Witten-Shatashvili action.

7. Witten-Shatashvili action

In this section we shall compute the Witten-Shatashvili action. From the simple expression that relates the partition function to the renormalized tachyon (4.22) it is easy to deduce a simple and universal form for the WS action of the open bosonic string theory

$$S = \left(1 - \int \beta^T \frac{\delta}{\delta T_R} \right) Z[T_R] = K \int d^D X [1 - T_R(X) + \beta^T(X)]. \quad (7.1)$$

This can now be computed in both the cases analyzed in the previous sections. We shall show that the expressions for S that we will obtain are consistent both with the known results on the tachyon potential [7] and with the expected on-shell behavior. Thus a choice of coordinates in the space of couplings in which the tachyon β -function is non-linear allows one to find not only a simple general formula for the WS action, but provides also a space-time tachyon effective action that describes tachyon physics from the far-off shell to the near on-shell regions.

Let us start with the evaluation of (7.1) up to the third order in the expansion of the tachyon field using the non-linear β -function (6.6). A similar computation was done in [7, 24] by means of the linear β -function, $\beta(k) = (1 - k^2) T(k)$. We shall later compare the two results. From the renormalized field (4.18) and the β -function (6.6) we arrive at the following expression for the Witten action

$$S = K \left\{ 1 - \frac{1}{2} \int dk (2\pi)^D T_R(k) T_R(-k) \frac{\Gamma(2 - 2k^2)}{\Gamma^2(1 - k^2)} + \frac{1}{3!} \int dk_1 dk_2 dk_3 (2\pi)^D T_R(k_1) T_R(k_2) T_R(k_3) \delta(k_1 + k_2 + k_3) \times \right. \\ \left. \times \left[2 \left(1 + \sum_{i < j} k_i k_j \right) I(k_1, k_2, k_3) - \left(\frac{\Gamma(1 + 2k_2 k_3) \Gamma(2 + 2k_1 k_2 + 2k_1 k_3)}{\Gamma^2(1 + k_2 k_3) \Gamma^2(1 + k_1 k_2 + k_1 k_3)} + \text{cycl.} \right) \right] \right\}. \quad (7.2)$$

The propagator coming from the quadratic term in (7.2) exhibits the required pole at $k^2 = 1$. There are however also an infinite number of other zeroes and poles. We shall

show that these are due to the metric in the coupling space appearing in (1.3). The equations of motion derived from the action (7.2) are

$$\begin{aligned} \frac{\delta S}{\delta T_R(-k)} &= -K \frac{\Gamma(2-2k^2)}{\Gamma^2(1-k^2)} (2\pi)^D T(k) + \frac{K}{2} \int dk_1 dk' (2\pi)^D \delta(k_1+k'-k) T_R(k_1) T_R(k') \times \\ &\times \left\{ 2(1-k_1k+k_1k'-kk') I(-k, k_1, k') - \frac{\Gamma(1-2kk_1)\Gamma(2-2kk'+2k_1k')}{\Gamma^2(1-kk_1)\Gamma^2(1-kk'+k'k_1)} - \right. \\ &\quad \left. - \frac{\Gamma(1-2kk')\Gamma(2-2kk_1+2k'k_1)}{\Gamma^2(1-kk')\Gamma^2(1-kk_1+k'k_1)} - \frac{\Gamma(1+2k'k_1)\Gamma(2-2kk'-2kk_1)}{\Gamma^2(1+k'k_1)\Gamma^2(1-kk'-kk_1)} \right\}. \end{aligned} \quad (7.3)$$

As we did for the equation $\beta^T = 0$ in the previous section, by solving these equations perturbatively it is possible to recover the scattering amplitudes for three on-shell tachyons. To the lowest order the equation is

$$\frac{\Gamma(2-2k^2)}{\Gamma^2(1-k^2)} T_0(k) = 0. \quad (7.4)$$

At variance with the lowest order solution of $\beta^T = 0$, there are infinite possible solutions of (7.4). We choose the solution for which the tachyon field $T_0(k)$ is on the mass-shell, which corresponds to a consistent string theory background. This choice is also a solution of $\beta^T = 0$ to the lowest order. As we shall show, the other possible zeroes of (7.4) could be interpreted as zeroes of the metric in the space of couplings through eq. (1.3). With such a choice of $T_0(k)$, to the next order we recover the scattering amplitudes for three on-shell tachyons. By writing $T(k) = T_0(k) + T_1(k)$ and substituting it into (7.3) we find

$$\begin{aligned} T_1(k) &= \frac{\Gamma^2(1-k^2)}{2\Gamma(2-2k^2)} \int dk_1 dk' (2\pi)^D \delta(k-k_1-k') T_0(k_1) T_0(k') \times \\ &\times \left\{ 2(1-k_1k+k_1k'-kk') I(-k, k_1, k') - \frac{\Gamma(1-2kk_1)\Gamma(2-2kk'+2k_1k')}{\Gamma^2(1-kk_1)\Gamma^2(1-kk'+k'k_1)} - \right. \\ &\quad \left. - \frac{\Gamma(1-2kk')\Gamma(2-2kk_1+2k'k_1)}{\Gamma^2(1-kk')\Gamma^2(1-kk_1+k'k_1)} - \frac{\Gamma(1+2k'k_1)\Gamma(2-2kk'-2kk_1)}{\Gamma^2(1+k'k_1)\Gamma^2(1-kk'-kk_1)} \right\}. \end{aligned} \quad (7.5)$$

Since the two couplings T_0 satisfy the on-shell condition, k_1 and k' are on-shell. To pick out the propagator pole corresponding to the third k we set it on-shell too. The scattering amplitude for three on-shell tachyons is given again by the residue of the pole and with our normalization is $(2\pi)^{-1}$, in precise agreement with the result obtained in the previous section.

The equations (7.3) must be related to the equation $\beta^T = 0$ through a metric $G_{T(k)T(k')}$ as in (1.3), which in this case becomes

$$\frac{\delta S}{\delta T_R(k)} = - \int dk' G_{T(k)T(k')} \beta^{T(k')}. \quad (7.6)$$

The Witten-Shatashvili formulation of string field theory provides a prescription for the metric $G_{T(k)T(k')}$ which can then be computed explicitly. To the first two orders in powers of T_R , it is given by

$$G_{T(k)T(k')} = K \frac{(2\pi)^D \Gamma(2-2k^2)}{(1-k^2)\Gamma^2(1-k^2)} \delta(k+k') - \frac{K}{2} \int dk_1 (2\pi)^D \delta(k+k'+k_1) \frac{T_R(k_1)}{1-k'^2} \times$$

$$\times \left\{ 2(1 + k_1 k + k_1 k' + k k') I(k_1, k, k') - \frac{\Gamma(1 + 2k k_1) \Gamma(2 + 2k k' + 2k_1 k')}{\Gamma^2(1 + k k_1) \Gamma^2(1 + k k' + k' k_1)} - \frac{\Gamma(1 + 2k k') \Gamma(2 + 2k k_1 + 2k' k_1)}{\Gamma^2(1 + k k') \Gamma^2(1 + k k_1 + k' k_1)} - \frac{\Gamma(1 + 2k' k_1) \Gamma(2 + 2k k' + 2k k_1)}{\Gamma^2(1 + k' k_1) \Gamma^2(1 + k k' + k k_1)} - \frac{\Gamma(2 + 2k' k_1) \Gamma(2 + 2k k' + 2k k_1)}{\Gamma^2(1 + k' k_1) \Gamma^2(1 + k k' + k k_1)(1 + k k' + k k_1)} \right\}. \quad (7.7)$$

The first term in this metric coincides with (2.9) for a conformal primary given by the tachyon vertex. From (7.7) it is possible to see that the infinite number of zeroes and poles that the second order term in eq. (7.2) exhibits at $k^2 = 1 + n$ and $k^2 = 3/2 + n$, respectively, is in fact due to the metric. This is true except for the zero corresponding to the tachyon mass-shell $k^2 = 1$. In fact the metric (7.7) is regular for $k^2 = 1$. This indicates that the kinetic term in eq. (7.2) exhibits the required zero at the tachyon mass-shell and the metric (7.7) can be made responsible for the other extra zeroes and poles. If these zeroes and poles are just an artifact of the expansion in powers of T , it is an open question. It would be interesting to consider for example an expansion around $k^2 = 1 + n$ to all orders in T and check if in this case one would still find that the kinetic term exhibits a zero at $k^2 = 1 + n$.

Let us turn now to the cubic term in eq. (7.2). If one or two tachyons are on-shell, then the cubic term vanishes. This means that any exchange diagram involving the cubic term vanishes [24]. When all the three tachyons are on-shell, the scattering amplitude for three on-shell tachyons should arise directly as the coefficient of the cubic term. However, the cubic term in (7.2) is ill-defined on shell. Nonetheless, with the most obvious regularization (i.e. by going on-shell symmetrically by giving to the three tachyons an identical small mass m , $k_i^2 = 1 + m^2$ and then by taking the $m \rightarrow 0$ limit) one gets a finite result for the scattering amplitude [24]. Recalling the first of eqs. (6.5) we conclude that this scattering amplitude is $(2\pi)^{-1}$ with our normalization. Also the cubic term in (7.2) has a sequence of poles at finite distances from the tachyon mass-shell. This is related to the fact that the set of couplings that we have taken into account is not complete. If we get far enough from the tachyon mass-shell, we run into the poles due to all the other string states which have not been subtracted.

In the next section we shall compare (7.2) with the corresponding action derived from the cubic string field theory. Here we would like to show that, by means of a field redefinition, (7.2) can be rewritten in the form of the WS action obtained from a linear β -function [24], but that this field redefinition becomes singular on-shell. The partition function up to the third order in the bare tachyon field is again given by

$$\begin{aligned} Z(k) = & K\delta(k) - K\epsilon^{k^2-1} \times \\ & \times \left[T(k) - \frac{1}{2} \int dk_1 dk_2 (2\pi)^D \delta\left(k - k_1 - k_2\right) \epsilon^{-(1+2k_1 k_2)} T(k_1) T(k_2) \frac{\Gamma(1 + 2k_1 k_2)}{\Gamma^2(1 + k_1 k_2)} + \right. \\ & + \frac{1}{3!} \int dk_1 dk_2 dk_3 (2\pi)^D \delta\left(k - \sum_{i=1}^3 k_i\right) \times \\ & \left. \times \epsilon^{-2(1+\sum_{i<j} k_i k_j)} T(k_1) T(k_2) T(k_3) I(k_1, k_2, k_3) \right], \quad (7.8) \end{aligned}$$

where we have used (4.18). If instead of following the general procedure of ref. [23] one renormalizes the theory simply by normal ordering, the β -function turns out to be linear. Thus the renormalized field to all orders in the bare field would just be

$$\phi_R(k) = T(k)\epsilon^{k^2-1}, \quad (7.9)$$

so that $\beta(k) = (1 - k^2)\phi_R(k)$. The WS action with a linear β -function up to the third order in the tachyon field then reads

$$\begin{aligned} S_L = K \left\{ 1 - \frac{1}{2} \int dk (2\pi)^D \phi_R(k) \phi_R(-k) \frac{\Gamma(2 - 2k^2)}{\Gamma^2(1 - k^2)} + \right. \\ \left. + \frac{1}{3!} \int dk_1 dk_2 dk_3 (2\pi)^D \phi_R(k_1) \phi_R(k_2) \phi_R(k_3) \delta\left(\sum_{i=1}^3 k_i\right) 2 \left(1 + \sum_{i < j=2}^3 k_i k_j\right) \times \right. \\ \left. \times I(k_1, k_2, k_3) \right\} \quad (7.10) \end{aligned}$$

in agreement with what found in [24]. If we assume that the fields ϕ_R and T_R are related as follows

$$\phi_R(k) = T_R(k) + \int dk_1 f(k, k_1) T_R(k_1) T_R(k - k_1) + \dots, \quad (7.11)$$

by comparing the cubic terms in (7.2) and (7.10) one finds

$$\begin{aligned} \left[f(k_2 + k_3, k_2) \frac{\Gamma(2 + 2k_1 k_2 + 2k_1 k_3)}{\Gamma^2(1 + k_1 k_2 + k_1 k_3)} + \text{cycl.} \right] = \\ = \frac{1}{2} \left[\frac{\Gamma(1 + 2k_2 k_3)}{\Gamma^2(1 + k_2 k_3)} \frac{\Gamma(2 + 2k_1 k_2 + 2k_1 k_3)}{\Gamma^2(1 + k_1 k_2 + k_1 k_3)} + \text{cycl.} \right], \quad (7.12) \end{aligned}$$

so that the solution for f is $f(k_1 + k_2, k_1) = \Gamma(1 + 2k_1 k_2) / (2\Gamma^2(1 + k_1 k_2))$ and the field redefinition becomes

$$\phi_R(k) = T_R(k) + \int dk_1 dk_2 \frac{\Gamma(1 + 2k_1 k_2)}{2\Gamma^2(1 + k_1 k_2)} T_R(k_1) T_R(k_2) \delta(k - k_1 - k_2). \quad (7.13)$$

It is not difficult to see that if we evaluate this relation when the three tachyon fields are on-shell it becomes singular since $f(k, k_1)$ has a pole. This is in agreement with the Poincaré-Dulac theorem [25] used in [5] to prove that when the resonant condition (2.11) holds, namely near the on-shellness, the β -function has to be non-linear. We showed in fact that the field redefinition that gives from S the WS action constructed in terms of a linear β -function, S_L , becomes singular on-shell.

Let us now turn to the WS action computed in an expansion to the leading order in derivatives and to all orders in the powers of the tachyon fields. If we keep the renormalization ambiguity c undetermined, the β -function is given in (6.19). Using (1.4), S then reads

$$S = K \int dX (1 - T_R) \left[1 - \log(1 - T_R) + \left(1 + \frac{1}{2} \log \frac{c}{4}\right) \frac{\partial_\mu T_R \partial_\mu T_R}{(1 - T_R)^2} \right], \quad (7.14)$$

where $-\infty \leq T_R \leq 1$. With the field redefinition

$$1 - T_R = e^{-\tilde{T}} \tag{7.15}$$

S becomes

$$S = K \int dX e^{-\tilde{T}} \left[\left(1 + \frac{1}{2} \log \frac{c}{4} \right) \partial_\mu \tilde{T} \partial_\mu \tilde{T} + 1 + \tilde{T} \right], \tag{7.16}$$

which for $c = 4$ coincides with the space-time tachyon action found in [6, 7]. In particular we shall show in the next section that K coincides with the tension of the D25-brane, $K = T_{25}$, in agreement with the results of ref. [7]. It is not difficult to show that (7.16) can be rewritten, by means of a field redefinition, in the form found in [8] where the renormalization ambiguity was also discussed.

Note that (7.15) is the coordinate transformation in the coupling space that leads, for $c = 4$, from the non-linear β -function (6.17) to the linear beta function $\beta^T = (1 + \Delta)T$. The β -function in fact is a covariant vector in the coupling space and as such it transforms.

We have left the ambiguity c in (7.16) undetermined because we want to show that it is possible to fix c in such a way that the equation of motion deriving from (7.16) coincides with the equation $\beta^T = 0$ with β^T given in (6.19). In fact, in terms of the coordinates (7.15), this equation reads

$$\beta^{\tilde{T}} = \tilde{T} + \Delta \tilde{T} + \frac{1}{2} \log \frac{c}{4} \partial_\mu \tilde{T} \partial_\mu \tilde{T} = 0. \tag{7.17}$$

where we have kept into account that $\beta^{\tilde{T}}$ transforms like a covariant vector in the space of worldsheet theories. Choosing $\log(c/4) = -1$, eq. (7.17) becomes the equation of motion of the action (7.16). This is important because if we find finite action solutions of the equation (7.17), these would be at the same time solutions of the renormalization group equations *and* solitons of the tachyon effective action (7.16). These could then be interpreted as lower dimensional branes. Being solutions of the renormalization group equations they are interpreted as background consistent with the string dynamics, being solitons they must describe branes. The finite action solutions of eq. (7.17) are easy to find

$$\tilde{T}(X) = -n + \frac{1}{2} \sum_{i=1}^n (X^i)^2. \tag{7.18}$$

These codimension n solitons can be interpreted as D(25 - n)-branes. $26 - n$ are in fact the number of coordinates on which the profile $\tilde{T}(X)$ does not depend. Substituting the solution (7.18) into the action (7.16) with $\log(c/4) = -1$ we get

$$S = T_{25} (e\sqrt{2\pi})^n V_{26-n}. \tag{7.19}$$

Comparing this with the expected result $T_{25-n} V_{26-n}$ we derive the following ratio between the brane tensions

$$R_n = \frac{T_{25-n}}{T_{25}} = \left(\frac{e}{\sqrt{2\pi}} 2\pi \right)^n. \tag{7.20}$$

With our notation, $\alpha' = 1$, the exact tension ratio should be $R_n = (2\pi)^n$. Thus R_n differs from the one given in (7.20) by a factor $e/\sqrt{2\pi} = 1.084$. It is remarkable that a small derivatives expansion of the WS action truncated just to the second order provides a result with the 93% of accuracy. In particular the result (7.20) is much closer to the exact tension ratio than the one found in [7] with analogous procedure. The solutions of the equations of motion of the WS action considered in [7] were not in fact solutions of the equation $\beta^T = 0$, so that they could not be interpreted as consistent string backgrounds (this was already noticed by the authors of [7] and for this reason the exact tension ratio was obtained with a different procedure). The equations of motion deriving from the WS action are in fact related to the β -function through (1.3) where the metric should in principle be non-degenerate. However, if the metric is computed in some approximation, it could be singular and present solutions that introduce physics beyond that contained in the β -functions. The action (7.16) with $\log(c/4) = -1$ gives an equation of the form (1.3) with the non-degenerate metric $e^{-\hat{T}}$. The solution of this equation can be at the same time a soliton and a conformal RG fixed point.

In conclusion the general formula (1.4) reproduces all the expected results on tachyon effective actions both in the far off-shell and in the near on-shell regions.

8. Cubic vs. Witten-Shatashvili tachyon effective actions

In this section we shall compare the result (7.2), which gives the WS action up to the third order in the powers of the renormalized tachyon field $T_R(k)$, with the tachyon effective action S_C computed with the cubic open string field theory of [1]. Like (7.2), S_C is known exactly up to the third power in the tachyon field. A similar comparison was already done in [7] where, however, the WS action constructed in terms of the linear tachyon β -function $\beta^T(k) = (1 - k^2)T_R(k)$ was used. With such a choice of coordinates in the space of string fields, the relation between the tachyon fields of the cubic and the WS string field theory becomes singular on-shell [7]. The cubic string field theory parametrization of worldsheet RG is regular on-shell and it very well reproduces the tachyon scattering amplitudes [26], thus indicating that to it should correspond a non-linear β -function. We shall show that, comparing the result (7.2) for the WS action with the corresponding cubic string field theory action, a field redefinition between the tachyon fields in the two formulations can be found which is non-singular on-shell. In particular, by requiring the regularity of the coordinate transformation that links the cubic tachyon effective action to the WS action (7.2) we find that the overall normalization constant K in the WS action (7.2) is precisely the tension of the D25-brane. This is in agreement with all the conjectures involving tachyon condensation and with the result $K = T_{25}$ derived from the tachyon potential.

For a tachyon field $\Phi(k)$, the cubic string field theory action can be written as [34, 35]

$$\begin{aligned}
 S_C = 2\pi^2 T_{25} \left[-\frac{1}{2} \int dk (2\pi)^D \Phi(k) \Phi(-k) (1 - k^2) + \frac{1}{3} \int dk_1 dk_2 dk_3 (2\pi)^D \times \right. \\
 \left. \times \delta(k_1 + k_2 + k_3) \Phi(k_1) \Phi(k_2) \Phi(k_3) \left(\frac{3\sqrt{3}}{4} \right)^{3 - k_1^2 - k_2^2 - k_3^2} \right]. \quad (8.1)
 \end{aligned}$$

The normalization factor $2\pi^2 T_{25}$ was derived in [27] and will be important for our analysis. Let us assume that the relation between the fields $\Phi(k)$ and $T_R(k)$ of the two theories is of the form

$$\Phi(k) = f_1(k)T_R(k) + \int dk_1 dk_2 f_2(k, k_1)T_R(k_1)T_R(k_2)\delta(k - k_1 - k_2) + \dots, \quad (8.2)$$

where $f_1(k) = f_1(-k)$ from the reality of the tachyon field. The cubic string field theory action (8.1) becomes

$$S_C = 2\pi^2 T_{25} \left\{ -\frac{1}{2} \int dk (2\pi)^D T_R(k)T_R(-k)(1 - k^2)(f_1(k))^2 - \int dk_1 dk_2 dk_3 (2\pi)^D \times \right. \\ \times \delta(k_1 + k_2 + k_3)T_R(k_1)T_R(k_2)T_R(k_3) \times \\ \times \left[(1 + k_1 k_2 + k_1 k_3)f_1(k_1)f_2(k_2 + k_3, k_3) - \right. \\ \left. \left. - \frac{1}{3}f_1(k_1)f_1(k_2)f_1(k_3) \left(\frac{3\sqrt{3}}{4} \right)^{3 - \sum_i k_i^2} \right] \right\}. \quad (8.3)$$

By comparing the second order term of eq. (8.1) with the corresponding term of eq. (7.2) we find

$$(f_1(k))^2 = \frac{K}{2\pi^2 T_{25}} \frac{\Gamma(2 - 2k^2)}{(1 - k^2)\Gamma^2(1 - k^2)}. \quad (8.4)$$

When the tachyon field is on the mass-shell, $f_1(k)$ is regular and takes the value

$$f_1 = \frac{1}{2\pi} \sqrt{\frac{K}{T_{25}}}. \quad (8.5)$$

From the comparison of the cubic terms in (8.3) and in (7.2) we get

$$[1 - (k_2 + k_3)^2]f_2(k_2 + k_3, k_3) = \frac{f_1(k_2)f_1(k_3)}{3} \left(\frac{3\sqrt{3}}{4} \right)^{3 - \sum_i k_i^2} - \frac{K}{3!2\pi^2 T_{25} f_1(k_2 + k_3)} \times \\ \times \left[2(1 - k_2 k_3 - k_2^2 - k_3^2)I(-k_2 - k_3, k_2, k_3) - \right. \\ \left. - \left(\frac{\Gamma(1 + 2k_2 k_3)\Gamma(2 - 2(k_2 + k_3)^2)}{\Gamma^2(1 + k_2 k_3)\Gamma^2(1 - (k_2 + k_3)^2)} + \text{cycl.} \right) \right]. \quad (8.6)$$

We can fix the value of the normalization constant K by requiring the regularity of the function $f_2(k_2 + k_3, k_3)$ when the three tachyons are on-shell. The on-shell condition is

$$2(k_1 k_2 + k_2 k_3 + k_1 k_3) + 3 = 0,$$

and the factor between square brackets in (8.6) is, on-shell, $(2\pi)^{-1}$. Consequently, requiring the regularity of the function f_2 when the three tachyons are on-shell, eq. (8.6) simply becomes

$$K = T_{25}. \quad (8.7)$$

When (8.7) is satisfied, the field redefinition that links the boundary and the cubic string field theory tachyon effective actions is regular on-shell. This result shows that the tachyon dynamics described by the WS string field theory reproduces the conjectured relations involving tachyon condensation. With analogous procedure, it is not difficult to show that the coordinate transformation between the WS tachyon effective action constructed in terms of the linear β -function (7.10), with $K = T_{25}$, and (8.1) is, as expected, singular on-shell.

9. Conclusions

In this paper we have derived some exact results for the non-linear tachyon β -function of the open bosonic string theory. We have shown its relevance in the construction of the Witten-Shatashvili bosonic string field theory. When a non-linear renormalization of the tachyon field is considered [23], the WS action in fact is simply given by (1.4). This formula has a wide range of validity. It can be applied to the case in which the tachyon profile is a slowly varying function of the embedding coordinates of the string to derive the exact tachyon potential and the first derivative terms of the effective action. Eq. (1.4) holds also when the tachyon coupling $T(k)$ is small and has support near the mass-shell. For such tachyon profiles we showed that perturbative solutions of the equation $\beta^T = 0$ provide the expected scattering amplitudes of on-shell tachyons.

The explicit form of the WS action constructed from the tachyon non-linear β -function is in precise agreement with all the conjectures involving tachyon condensation. In particular its normalization can be fixed either by studying the exact tachyon potential or by finding the field redefinition that maps the WS action into the effective tachyon action coming from the cubic string field theory. This field redefinition is non-singular on-shell only if the normalization constant coincides with the tension of the D25-brane.

The knowledge of the non-linear tachyon β -function is very important also for another reason. The solutions of the equation $\beta^T = 0$ give the conformal fixed points, the backgrounds that are consistent with the string dynamics. In the case of slowly varying tachyon profiles, we showed that the equations of motion for the WS action can be made identical to the RG fixed point equation $\beta^T = 0$. This can be done for a particular choice of the renormalization prescription ambiguity. We have found soliton solutions of this equation to which correspond a finite value of the WS action. Being solutions of the RG equations these solitons are lower dimensional D-branes for which the finite value of S provides a very accurate estimate of the D-brane tension.

When other excited string modes are present, say modes of the vector field A_μ , the form of the WS action should still be given by (1.4) where the renormalized tachyon field depends also on the other string couplings. In particular it would be interesting to apply our renormalization procedure to the other renormalizable case in which the boundary perturbation contains also a vector field [36]. Whether our approach would help in treating also non-renormalizable interactions it is yet not clear.

The decay of unstable systems of D-branes, pictured as a tachyon field rolling down a potential toward a stable minimum, can also be addressed in the context of the boundary string field theory. It involves deforming the world sheet conformal field theory of the

unstable D-brane by a conformal, time dependent tachyon profile. It is useful then to construct β -functions which are valid away from the RG fixed point to demonstrate that the renormalization flow exists, to draw the RG-trajectories and to understand where the endpoints of the RG flux are. Thus our approach should reveal important in studying the physics around a conformal fixed points and in particular about the time dependent solutions describing rolling tachyons.

Acknowledgments

E.C. and G.G. would like to thank R. Jackiw for an helpful discussion and the Bruno Rossi MIT-INFN fellowship for financial support. E.C. thanks I. Sigalov and W. Taylor for useful discussions. G.G. acknowledges the M.I.T. Center for Theoretical Physics for hospitality during the preparation of this work.

A. Computation of $I(k_1, k_2, k_3)$

In this appendix we shall compute the integral $I(k_1, k_2, k_3)$ appearing in eq. (4.15)

$$I(k_1, k_2, k_3) = \frac{2^{2k_1 k_2 + 2k_2 k_3 + 2k_1 k_3}}{(2\pi)^3} \int_0^{2\pi} d\tau_1 d\tau_2 d\tau_3 \left[\sin^2 \left(\frac{\tau_1 - \tau_2}{2} \right) \right]^{k_1 k_2} \times \\ \times \left[\sin^2 \left(\frac{\tau_2 - \tau_3}{2} \right) \right]^{k_2 k_3} \left[\sin^2 \left(\frac{\tau_1 - \tau_3}{2} \right) \right]^{k_1 k_3}. \quad (\text{A.1})$$

Introducing the variables

$$x = \frac{\tau_1 - \tau_2}{2}, \quad y = \frac{\tau_3 - \tau_1}{2}$$

the integral over τ_1, τ_2 and τ_3 can be written as

$$I = -\frac{4^{k_1 k_2 + k_2 k_3 + k_1 k_3 + 1}}{2\pi^3} \int_0^{2\pi} d\tau_1 \int_{\frac{\tau_1}{2}}^{\frac{\tau_1}{2} - \pi} dx \int_{-\frac{\tau_1}{2}}^{\pi - \frac{\tau_1}{2}} dy [\sin^2 x]^{k_1 k_2} [\sin^2 y]^{k_1 k_3} [\sin^2(x+y)]^{k_2 k_3}.$$

With a suitable shift of the integration variables we obtain

$$I = \frac{4^{k_1 k_2 + k_2 k_3 + k_1 k_3}}{\pi^2} \int_0^\pi dx \int_0^\pi dy [\sin x]^{2k_1 k_2} [\sin y]^{2k_1 k_3} [\sin^2(x+y)]^{k_2 k_3} \\ = \frac{4^{k_1 k_2 + k_2 k_3}}{\pi^2} \int_0^\pi dx \int_0^\pi dy [\sin x]^{2k_1 k_2} [\sin y]^{2k_1 k_3} [1 - e^{2i(x+y)}]^{k_1 k_3} [1 - e^{-2i(x+y)}]^{k_2 k_3} \\ = \frac{4^{k_1 k_2 + k_1 k_3}}{\pi^2} \int_0^\pi dx \int_0^\pi dy [\sin x]^{2k_1 k_2} [\sin y]^{2k_1 k_3} \times \\ \times \sum_{n,m=0}^{\infty} \frac{\Gamma(n - k_2 k_3) \Gamma(m - k_2 k_3)}{n! m! \Gamma^2(-k_2 k_3)} e^{2i(x+y)(n-m)}. \quad (\text{A.2})$$

Integrating over x and y we have

$$I = \sum_{n,m=0}^{\infty} \frac{\Gamma(n - a_3) \Gamma(m - a_3)}{n! m! \Gamma(1+a_1+n-m) \Gamma(1+a_1-n+m) \Gamma(1+a_2+n-m) \Gamma(1+a_2-n+m)} \times$$

$$\times \frac{\Gamma(1+2a_1)\Gamma(1+2a_2)}{\Gamma^2(-a_3)} \quad (\text{A.3})$$

where $a_1 = k_1 k_2$, $a_2 = k_2 k_3$ and $a_3 = k_1 k_3$. Shifting $m \rightarrow m - n$ in the sum over m we have

$$\begin{aligned} I = & \sum_{n,m=0}^{\infty} \frac{\Gamma(n-a_2)\Gamma(n+m+a_2)\Gamma(1+2a_1)\Gamma(1+2a_3)}{n!(n+m)!\Gamma^2(-a_2)\Gamma(1+a_1+m)\Gamma(1+a_1-m)\Gamma(1+a_3+m)\Gamma(1+a_3-m)} + \\ & + \sum_{n=0}^{\infty} \sum_{m=-n}^0 \frac{\Gamma(n-a_2)\Gamma(n+m+a_2)\Gamma(1+2a_1)\Gamma(1+2a_3)}{n!(n+m)!\Gamma^2(-a_2)\Gamma(1+a_1+m)\Gamma(1+a_1-m)\Gamma(1+a_3+m)\Gamma(1+a_3-m)} - \\ & - \frac{\Gamma(1+2a_1)\Gamma(1+2a_3)}{\Gamma^2(1+a_1)\Gamma^2(1+a_3)} {}_2F_1(-a_2, -a_2; 1; 1) \end{aligned} \quad (\text{A.4})$$

where ${}_2F_1(\alpha, \beta; \gamma; z)$ is the Hypergeometric function. Changing the sign of the integer m in the second term of the previous equation and noting that

$$\sum_{n=0}^{\infty} \sum_{m=0}^n = \sum_{n=m}^{\infty} \sum_{m=0}^{\infty}$$

we find

$$\begin{aligned} I = & 2 \sum_{m=0}^{\infty} \frac{\Gamma(m-a_2)\Gamma(1+2a_1)\Gamma(1+2a_3)}{\Gamma(-a_2)\Gamma(1+a_1+m)\Gamma(1+a_1-m)\Gamma(1+a_3+m)\Gamma(1+a_3-m)} \times \\ & \times {}_2F_1(m-a_2, -a_2; m+1; 1) - \frac{\Gamma(1+2a_1)\Gamma(1+2a_3)}{\Gamma^2(1+a_1)\Gamma^2(1+a_3)} {}_2F_1(-a_2, -a_2; 1; 1). \end{aligned} \quad (\text{A.5})$$

It is not difficult to show that the sum over m can be extended to negative values so that we find

$$\begin{aligned} I = & - \frac{\Gamma(1+2a_1)\Gamma(1+2a_2)\Gamma(1+2a_3)}{\Gamma(1-a_1)\Gamma(a_1)\Gamma(1-a_2)\Gamma(a_2)\Gamma(1-a_3)\Gamma(a_3)} \times \\ & \times \sum_{m=-\infty}^{\infty} \frac{\Gamma(m-a_1)\Gamma(m-a_2)\Gamma(m-a_3)}{\Gamma(1+m+a_1)\Gamma(1+m+a_2)\Gamma(1+m+a_3)}. \end{aligned} \quad (\text{A.6})$$

The series in the second line of the right-hand side of (A.6) is convergent for $1+a_1+a_2+a_3 > 0$. To sum over m we use a standard procedure. Consider the path in figure 1. Defining

$$S = \sum_{m=-\infty}^{\infty} \frac{\Gamma(m-a_1)\Gamma(m-a_2)\Gamma(m-a_3)}{\Gamma(1+m+a_1)\Gamma(1+m+a_2)\Gamma(1+m+a_3)} \equiv \sum_{m=-\infty}^{\infty} f(m) \quad (\text{A.7})$$

we can write

$$\oint_C \pi \cotg \pi z f(z) dz = \sum_{m=-N}^N f(m) + \tilde{S} \quad (\text{A.8})$$

where \tilde{S} is the sum of the residues of $\pi \cotg \pi z f(z)$ evaluated in the poles of $f(z)$. In the limit $N \rightarrow \infty$ the left-hand side of the previous equation vanishes reducing S to

$$S = - \frac{\Gamma(1+a_1+a_2+a_3)}{\Gamma(1+a_1)\Gamma(1+a_2)\Gamma(1+a_3)\Gamma(1+a_1+a_2)\Gamma(1+a_1+a_3)\Gamma(1+a_2+a_3)} \times$$




Figure 1: Contour C.

$$\times \left[\frac{\pi^3 \cos^2 \pi a_1}{\sin \pi a_1 \sin \pi(a_1 - a_2) \sin \pi(a_1 - a_3)} + \text{cycl.} \right]. \quad (\text{A.9})$$

So that I becomes

$$I = \frac{\Gamma(1 + a_1 + a_2 + a_3) \Gamma(1 + 2a_1) \Gamma(1 + 2a_2) \Gamma(1 + 2a_3)}{\Gamma(1 + a_1) \Gamma(1 + a_2) \Gamma(1 + a_3) \Gamma(1 + a_1 + a_2) \Gamma(1 + a_2 + a_3) \Gamma(1 + a_1 + a_3)}. \quad (\text{A.10})$$

References

- [1] E. Witten, *Noncommutative geometry and string field theory*, *Nucl. Phys.* **B 268** (1986) 253.
- [2] E. Witten, *On background independent open string field theory*, *Phys. Rev.* **D 46** (1992) 5467 [[hep-th/9208027](#)].
- [3] E. Witten, *Some computations in background independent off-shell string theory*, *Phys. Rev.* **D 47** (1993) 3405 [[hep-th/9210065](#)].
- [4] S.L. Shatashvili, *Comment on the background independent open string theory*, *Phys. Lett.* **B 311** (1993) 83 [[hep-th/9303143](#)].
- [5] S.L. Shatashvili, *On the problems with background independence in string theory*, [hep-th/9311177](#).
- [6] A.A. Gerasimov and S.L. Shatashvili, *On exact tachyon potential in open string field theory*, *J. High Energy Phys.* **10** (2000) 034 [[hep-th/0009103](#)].
- [7] D. Kutasov, M. Marino and G.W. Moore, *Some exact results on tachyon condensation in string field theory*, *J. High Energy Phys.* **10** (2000) 045 [[hep-th/0009148](#)].
- [8] A.A. Tseytlin, *Sigma model approach to string theory effective actions with tachyons*, *J. Math. Phys.* **42** (2001) 2854 [[hep-th/0011033](#)].

- [9] G. Grignani, M. Laidlaw, M. Orselli and G.W. Semenoff, *O(d) invariant tachyon condensates in the 1/D expansion*, *Phys. Lett.* **B 543** (2002) 127 [[hep-th/0206025](#)].
- [10] A. Sen, *Descent relations among bosonic D-branes*, *Int. J. Mod. Phys.* **A 14** (1999) 4061 [[hep-th/9902105](#)].
- [11] S. Elitzur, E. Rabinovici and G. Sarkissian, *On least action D-branes*, *Nucl. Phys.* **B 541** (1999) 246 [[hep-th/9807161](#)].
- [12] J.A. Harvey, S. Kachru, G.W. Moore and E. Silverstein, *Tension is dimension*, *J. High Energy Phys.* **03** (2000) 001 [[hep-th/9909072](#)].
- [13] A. Sen, *Rolling tachyon*, *J. High Energy Phys.* **04** (2002) 048 [[hep-th/0203211](#)].
- [14] A. Sen, *Tachyon matter*, *J. High Energy Phys.* **07** (2002) 065 [[hep-th/0203265](#)].
- [15] A. Sen, *Field theory of tachyon matter*, *Mod. Phys. Lett.* **A 17** (2002) 1797 [[hep-th/0204143](#)].
- [16] A. Sen, *Time evolution in open string theory*, *J. High Energy Phys.* **10** (2002) 003 [[hep-th/0207105](#)].
- [17] A. Sen, *Time and tachyon*, *Int. J. Mod. Phys.* **A 18** (2003) 4869 [[hep-th/0209122](#)].
- [18] M.R. Garousi, *Off-shell extension of s-matrix elements and tachyonic effective actions*, *J. High Energy Phys.* **04** (2003) 027 [[hep-th/0303239](#)].
- [19] M.R. Garousi, *S-matrix elements and off-shell tachyon action with non-abelian gauge symmetry*, *J. High Energy Phys.* **12** (2003) 036 [[hep-th/0307197](#)].
- [20] D. Kutasov and V. Niarchos, *Tachyon effective actions in open string theory*, *Nucl. Phys.* **B 666** (2003) 56 [[hep-th/0304045](#)].
- [21] A. Fotopoulos and A.A. Tseytlin, *On open superstring partition function in inhomogeneous rolling tachyon background*, *J. High Energy Phys.* **12** (2003) 025 [[hep-th/0310253](#)].
- [22] V. Niarchos, *Notes on tachyon effective actions and veneziano amplitudes*, [hep-th/0401066](#).
- [23] I.R. Klebanov and L. Susskind, *Renormalization group and string amplitudes*, *Phys. Lett.* **B 200** (1988) 446.
- [24] S.A. Frolov, *On off-shell structure of open string sigma model*, *J. High Energy Phys.* **08** (2001) 020 [[hep-th/0104042](#)].
- [25] A.B. Zamolodchikov, *Integrable field theory from conformal field theory*, *Adv. Stud. Pure Math.* **19** (1989) 641.
- [26] W. Taylor, *A perturbative analysis of tachyon condensation*, *J. High Energy Phys.* **03** (2003) 029 [[hep-th/0208149](#)].
- [27] A. Sen, *Universality of the tachyon potential*, *J. High Energy Phys.* **12** (1999) 027 [[hep-th/9911116](#)].
- [28] A. Sen, *Tachyon condensation on the brane antibrane system*, *J. High Energy Phys.* **08** (1998) 012 [[hep-th/9805170](#)].
- [29] C.G. Callan and L. Thorlacius, *Sigma models and string theory*, in proceedings *Particles, strings and supernovae*, vol. 2, pp 795–878, Providence 1988;
C.G. Callan Jr. and L. Thorlacius, *Open string theory as dissipative quantum mechanics*, *Nucl. Phys.* **B 329** (1990) 117; *Using reparametrization invariance to define vacuum infinities in string path integrals*, *Nucl. Phys.* **B 319** (1989) 133.

- [30] V.A. Kostelecky, M. Perry and R. Potting, *Off-shell structure of the string sigma model*, *Phys. Rev. Lett.* **84** (2000) 4541 [[hep-th/9912243](#)].
- [31] M.R. Garousi, *On-shell s-matrix and tachyonic effective actions*, *Nucl. Phys. B* **647** (2002) 117 [[hep-th/0209068](#)].
- [32] F. Larsen, A. Naqvi and S. Terashima, *Rolling tachyons and decaying branes*, *J. High Energy Phys.* **02** (2003) 039 [[hep-th/0212248](#)].
- [33] W. Taylor, *Perturbative diagrams in string field theory*, [hep-th/0207132](#).
- [34] K. Ohmori, *A review on tachyon condensation in open string field theories*, [hep-th/0102085](#).
- [35] D. Ghoshal and A. Sen, *Normalisation of the background independent open string field theory action*, *J. High Energy Phys.* **11** (2000) 021 [[hep-th/0009191](#)].
- [36] E. Coletti, V. Forini, G. Grignani, G. Nardelli and M. Orselli, in preparation.