



## City Research Online

### City, University of London Institutional Repository

---

**Citation:** Sanders, J., Bowden, T., Woolfe-Loftus, N., Sekhon, M. & Aitken, L. M. (2022). Predictors of health-related quality of life after cardiac surgery: a systematic review. *Health and Quality of Life Outcomes*, 20(1), 79. doi: 10.1186/s12955-022-01980-4

This is the published version of the paper.

This version of the publication may differ from the final published version.

---

**Permanent repository link:** <https://city-test.eprints-hosting.org/id/eprint/28209/>

**Link to published version:** <https://doi.org/10.1186/s12955-022-01980-4>

**Copyright:** City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

**Reuse:** Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

---

---



**PREDICTORS OF HEALTH-RELATED QUALITY OF LIFE AFTER CARDIAC SURGERY: A SYSTEMATIC  
REVIEW**

Julie Sanders<sup>1,2</sup> Tracey Bowden<sup>3</sup>, Nicholas Woolfe-Loftus<sup>1</sup>, Mandeep Sekhon<sup>3</sup> Leanne M Aitken<sup>3,4</sup>

<sup>1</sup> St Bartholomew's Hospital, West Smithfield, Barts Health NHS Trust. London, UK EC1A 7DN

<sup>2</sup> William Harvey Research Institute, Charterhouse Square, Queen Mary University of London,  
London, UK, EC1M 6BQ

<sup>3</sup> School of Health Sciences, City, University of London, London, UK. EC1V 0HB

<sup>4</sup> School of Nursing and Midwifery, Griffith University, Nathan, Queensland, Australia. QLD 4111.

**CORRESPONDING AUTHOR:** Professor Julie Sanders, Director of Clinical Research, St Bartholomew's  
Hospital, West Smithfield, Barts Health NHS Trust. West Smithfield. London. EC1A 7DN. Telephone:  
+44 (0)7960 311979. Email address: [j.sanders@qmul.ac.uk](mailto:j.sanders@qmul.ac.uk) Twitter: @julesanders2

# **PREDICTORS OF HEALTH-RELATED QUALITY OF LIFE AFTER CARDIAC SURGERY: A SYSTEMATIC**

## **REVIEW**

### **ABSTRACT**

#### **BACKGROUND**

Health-related quality of life (HRQoL) is important in determining surgical success, particularly from the patients' perspective.

#### **AIMS**

To identify predictors for HRQoL outcome after cardiac surgery in order to identify potentially modifiable risk factors where interventions to improve patient outcomes could be targeted.

#### **METHODS**

Electronic databases (including MEDLINE, CINAHL, Embase) were searched between January 2001 and December 2020 for studies determining risk predictors of HRQoL (using a recognised and validated tool) in adult patients undergoing cardiac surgery. Data extraction and quality assessments were undertaken and data was summarised using descriptive statistics and narrative synthesis, as appropriate.

#### **RESULTS**

Overall, 3924 papers were screened with 41 papers included in the review. Considerable methodological heterogeneity between studies was observed. Most were single-centre (75.6%) prospective observational studies (73.2%) conducted in patients undergoing coronary artery bypass graft (CABG) (n=51.2%) using a version of the SF-36 (n=63.4%). Overall, 103 independent predictors (62 pre-operative, five intra-operative and 36 post-operative) were identified, where 34 (33.0%) were reported in more than one study. Potential pre-operative modifiable risk factors include

alcohol use, BMI/weight, depression, pre-operative quality of life and smoking while in the post-operative period pain and strategies to reduce post-operative complications and intensive care and hospital length of stay are potential therapeutic targets.

## **CONCLUSION**

Despite a lack of consistency across studies, several potentially modifiable risk factors were identified that could be targeted in interventions to improve patient or treatment outcomes. This may contribute to delivering more person-centred care involving shared decision-making to improve patient HRQoL after cardiac surgery.

**KEYWORDS:** Health-related quality of life; predictors; cardiac surgery; quality of life; patient reported outcome

# **PREDICTORS OF HEALTH-RELATED QUALITY OF LIFE AFTER CARDIAC SURGERY: A SYSTEMATIC REVIEW**

## **INTRODUCTION**

High quality surgical care should include mortality, morbidity and patient-centred outcome measurement<sup>1</sup>. However, patient reported outcomes (PRO) are rarely recorded. Even in research contexts, PROs have only been reported in 29% of cardiac surgery trials<sup>2</sup>, despite the fact that those experiencing post-operative complications have worse quality of life<sup>3</sup>, which can last three years after surgery<sup>4</sup>.

Despite clinicians previously considering health-related quality of life (HRQoL) less important than clinical measures<sup>5</sup>, globally health ministers have stated the need to invest in measures that matter most to people<sup>6</sup>. HRQoL measurement allows a holistic, patient-centred perspective of recovery and it is becoming increasingly recognised that HRQoL is important in determining surgical success both from the patients<sup>7</sup> and health-care commissioners<sup>8</sup> perspective.

Factors that predict cardiac surgery mortality do not predict post-operative HRQoL outcome<sup>9</sup>. Thus, an understanding of the factors that do predict HRQoL would be useful to inform patients of the implications of surgery and interventions to improve potentially modifiable risk factors could be implemented. Certainly in the UK, HRQoL, and factors associated with it, was identified as the top ten research priority for adult cardiac surgery research<sup>10</sup>. We therefore undertook a literature review to ascertain the predictors of HRQoL after cardiac surgery, to identify potentially modifiable risk factors that could be targeted for intervention.

## **METHODS**

### Protocol and registration

This review was registered on PROSPERO, an international prospective register of systematic review (February 2019, reference CRD42019120080) and conducted in accordance with the Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) guidelines.

### Eligibility criteria

All studies that undertook multivariable analysis to identify independent predictors of HRQoL after cardiac surgery were eligible for inclusion. The detailed inclusion and exclusion criteria are detailed in Table 1.

**Table 1. Inclusion and exclusion criteria.**

Inclusion criteria	Exclusion criteria
Adult patients ( $\geq 18$ years of age)	Surgical ablation procedures in isolation
Primary research	Ventricular Assist Device (VAD) procedures
English language	Studies that did not include multivariable analysis of predictors of HRQoL only
Published 2001 – 2020	Congenital heart disease
Patients undergoing cardiac surgical procedures	Heart transplantation
	Transcatheter aortic valve implantation
	Descriptive exploration of interventions such as cardiac rehabilitation
	Studies that did not use a recognised quality of life instrument
	Comparison of quality of life in patients who underwent cardiac surgery with those who received percutaneous coronary intervention

## **Information sources, search strategy and study selection**

A search of MEDLINE, Cumulated Index of Nursing and Allied Health Literature (CINAHL), Embase, Cochrane Library and clinicaltrials.gov ([www.clinicaltrials.gov](http://www.clinicaltrials.gov)) was undertaken for relevant papers in English between January 2001 and December 2020. Search terms included cardiac surgery OR Cardiac Surgical Procedures AND quality of life OR outcome assessment and were adapted for each database (Supplementary material 1). Two authors screened the title and abstracts of all citations for suitability against the inclusion and exclusion criteria (Table 1). The reference lists of any identified systematic reviews were also screened for eligible papers.

## **Data collection and syntheses (data items and data collection process)**

Data were extracted by two authors into a standardised proforma with disagreements resolved through discussion until consensus was achieved. Data extraction included author, country, year, study design, type of surgery, sample size, HRQoL tool used including the time-points where HRQoL was measured, and the independent predictors of HRQoL.

## **Risk of bias and quality assessment**

All included papers were quality reviewed using an adapted Critical Appraisal Skills Programme (CASP) template for cohort studies ([https://casp-uk.net/wp-content/uploads/2018/01/CASP-Clinical-Prediction-Rule-Checklist\\_2018.pdf](https://casp-uk.net/wp-content/uploads/2018/01/CASP-Clinical-Prediction-Rule-Checklist_2018.pdf)). Initial papers were reviewed independently by two authors to ensure consistency and subsequent papers were reviewed by two of four authors with additional random checks undertaken at the end to be assured of continued assurance. A risk of bias graph was generated. Studies were not excluded on the basis of the quality assessment.



## **Analysis**

Following data extraction, results were summarised using descriptive statistics, tables and narrative synthesis, as appropriate. Interpretation of the analysis was discussed and agreed by all members of the authorship team. Meta-analysis was not possible due to the heterogeneity of studies.

## **RESULTS**

### **Study selection**

A total of 3924 papers were identified for possible inclusion (Figure 1) with 100 papers undergoing independent full-text assessment. This resulted in 41 papers being included for data synthesis.

### **Study characteristics**

Thirty-two studies were conducted in Europe (two of which were in the UK), four each in Australia the USA, and one in Canada (Table 2). The vast majority were single centre (n=31) with seven studies conducted in two centres and three studies conducted in multiple centres. Most were prospective observational studies (n=30) on patients undergoing coronary artery bypass graft (CABG) (n=21), CABG and/or valve surgery (n=10), valve only (n=1) or all types of cardiac surgery (n=9), with sample sizes in the HRQoL analysis ranging from 34 to 8676. A version of the SF-36 was used in the majority of studies (n=26).

In most studies HRQoL was measured pre-operatively (n=35) in addition to at least one post-operative assessment (Table 2), usually within six months of surgery (n=20) with twenty-four studies assessing outcome at one year or beyond (some studies assessed at more than one time-point).

HRQoL was not the primary outcome in all studies.

## **Risk of Bias**

Figures 2 and 3 demonstrate the variable risk of bias across studies and also in considering studies individually.

## **Independent predictors of HRQoL**

The independent predictors associated with HRQoL by operative and follow-up time-frame are detailed in Table 3 (and by study are included in are detailed in Supplementary Table 1). Overall, variables that were examined were predominantly focused on the clinical condition and experience of patients across the pre-, intra- and post-operative course. Of note, few demographic, social or psychological factors were incorporated into the analysis. Despite 26 studies (63.4%) using a version of the SF-36, how it was implemented and categorised to determine predictors varied across studies. For example, four studies explored predictors in relation to the overall score<sup>4,11,12,13</sup>, 13 explored the physical component (PCS) and mental component score (MCS) separately<sup>14,15,16,17,18,19,20,21,22,23,24,25,26</sup>, four studies explored one domain<sup>27,28,29,30</sup> two studies explored predictors in all SF-36 domains<sup>31,32</sup> while only Falcoz and colleagues explored both PCS and MCS and all domains<sup>33</sup>. Furthermore, Kube and colleagues used the SF-12 which measures physical and psychological quality of life<sup>34</sup>.

Due to the variation in analysis and reporting across the studies, the independent predictors identified were grouped by operative and follow-up time-frame (Table 3). In total, 103 independent predictors (62 pre-operative, 5 intra-operative and 36 post-operative) were identified. Of those 103 variables 34 (33.0%) were identified as significant in more than one study and almost all of those (n=33 (97.1%)) were also found to be non-significant in other studies (non-significant variable data detailed in Supplementary Table 2). Variables found to be predictive at all three time-points were age, angina, chronic obstructive pulmonary disease (COPD), diabetes, gender, hypertension and NYHA class and peripheral vascular disease.

## **Potentially modifiable risk factors**

Of the 62 pre-operative variables identified as independent risk factors for HRQoL outcome those that are potentially modifiable pre-surgery include alcohol use, body mass index (BMI)/weight, depression, pre-operative quality of life and smoking (Table 3). Similarly, in the post-operative period independent predictors with the potential to be modified to improve HRQoL outcome were pain, traumatic memories and restlessness in the intensive care unit (ICU). Furthermore, general strategies to reduce post-operative complications (including infection, myocardial infarction, arrhythmias and readmission) and shorten ICU and hospital length of stay are also identified as potential targets to improve post-surgical HRQoL (Table 3).

## **DISCUSSION**

The inclusion, measurement and use of HRQoL and PRO in cardiac surgery is lacking. Healthcare organisations need to work with patients to deliver more person-centred care, sharing decision-making, to meaningfully improve care outcomes<sup>35</sup>. The ‘holy grail’ of prognostic factor research is to improve patient outcomes by providing a personalised approach to healthcare and risk prediction and how such factors could be used to improve patient or treatment outcomes<sup>36</sup>. Thus, we sought to identify known risk predictors for HRQoL outcome after cardiac surgery, specifically to focus on modifiable risk factors where interventions to improve patient or treatment outcomes could be targeted. We identified 41 studies, which were predominantly European-based single-centre prospective observational studies conducted in CABG patients. Certainly, recognition of the non-modifiable risk factors found to be particularly impactful both on short and longer-term HRQoL (age, angina, COPD, diabetes, gender, hypertension and NYHA class and peripheral vascular disease) may assist in identifying high risk patients and the identification of interventions and associated resources that might then be directed to assisting these patients to recover. In terms of potential modifiable risk factors, pre-operative factors include alcohol use, smoking, BMI/weight depression, and pre-operative quality of life, while ongoing pain management, prevention of post-operative

complications and general strategies to reduce ICU and hospital length of stay could also be beneficial.

Individually focused lifestyle and therapeutic interventions have shown effectiveness in weight and BMI reduction<sup>37</sup>, decreasing alcohol consumption<sup>38</sup>, psychological preparation (including depression and anxiety)<sup>39</sup> and smoking cessation<sup>40</sup>. Given that BMI<sup>41</sup>, alcohol use<sup>42</sup>, depression and anxiety<sup>43</sup> and smoking<sup>44</sup> have also been identified to be associated with many in-hospital post-operative complications, strategies to encourage their reduction are likely to have beneficial impacts on improving overall morbidity and general recovery. As yet, interventions specifically targeting pre-operative HRQoL do not exist. While most tools combine physical, mental and social wellbeing traditionally greater emphasis clinically has placed on physical health. Nonetheless, the importance of psychological readiness and inclusion of social support and anxiety reduction in prehabilitation programmes is now recognised as part of cardiac surgery enhanced recovery<sup>45</sup>. Furthermore, we found that severe pain during the ICU stay was an independent predictor of HRQoL at six months<sup>46</sup>, while high pain scores at 15 months were predictive of HRQoL eight years after surgery in elderly patients<sup>47</sup>. Since up to 10% of cardiac surgery patients develop severe chronic post-surgical pain<sup>48</sup>, with predictors of chronic pain including early severe pain<sup>49</sup> personalised effective pain management is vital. Current recommendations suggest the use of multimodal opioid-sparing pain management alongside the use of a pain assessment tool to ensure the lowest opioid dose<sup>45</sup>.

Certainly, future work requires more methodologically robust studies, including large multi-site studies with appropriate control of confounding factors. However, generally a greater emphasis on HRQoL as an outcome measure after cardiac surgery, both clinically and in research, is needed. Although HRQoL has been previously undervalued by clinicians<sup>5</sup>, the landscape is changing with the importance of HRQoL now recognised in cardiac surgery clinical guidelines<sup>50</sup>, the enhanced recovery recommendations<sup>45</sup>, the cardiac surgery core outcome dataset<sup>51</sup> and that PROs are included in the

Swedish national registers<sup>52</sup> and emerging in the USA STS National Database<sup>53</sup>. Similarly, HRQoL is reported as a top research priority in cardiac surgery, both in the UK<sup>10</sup> and in the USA<sup>54</sup>. Therefore, our review is timely, in that it collates the available evidence on predictors of HRQoL, highlights potential modifiable factors on which interventions could be based to improve patient outcome and emphasises where greater research quality in prognosis factor research should reside in this area.

## **Strengths and limitations**

Despite the methodological robustness of this review, there are three main limitations. Firstly, the methodological heterogeneity of the included studies restricts the ability to make strong conclusions or undertake a meta-analysis. Our review reflects that despite the considerable growth in prognosis research, the quality is often sub-standard<sup>55</sup>. Secondly, although only English language publications were included, studies from around the World have been included, providing a relatively wide base of evidence. Finally, included studies were limited to those published from 2001. A balance was struck between including all evidence and ensuring the results of this review were clinically appropriate outcome risk factors for the current time. A period of 20-years was deemed sufficient to address the balance needed.

In conclusion, despite a lack of consistency across studies, several potentially modifiable risk factors on which interventions to improve patient HRQoL outcomes could be targeted were identified. While this review has robustly collated the current best prognosis factor evidence relating to risk predictors of HRQoL after cardiac surgery, there is still a need for large multi-site studies, with appropriate control of confounding factors, to examine the role of these factors in affecting HRQoL outcome. Now that considerably more emphasis is placed on the importance of HRQoL and PROs after cardiac surgery, the hope is that this will contribute to delivering more person-centred care involving shared decision-making to improve patient short- and longer-term recovery.

## 271 **IMPLICATIONS FOR PRACTICE**

- 272 • Cardiac surgery and enhanced recovery guidelines highlight the importance of HRQoL
- 273 • Pre-operative lifestyle and therapeutic interventions relating to weight, alcohol use,
- 274 psychological preparation and smoking cessation may improve HRQoL
- 275 • Reducing chronic post-operative pain, in-hospital complications and length of hospital stay
- 276 could also improve HRQoL.
- 277 • More person-centred care, including HRQoL and shared decision-making, is needed to
- 278 improve patient short- and longer-term recovery

279

## 280 **ABBREVIATIONS**

281	BMI	body mass index
282	CABG	coronary artery bypass graft
283	CASP	Critical Appraisal Skills Programme
284	COPD	chronic obstructive pulmonary disease
285	HRQoL	health-related quality of life
286	ICU	intensive care unit
287	MCS	mental component score
288	NYHA	New York Heart Association
289	PCS	physical component score
290	PRISMA	Preferred Reporting Items for Systematic reviews and Meta-Analyses
291	PRO	patient reported outcomes
292	PROSPERO	International Prospective Register of Systematic Reviews
293	SF-36	Short-Form 36

294

295

296

297 **DECLARATIONS**

298 **Ethics approval and consent to participate**

299 Not applicable

300

301 **Consent for publication**

302 Not applicable

303

304 **Availability of data and materials**

305 Not applicable

306

307 **Competing interests**

308 The authors declare that they have no competing interests

309

310 **Funding**

311 This work was supported by Barts Charity (Grant Reference Number: MRD0181) to Leanne Aitken for  
312 research associate support.

313

314 **Authors' contributions**

315 **JS:**Conceptualisation, data curation, formal analysis, investigation, methodology, project  
316 administration, supervision, validation, visualisation, writing-original draft; **TB:**data curation, formal  
317 analysis, investigation, validation, writing-review and editing; **NW:**data curation, investigation,  
318 writing-review and editing; **MS:**data curation, formal analysis, investigation, writing-review and  
319 editing; **LA:**Conceptualisation, data curation, formal analysis, funding acquisition, investigation,  
320 methodology, validation, writing-review and editing.

321

322 **Acknowledgements**

323 Not applicable

324

325

326

327

328 **REFERENCES**

329 1. Birkmeyer JD, Dimick JB, Birkmeyer NJO. Measuring the quality of surgical care: structure,  
330 process, or outcomes? *J Am Coll Surg* 2004; 198: 626–632.

331 2. Goldfarb M, Drudi L, Almohammadi M, et al. Outcome reporting in cardiac surgery trials:  
332 Systematic review and critical appraisal. *J Am Heart Assoc* 2015; 4: 1–9.

333 3. Martin C, Holmes S, Martin L, et al. Abstract P331: The Impact of In-Hospital Postoperative  
334 Complications on Health Related Quality of Life in Cardiac Surgery Patients. In: *Circulation:*  
335 *Cardiovascular Quality and Outcomes*,

336 4. Myles PS, Viira D, Hunt JO. Quality of life at three years after cardiac surgery: relationship  
337 with preoperative status and quality of recovery. *Anaesth Intensive Care* 2006; 34: 176–83.

338 5. Calvert M, Kyte D, Price G, et al. Maximising the impact of patient reported outcome  
339 assessment for patients and society. *BMJ* 2019; 364: 1–8.

340 6. Ministers OH. *Ministerial Statement: The Next Generation of Health Reforms Ministerial*  
341 *Statement*. Epub ahead of print 2017. DOI: 10.1016/j.palaeo.2006.08.010.

342 7. Colak Z, Segotic I, Uzun S, et al. Health related quality of life following cardiac surgery -  
343 correlation with EuroSCORE. *Eur J Cardio-thoracic Surg* 2008; 33: 72–76.

344 8. Devlin NJ, Buxton M, Vallance-Owen A, et al. *Getting the most out of PROMs: health*  
345 *outcomes and NHS decision-making*, [www.kingsfund.org.uk/publications](http://www.kingsfund.org.uk/publications) (2010, accessed 13  
346 April 2018).

347 9. Vainiola T, Pettilä V, Roine RP, et al. Comparison of two utility instruments, the EQ-5D and  
348 the 15D, in the critical care setting. *Intensive Care Med* 2010; 36: 2090–3.

349 10. Lai FY, Abbasciano RG, Tabberer B, et al. Identifying research priorities in cardiac surgery: A  
350 report from the James Lind Alliance Priority Setting Partnership in adult heart surgery. *BMJ*  
351 *Open* 2020; 10: 1–9.



- 352 11. Myles PS, Hunt JO, Fletcher H, et al. Relation between Quality of Recovery in Hospital and  
353 Quality of Life at 3 Months after Cardiac Surgery. *Anesthesiology* 2001; 95: 862–867.
- 354 12. Norkienė I, Urbanaviciute I, Kezyte G, et al. Impact of pre-operative health-related quality of  
355 life on outcomes after heart surgery. *ANZ J Surg* 2018; 88: 332–336.
- 356 13. Kurfist V, Mokráček A, Krupauerová M, et al. Health-related quality of life after cardiac  
357 surgery--the effects of age, preoperative conditions and postoperative complications. *J*  
358 *Cardiothorac Surg* 2014; 9: 46.
- 359 14. Baldassarre FG, Arthur HM, DiCenso A, et al. Effect of coronary artery bypass graft surgery on  
360 older women's health-related quality of life. *Hear Lung J Acute Crit Care* 2002; 31: 421–431.
- 361 15. Schelling G, Richter M, Roozendaal B, et al. Exposure to high stress in the intensive care unit  
362 may have negative effects on health-related quality-of-life outcomes after cardiac surgery.  
363 *Crit Care Med* 2003; 31: 1971–1980.
- 364 16. Baberg HT, Dirlich M, Laczkovics A, et al. Determinants of health-related quality of life after  
365 aortic valve replacement in six-month survivors of intervention. *J Heart Valve Dis* 2004; 13:  
366 914–920.
- 367 17. Rumsfeld JS, Ho PM, Magid DJ, et al. Predictors of health-related quality of life after coronary  
368 artery bypass surgery. *Ann Thorac Surg* 2004; 77: 1508–1513.
- 369 18. Al-Ruzzeh S, Athanasiou T, Mangoush O, et al. Predictors of poor mid-term health related  
370 quality of life after primary isolated coronary artery bypass grafting surgery. *Heart* 2005; 91:  
371 1557–62.
- 372 19. Le Grande MR, Elliott PC, Murphy BM, et al. Health related quality of life trajectories and  
373 predictors following coronary artery bypass surgery. *Health Qual Life Outcomes* 2006; 4: 49.
- 374 20. Grady KL, Lee R, Subačius H, et al. Improvements in Health-Related Quality of Life Before and  
375 After Isolated Cardiac Operations. *Ann Thorac Surg* 2011; 91: 777–783.
- 376 21. Patron E, Messerotti Benvenuti S, Palomba D. Preoperative biomedical risk and depressive  
377 symptoms are differently associated with reduced health-related quality of life in patients

378 1 year after cardiac surgery. *Gen Hosp Psychiatry* 2016; 40: 47–54.

379 22. Grand N, Bouchet JB, Zufferey P, et al. Quality of Life After Cardiac Operations Based on the  
380 Minimal Clinically Important Difference Concept. *Ann Thorac Surg* 2018; 106: 548–554.

381 23. Coelho PNMP, Miranda LMRPC, Barros PMP, et al. Quality of life after elective cardiac surgery  
382 in elderly patients. *Interact Cardiovasc Thorac Surg* 2019; 28: 199–205.

383 24. Blokzijl F, Houterman S, Van Straten BHM, et al. Quality of life after coronary bypass: A  
384 multicentre study of routinely collected health data in the Netherlands. *Eur J Cardio-thoracic  
385 Surg* 2019; 56: 526–533.

386 25. Joskowiak D, Meusel D, Kamla C, et al. Impact of Preoperative Functional Status on Quality of  
387 Life after Cardiac Surgery. *Thorac Cardiovasc Surg*. Epub ahead of print 2019. DOI: 10.1055/s-  
388 0039-1696953.

389 26. Perrotti A, Ecamot F, Monaco F, et al. Quality of life 10 years after cardiac surgery in adults: a  
390 long-term follow-up study. *Health Qual Life Outcomes* 2020; 18: 1–10.

391 27. Jarvinen O, Julkunen J, Saarinen T, et al. Perioperative myocardial infarction has negative  
392 impact on health-related quality of life following coronary artery bypass graft surgery. *Eur J  
393 Cardio-Thoracic Surg* 2004; 26: 621–627.

394 28. Rijnhart-De Jong H, Haenen J, Bol Raap G, et al. Determinants of non-recovery in physical  
395 health-related quality of life one year after cardiac surgery: A prospective single Centre  
396 observational study. *J Cardiothorac Surg* 2020; 15: 1–10.

397 29. Juergens MC, Seekatz B, Moosdorf RG, et al. Illness beliefs before cardiac surgery predict  
398 disability, quality of life, and depression 3 months later. *J Psychosom Res* 2010; 68: 553–560.

399 30. Deaton C, Thourani V. Patients with Type 2 Diabetes Undergoing Coronary Artery Bypass  
400 Graft Surgery: Predictors of Outcomes. *Eur J Cardiovasc Nurs* 2009; 8: 48–56.

401 31. El Baz N, Middel B, Van Dijk JP, et al. EuroSCORE predicts poor health-related physical  
402 functioning six month postcoronary artery bypass graft surgery. *J Cardiovasc Surg (Torino)*  
403 2008; 49: 663–72.

- 404 32. Humphreys JM, Denson LA, Baker RA, et al. The importance of depression and alcohol use in  
405 coronary artery bypass graft surgery patients: Risk factors for delirium and poorer quality of  
406 life. *J Geriatr Cardiol* 2016; 13: 51–57.
- 407 33. Falcoz PE, Chocron S, Stoica L, et al. Open Heart Surgery: One-Year Self-Assessment of Quality  
408 of Life and Functional Outcome. *Ann Thorac Surg* 2003; 76: 1598–1604.
- 409 34. Kube T, Meyer J, Grieshaber P, et al. Patients' pre- and postoperative expectations as  
410 predictors of clinical outcomes six months after cardiac surgery. *Psychol Heal Med* 2020; 25:  
411 781–792.
- 412 35. Alderwick H, Dixon J. The NHS long term plan. *BMJ* 2019; 364: 1–136.
- 413 36. Riley RD, van der Windt DA, Croft P, et al. (eds). *Prognosis research in healthcare: Concepts,*  
414 *methods and impact*. Oxford: Oxford University Press, 2019.
- 415 37. Johnson M, Desiree Backman M, Neal Kohatsu R, et al. *Interventions for Reducing Body Mass*  
416 *Index and Other Weight-related Indicators: A Review of Systematic Reviews,*  
417 [https://www.ucdmc.ucdavis.edu/iphi/Programs/OP/resources/Interventions for Reducing](https://www.ucdmc.ucdavis.edu/iphi/Programs/OP/resources/Interventions%20for%20Reducing%20BMI%20and%20Other%20Weight-related%20Indicators%20Review%20of%20Systematic%20Reviews_12-15-16.pdf)  
418 [BMI and Other Weight-related Indicators Review of Systematic Reviews\\_12-15-16.pdf](https://www.ucdmc.ucdavis.edu/iphi/Programs/OP/resources/Interventions%20for%20Reducing%20BMI%20and%20Other%20Weight-related%20Indicators%20Review%20of%20Systematic%20Reviews_12-15-16.pdf) (2016,  
419 accessed 19 April 2018).
- 420 38. Budworth L, Prestwich A, Lawton R, et al. Preoperative Interventions for Alcohol and Other  
421 Recreational Substance Use: A Systematic Review and Meta-Analysis. *Front Psychol* 2019; 10:  
422 34.
- 423 39. Salzmann S, Salzmann-Djufri M, Wilhelm M, et al. Psychological Preparation for Cardiac  
424 Surgery. *Curr Cardiol Rep* 2020; 22: 172.
- 425 40. Cahill, Stevens, Perera, et al. Pharmacological interventions for smoking cessation: an  
426 overview and network meta-analysis. *Cochrane Database Syst Rev* 2013; N.PA-N.PA.
- 427 41. Gao M, Sun J, Young N, et al. Impact of Body Mass Index on Outcomes in Cardiac Surgery. *J*  
428 *Cardiothorac Vasc Anesth* 2016; 30: 1308–16.
- 429 42. Eliassen M, Grønkjær M, Skov-Ettrup LS, et al. Preoperative alcohol consumption and

430 postoperative complications: a systematic review and meta-analysis. *Ann Surg* 2013; 258:  
 431 930–42.

432 43. Pignay-Demaria V, Lespérance F, Demaria RG, et al. Depression and anxiety and outcomes of  
 433 coronary artery bypass surgery. *Ann Thorac Surg* 2003; 75: 314–321.

434 44. Sepehrpour AH, Lo TT, McCormack DJ, et al. Is there benefit in smoking cessation prior to  
 435 cardiac surgery? *Interact Cardiovasc Thorac Surg* 2012; 15: 726–732.

436 45. Engelman DT, Ben Ali W, Williams JB, et al. Guidelines for Perioperative Care in Cardiac  
 437 Surgery: Enhanced Recovery after Surgery Society Recommendations. *JAMA Surg* 2019; 12:  
 438 1–12.

439 46. Vainiola T, Roine RP, Suojaranta-Ylinen R, et al. Can factors related to mortality be used to  
 440 predict the follow-up health-related quality of life (HRQoL) in cardiac surgery patients?  
 441 *Intensive Crit Care Nurs* 2013; 29: 337–43.

442 47. Jokinen JJ, Hippelainen MJ, Hanninen T, et al. Prospective assessment of quality of life of  
 443 octogenarians after cardiac surgery: factors predicting long-term outcome. *Interact*  
 444 *Cardiovasc Thorac Surg* 2008; 7: 813–818.

445 48. Gjeilo KH, Stenseth R, Klepstad P. Risk Factors and Early Pharmacological Interventions to  
 446 Prevent Chronic Postsurgical Pain Following Cardiac Surgery. *Am J Cardiovasc Drugs* 2014; 14:  
 447 335–342.

448 49. Bjørnnes AK, Parry M, Lie I, et al. Pain experiences of men and women after cardiac surgery. *J*  
 449 *Clin Nurs* 2016; 25: 3058–3068.

450 50. Sousa-Uva M, Neumann FJ, Ahlsson A, et al. 2018 ESC/EACTS Guidelines on myocardial  
 451 revascularization. *Eur J Cardio-Thoracic Surg* 2019; 55: 4–90.

452 51. Benstoem C, Moza A, Meybohm P, et al. A core outcome set for adult cardiac surgery trials: A  
 453 consensus study. *PLoS One* 2017; 12: 1–12.

454 52. Nilsson E, Orwelius L, Kristenson M. Patient-reported outcomes in the Swedish National  
 455 Quality Registers. *J Intern Med* 2016; 279: 141–153.

- 456 53. D'Agostino RS, Jacobs JP, Badhwar V, et al. The Society of Thoracic Surgeons Adult Cardiac  
457 Surgery Database: 2019 Update on Outcomes and Quality. *Ann Thorac Surg* 2019; 107: 24–  
458 32.
- 459 54. Khazanie P, Krumholz HM, Kiefe CI, et al. Priorities for Cardiovascular Outcomes Research: A  
460 Report of the National Heart, Lung, and Blood Institute's Centers for Cardiovascular  
461 Outcomes Research Working Group. *Circ Cardiovasc Qual Outcomes* 2017; 10: 1–8.
- 462 55. Riley R, Moons KG, Hayden J, et al. Prognostic Factor Research. In: Riley R, van der Windt D,  
463 Croft P, et al. (eds) *Prognosis Research in Healthcare*. Oxford: Oxford University Press, 2019,  
464 pp. 107–138.
- 465 56. Herlitz J, Brandrup-Wognsen G, Caidahl K, et al. Improvement and factors associated with  
466 improvement in quality of life during 10 years after coronary artery bypass drafting. *Coron*  
467 *Artery Dis* 2003; 14: 509–517.
- 468 57. Schelling G, Richter M, Roozendaal B, et al. Exposure to high stress in the intensive care unit  
469 may have negative effects on health-related quality-of-life outcomes after cardiac surgery.  
470 *Crit Care Med*. 2003; 31: 1971–1980.
- 471 58. Herlitz J, Brandrup-Wognsen G, Caidahl K, et al. Determinants for an impaired quality of life  
472 10 years after coronary artery bypass surgery. *Int J Cardiol* 2005; 98: 447–452.
- 473 59. Peric V, Borzanovic M, Jovanovic A, et al. The relationship between EuroSCORE preoperative  
474 risk prediction and quality of life changes after coronary artery by-pass surgery. *Interact*  
475 *Cardiovasc Thorac Surg* 2005; 4: 622–626.
- 476 60. Noyez L, Markou A, van Breugel F. Quality of life one year after myocardial revascularization.  
477 Is preoperative quality of life important? *Interact Cardiovasc Thorac Surg* 2006; 5: 115–120.
- 478 61. Panagopoulou E, Montgomery A, Benos A. Quality of life after coronary artery bypass  
479 grafting: evaluating the influence of preoperative physical and psychosocial functioning. *J*  
480 *Psychosom Res* 2006; 60: 639–644.
- 481 62. Dunning J, Waller JRL, Smith B, et al. Coronary Artery Bypass Grafting is Associated With

- 482           Excellent Long-Term Survival and Quality of Life: A Prospective Cohort Study. *Ann Thorac Surg*  
483           2008; 85: 1988–1993.
- 484   63.   Peric V, Borzanovic M, Stolic R, et al. Predictors of Worsening of Patients' Quality of Life Six  
485           Months After Coronary Artery Bypass Surgery. *J Card Surg* 2008; 23: 648–654.
- 486   64.   Herlitz J, Brandrup-Wognsen G, Evander MH, et al. Quality of life 15 years after coronary  
487           artery bypass grafting. *Coron Artery Dis* 2009; 20: 363–369.
- 488   65.   Maisano F, Viganò G, Calabrese C, et al. Quality of life of elderly patients following valve  
489           surgery for chronic organic mitral regurgitation. *Eur J Cardio-thoracic Surg* 2009; 36: 261–266.
- 490   66.   Rantanen A, Kaunonen M, Tarkka M, et al. Patients' and significant others' health-related  
491           quality of life one month after coronary artery bypass grafting predicts later health-related  
492           quality of life. *Heart Lung J Acute Crit Care* 2009; 38: 318–329.
- 493   67.   Peric V, Borzanovic M, Stolic R, et al. Quality of life in patients related to gender differences  
494           before and after coronary artery bypass surgery☆. *Interact Cardiovasc Thorac Surg* 2010; 10:  
495           232–238.
- 496   68.   Bjørnnes AK, Parry M, Falk R, et al. Impact of marital status and comorbid disorders on  
497           health-related quality of life after cardiac surgery. *Qual Life Res* 2017; 26: 2421–2434.
- 498   69.   Bishawi M, Hattler B, Almassi GH, et al. Preoperative factors associated with worsening in  
499           health-related quality of life following coronary artery bypass grafting in the Randomized  
500           On/Off Bypass (ROOBY) trial. *Am Heart J* 2018; 198: 33–38.
- 501   70.   Schaal NK, Assmann A, Rosendahl J, et al. Health-related quality of life after heart surgery –  
502           Identification of high-risk patients: A cohort study. *Int J Surg* 2020; 76: 171–177.

505 **Table 2: Study characteristics (n=41).**

Study  (Author, year,  country)	Study design  (including (number of  sites))	Patients (population (type of surgery) and sample size)			HRQoL tool used  and time-point of  predictive model  (months unless  otherwise stated)
		Type of surgery	Sample size		
			Participation rate of eligible persons	Completed follow-up: total cohort	
Myles 2001 <sup>11</sup>  Australia	Pre-op post-op (1)	CABG, valve,  combined, other	120/125 (96%)	108/120 (90%)	SF-36(3)
Baldassarre  2002 <sup>14</sup>  Canada	Prospective cohort (1)	Isolated CABG  (primary)	34/64 (53%)	30/34 (88%)	SF-36(3)

Falcoz 2003 <sup>33</sup> France	Prospective cohort (1)	CABG, valve, combined, other (elective)	293/452 (65%)	264/293 (90%)	SF-36(12)
Herlitz 2003 <sup>56</sup> Sweden	Prospective cohort (2)	Isolated CABG (primary)	1225/2000 (61%)	976/2000 (49%)	NHP(10years)
Schelling 2003 <sup>57</sup> Germany	Prospective cohort (1)	CABG, valve, combined	223/387 (58%)	148/223 (66%)	SF-36(6)
Baberg 2004 <sup>16</sup> Germany	Prospective and retrospective cohort (1)	AVR +/- MVR	201/414 (47%)	201/414 (49%)	SF-36(3years) <sup>a</sup>
Jarvinen 2004 <sup>27</sup> Finland	Prospective cohort (1)	Isolated CABG	501/1128 (44%)	458/501 (91%)	SF-36(12)



Rumsfeld 2004 <sup>17</sup> America	Prospective cohort (14)	Isolated CABG (primary)	2480/3956 (63%)	1973/2480 (80%)	SF-36(6)
Al-Ruzzeh 2005 <sup>18</sup> UK	Cross-sectional (1)	Isolated CABG (primary)	437/463 (94%)	NA	SF-36(12) <sup>a</sup>
Herlitz 2005 <sup>58</sup> Sweden	Prospective cohort (1)	Isolated CABG (primary)	1225/2000 (61%)	637/1225 (52%)	NHP(10years) <sup>b</sup>
Peric 2005 <sup>59</sup> Serbia and Montenegro	Prospective cohort (1)	Elective CABG	243 (no mention of consent/refusal rate)	226/243 (93%)	NHP(6)
Le Grande 2006 <sup>19</sup>	Pre-op post-op (1)	Elective CABG	182/407 (45%)	117/182 (64%)	SF-36(2, 6)

Australia					
Myles 2006 <sup>4</sup>  Australia	Pre-op post-op (1)	CABG, valve,  combined, other	108/120 (90%)	93/108 (86%)	SF-36(3, 3years)
Noyez 2006 <sup>60</sup>  Netherlands	Retrospective cohort  (1)	Isolated CABG	428/428 (100%)	428/428 (100%)	EQ-5D(12)
Panagopoulou2  006 <sup>61</sup>  Greece	Prospective cohort (1)	Elective CABG	157/256 (61%)	1mo:117/157 (75%)  6mo:104/157 (66%)	MNHD-Q(1,6)
Dunning 2008 <sup>62</sup>  England	Prospective cohort (1)	Isolated CABG	911/1180 (77%)	621/911 (68%)	EQ-5D(10) <sup>a</sup>
El Baz 2008 <sup>31</sup>  Netherlands	Prospective  observational (2)	CABG	198/256 (73%)	168/198 (85%)	SF-36(6)

Jokinen 2008 <sup>47</sup>  Finland	Prospective observational cohort  (1)	CABG, valve, combined, other	91/98 (93%)	46/91 (51%)	NHP(15, 8.2years) <sup>b</sup>
Peric 2008 <sup>63</sup>  Serbia	Pre-op post-op (1)	Isolated CABG	Not reported	192/208 (92%)	NHP(6)
Deaton 2009 <sup>30</sup>  USA	Prospective cohort (2)	Isolated CABG	317/442 (72%)	270/317 (85%)	SF-36(3) <sup>a</sup>
Herlitz 2009 <sup>64</sup>  Sweden	Prospective cohort (2)	Isolated CABG  (primary)	Not reported.  2000 screened	639/2000 (32%)	NHP(15 years)
Maisano 2009 <sup>65</sup>  Italy (implied by authorship, not stated)	Retrospective cohort with prospective assessment of HRQoL  (1) (implied)	Mitral valve surgery  (± AVR,± TV surgery,± CABG)	225/225 (100%)	220/225 (98%)	MLHF(3 years) <sup>a</sup>
Rantanen  2009 <sup>66</sup>	Prospective cohort (1)	Elective CABG	1mo:274/367 (75%)  6mo:244/271 (90%)	1mo:270/274 (99%)  6mo:240/244 (98%)	15D(1,6,12)

Finland			12mo:236/266 (89%)	12mo:235/236 (100%)	
Juergens 2010 <sup>29</sup> Germany	Prospective cohort (1)	Elective CABG, valve, combined	56/85(65%)	42/65(75%)	SF-12(3)
Peric 2010 <sup>67</sup> Serbia	Prospective pre-op post-op (1)	Elective CABG	243/243 (100%)	226/243 (93%)	NHP(6)
Grady 2011 <sup>20</sup> America	Prospective cohort (1)	CABG, valve, maze, combined	840/2524 (33%)	0mo:173/840 (21%) 6mo:177/840 (21%) 12mo:174/840 (21%) 24mo:129/840 (15%) 36mo:69/840 (8%) Total:816/840 (97%)	SF-36(3, 6, 12, 2years, 3years)
Vainiola 2013 <sup>46</sup> Finland	Prospective cohort (1)	CABG, valve, combined, aortic, other	785/980 (80%)	571/785 (73%)	15-D(6)

Kurfirst 2014 <sup>13</sup> Czech republic	Prospective cohort (1)	CABG, valve, combined (elective)	310 eligible.	260/310 (84%)	SF-36(12)
Humphreys 2016 <sup>32</sup> Australia	Prospective cohort (1)	Elective CABG	180 agreed to participate. No further details	173/180 (96%)	SF-36(6)
Patron 2016 <sup>21</sup> Italy	Pre-op post-op (1)	CABG, valve, combined (primary, elective)	92/92 (100%)	75/92 (82%)	SF-12(12)
Bjornnes 2017 <sup>68</sup> Norway	Secondary analysis of RCT (2)	CABG, valve, combined	416/525 (79%)	349/416 (84%)	15D(2 weeks, 3,6,12)

Norkiene 2018 <sup>12</sup> Lithuania	Prospective cohort (1)	CABG, valve, combined	210 / 210 (100%) No further details	105/210(50%)	SF-36(12)
Bishawi 2018 <sup>69</sup> America	Secondary analysis of RCT (18)	Isolated CABG (urgent or elective)	2203/3670 (60%)	1770 / 2203 (80%)	SAQVR-36(12)
Grand 2018 <sup>22</sup> France	Prospective cohort (1)	CABG, valve, combined (elective)	495/548 (90%)	326/495 (66%)	SF-36(6)
Coelho 2019 <sup>23</sup> Portugal	Prospective cohort (1) (implied)	CABG, valve, combined (elective)	Not stated	384/430 (89%)	SF-36(12)
Blokzijl 2019 <sup>24</sup> Netherlands	Retrospective cohort multicentre (3)	Elective CABG	2606/8643 (30%)	2606/8643 (30%)	SF-36 or SF12(10- 14)
Joskowiak 2019 <sup>25</sup> Germany	Prospective cohort (1)	CABG, valve, combined, aortic, other, redo (elective)	164 consented but does not state number	164/164 (100%)	SF-36(12)

			who were eligible and screened		
Perrotti 2019 <sup>26</sup> France	Prospective cohort (1)	Isolated CABG (elective)	272/272 (100%)	118/272 (43%)	SF-36(10years)
Kube 2020 <sup>34</sup> Germany	Prospective cohort (2)	CABG, valve, combined (elective)	70/110 (64%)	53/70 (76%)	SF-12(6)
Rijnhart-de Jong 2020 <sup>28</sup>	Prospective cohort (1)	Non-salvage cardiac surgery	1544/1773 (87%)	874/1544 (57%)	SF-36(12)
Schaal 2020 <sup>70</sup> Germany	Prospective cohort (1)	CABG, valve, combined, aortic,	8676/14043 (62%)	8676/14043 (62%)	NHP(6)

506 <sup>a</sup> no preoperative HRQoL assessment performed; <sup>b</sup> preoperative HRQoL assessment carried out in some, but not all patients; **Abbreviations:** AVR:aortic  
507 valve replacement; CABG:Coronary Artery Bypass Surgery; EQ-5D:EuroQoL- 5 Dimension; HRQoL:Health-related Quality of Life; MLHF:Minnesota Living with  
508 Heart Failure questionnaire; MNHD-Q:MacNew Heart Disease Quality of Life Questionnaire; MVR:mitral valve replacement; NHP:Nottingham Health Profile;

509 PF:physical function; SAQ:Seattle Angina Questionnaire; SF-12:12 item short form health survey; SF-36:36 item short form health survey; TV:tricuspid valve;

510 VR-36:Veteran's Rand(version of SF36)

511 **Table 3: Independent variables by operative and follow-up time period**

	Independent predictors of HRQoL between 1 and 6 months follow-up	Independent predictors of HRQoL between 1year and 3.5years follow-up	Independent predictors of HRQoL at 8.2 years and beyond
Pre-operative variables	Age <sup>14,17,32, 34,70</sup> Alcohol use <sup>32</sup> ASA score <sup>15,22</sup> Angina <sup>22, 31,70</sup> BMI <sup>30</sup> Cerebrovascular disease <sup>32, 33,69</sup> Charlton Comorbidity Index <sup>30</sup> Chronic Heart Failure <sup>32</sup> Chronic neurological disease <sup>17</sup>	Atrial fibrillation <sup>65</sup> Age <sup>20, 23,25,65,66, 68</sup> Angina class <sup>33</sup> Back/neck problems <sup>68</sup> BMI <sup>20</sup> Chronic Heart Failure <sup>18,20</sup> COPD <sup>20, 33,69</sup> Cerebrovascular accident <sup>69</sup> Depression <sup>21,68,69</sup>	Age <sup>56,58,64</sup> Angina <sup>26</sup> COPD <sup>58</sup> COPD/asthma <sup>62</sup> CCSC <sup>62</sup> Diabetes <sup>26,47,58,62,64</sup> Duration of cardiac symptoms pre-op >120 days <sup>47</sup> Dyspnea <sup>26</sup>



	<p>COPD<sup>17</sup></p> <p>Delirium<sup>32</sup></p> <p>Depression<sup>32</sup></p> <p>Diabetes mellitus<sup>46,63,67,</sup></p> <p>Ejection fraction<sup>17,63</sup></p> <p>Ethnicity (aboriginal)<sup>32</sup></p> <p>Everyday functioning questionnaire<sup>19</sup></p> <p>EuroSCORE<sup>31,59</sup></p> <p>FEV1<sup>17</sup></p> <p>Gender<sup>63,67</sup></p> <p>Gender:Male<sup>46,70</sup></p> <p>Gender:Female<sup>30</sup></p> <p>Geriatric depression scale<sup>30</sup></p> <p>Health behaviours<sup>32</sup></p> <p>History of renal insufficiency<sup>31</sup></p>	<p>Diabetes Mellitus<sup>16, 28,65,69</sup></p> <p>Education<sup>21,33,68</sup></p> <p>Ejection Fraction<sup>24,33</sup></p> <p>Ethnicity (non-white)<sup>20</sup></p> <p>EuroSCORE<sup>21,65</sup></p> <p>Gastrointestinal tract problems<sup>18</sup></p> <p>Gender<sup>23</sup></p> <p>Gender:Female<sup>20,28</sup></p> <p>Gender/marital status interaction<sup>68</sup></p> <p>Hypertension<sup>20</sup></p> <p>Infection<sup>28</sup></p> <p>MI<sup>20</sup></p> <p>Mobility<sup>60</sup></p> <p>Neurological disease<sup>25</sup></p> <p>NYHA class<sup>16,20, 33,65</sup></p>	<p>Gender:Female<sup>62,64</sup></p> <p>Height<sup>58</sup></p> <p>Hypertension<sup>56,58</sup></p> <p>NYHA class<sup>56,64</sup></p> <p>Obesity<sup>58,64</sup></p> <p>Pre-op Nottingham Health Profile<sup>56</sup></p> <p>Pre-op inferior QoL<sup>64</sup></p> <p>Protective use of statin<sup>47</sup></p> <p>Peripheral Vascular Disease<sup>62</sup></p> <p>Redo surgery<sup>62</sup></p> <p>Smoking<sup>62</sup></p>
--	---	--	---

	Hypertension <sup>17,32</sup> Illness perception questionnaire <sup>29</sup> Living alone <sup>30,70</sup> NYHA class <sup>17,19,31,70</sup> Peripheral vascular disease <sup>17</sup> Previous cardiac surgery <sup>19</sup> Pre-op 15-D <sup>46</sup> Pre-op expectations (IPQ-E) <sup>34</sup> Pre-op MacNew score <sup>61</sup> Pre-op MCS <sup>17</sup> Pre-op PCS <sup>17</sup> Pre-op SF-12 physical QoL <sup>34</sup> Pre-op SF-12 psychological Qo <sup>34</sup> Profile of mood state vigor-activity <sup>19</sup> Profile of mood state fatigue-inertia <sup>19</sup>	Other diseases <sup>66</sup> Pain intensity <sup>68</sup> Parsonnet score <sup>33</sup> PCI<1yr <sup>28</sup> Peripheral or cerebral vascular Disease <sup>33</sup> Peripheral Vascular Disease <sup>18,20</sup> Pre-op MCS <sup>13,12,23 24,25</sup> Pre-op PCS <sup>13,12,21, 23 24,25,28</sup> Pre-op SAQ <sup>69</sup> Pre-op SF-36 <sup>4</sup> Pre-op VAS <sup>60</sup> Pre-op VR36 <sup>69</sup> Presence coronary artery disease <sup>20</sup> Pulmonary disease <sup>24</sup> QoR-40 <sup>4</sup>	
--	--	--	--

	Psychiatric disease <sup>17</sup> Psychological distress <sup>61</sup> QoR-40 <sup>11</sup> Serum creatinine <sup>17</sup> Smoking <sup>17, 31,67</sup> Stress symptom score <sup>15</sup> Work:Not in workforce <sup>19,70</sup> Work manual occupation <sup>19</sup>	Redo surgery <sup>20</sup> Renal disease <sup>24</sup> Segment wall motion (abnormal) <sup>33</sup> Serum creatinine <sup>65</sup> Type D personality <sup>18</sup> Type valvular heart disease <sup>16</sup>	
Intra-operative variables	Cardiopulmonary bypass duration <sup>22</sup> Higher mean pulmonary pressure <sup>19</sup> CABG procedure <sup>70</sup>	On cardiopulmonary bypass <sup>33</sup> Prosthetic valve type <sup>16</sup> CABG procedure <sup>25</sup>	Inotropic drugs at time of surgery <sup>64</sup>
Post-operative variables	Aid from network members <sup>66</sup> Complications <sup>63,67</sup> Dobutamine <sup>22</sup> Length of hospital stay <sup>31</sup>	Infective complications <sup>18</sup> Ejection fraction at follow-up <sup>65</sup> Hospital LOS <sup>4,23</sup> ICU LOS <sup>23</sup>	High pain score at 15 months <sup>47</sup> ICU time <sup>64</sup> ICU 2 days <sup>62</sup> ICU >3 days <sup>47</sup>

	<p>MI<sup>66</sup></p> <p>Prolonged LOS<sup>30</sup></p> <p>New cardiac arrhythmia<sup>19</sup></p> <p>No of categories of traumatic memory<sup>15</sup></p> <p>Other diseases<sup>66</sup></p> <p>Pain:severe and unbearable<sup>46</sup></p> <p>Physical exertion causing symptoms<sup>66</sup></p> <p>Post-op expectations (IPQ-E)<sup>34</sup></p> <p>Post-op SF-12 physical QoL<sup>34</sup></p> <p>Post-op SF-12 psychological QoL<sup>34</sup></p> <p>QoL at 1month (15D)<sup>61,66</sup></p> <p>Readmission to hospital within 6weeks<sup>31</sup></p> <p>Reexploration<sup>31</sup></p> <p>Renal replacement for acute renal failure<sup>22</sup></p> <p>Restlessness during ICU treatment<sup>46</sup></p>	<p>Mitral regurgitation at follow-up echo<sup>65</sup></p> <p>Perioperative MI<sup>27</sup></p> <p>QoL at 1month (15D)<sup>66</sup></p> <p>Quality of life at 3months<sup>4</sup></p> <p>Poor QOR-40 at 1month<sup>4</sup></p> <p>Physical exertion causing symptoms<sup>66</sup></p> <p>Post-operative rhythm<sup>16</sup></p> <p>Sternal complications<sup>16</sup></p> <p>Symptoms on mild exertion or at rest<sup>66</sup></p> <p>Systolic pulmonary artery pressure at follow-up<sup>65</sup></p>	<p>Length ventilator time<sup>58</sup></p> <p>Low energy score at 15months<sup>47</sup></p>
--	--	--	---

	Sternal resuturing <sup>31</sup> Symptoms on mild exertion or at rest <sup>66</sup> Ventilation >48hrs <sup>22</sup>		
--	--	--	--

512 BMI:Body Mass Index; CABG:Coronary Artery Bypass Graft; CCSC: Canadian Cardiovascular Society; COPD:Chronic obstructive pulmonary disease;

513 FEV1:Forced Expiratory Volume; ICU:Intensive Care Unit; LOS:Length of Stay; MCS:mental component score; MI:Myocardial Infarction; NYHA:New York

514 Heart Association classification; PCI:Percutaneous Coronary Intervention; PCS:Physical Component Score; QoL:Quality of Life; SAQ:Seattle Angina

515 Questionnaire; VAS:Visual Analogue Scale; VR36:Veteran's Ran

