



City Research Online

City, University of London Institutional Repository

Citation: Walsh, T. S., Parker, R. A., Aitken, L., McKenzie, C. A., Glen, R. & Weir, C. J. (2025). Relative and bedside nurse assessment of comfort and communication during propofol, dexmedetomidine, or clonidine-based sedation: pre-planned analysis within the A2B RCT. Health Technology Assessment,

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/35668/>

Link to published version:

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Relative and bedside nurse assessment of comfort and communication during propofol, dexmedetomidine, or clonidine-based sedation: pre-planned analysis within the A2B RCT

Timothy S Walsh^{1*}, Richard A. Parker², Leanne M Aitken³, Cathrine A McKenzie⁴, Robert Glen⁵, Christopher J Weir², the A2B Trial Investigators Group†

¹ Usher Institute, The University of Edinburgh, Edinburgh, UK

² Edinburgh Clinical Trials Unit, Usher Institute, The University of Edinburgh, Edinburgh, UK

³ City St George's University of London, London, UK

⁴ National Institute of Health and Social Care Research (NIHR), Biomedical Research Centre (BRC), Southampton, Perioperative and Critical Care theme, University of Southampton, Southampton, UK

⁵ Lay member, Edinburgh

†Members of the A2B Trial Investigators Group are listed in Appendix 1

***Corresponding Author:**

Professor Timothy Walsh

Department of Anaesthesia, Critical Care & Pain Medicine

Usher Institute

University of Edinburgh

Room S8208, 2nd Floor, The Royal Infirmary of Edinburgh

NHS Lothian, 51 Little France Crescent

Edinburgh BioQuarter, Edinburgh, EH16 4SA

E-mail: timothy.walsh@ed.ac.uk

Key words

Critical Care; intensive care units; hypnotics and sedatives; respiration, artificial; adrenergic alpha-2 receptor agonists; dexmedetomidine; clonidine

Word count: 3441

Abstract

Background

Optimising comfort and ability to communicate for mechanically ventilated (MV) intensive care (ICU) patients is a priority for clinicians, ICU patients and their relatives. Current usual care is propofol-based sedation plus an opioid analgesic. The alpha2-agonists dexmedetomidine and clonidine are potential alternative sedatives.

Objective(s)

To explore whether nurses and relatives perceive patients sedated with dexmedetomidine and/or clonidine appear more awake, comfortable, and cooperative than patients receiving only propofol-based sedation.

Design and methods

Sub-study within an open-label three-arm trial.

Setting and participants

41 ICUs in the United Kingdom. 1437 adults receiving propofol $\pm$ opioid for sedation-analgesia within 48 hours of starting MV, expected to require ≥ 48 total hours of MV.

Interventions

Light sedation was targeted in all patients unless clinicians requested deeper sedation. In intervention groups algorithms promoted alpha2-agonist up-titration and propofol down-titration followed by sedation primarily with allocated alpha2-agonist. Usual care was propofol-based sedation. Intervention continued until patients were successfully extubated (primary outcome), or other pre-defined end-points.

Outcomes

For each 12-hours care period nurses responded to two 'yes/no' questions: *Is the patient able to communicate pain? Is the patient able to cooperate with care?* When the patients' personal legal representative visited, they were asked for 'yes/no' responses to three questions: *does the patient appear awake? Does the patient appear comfortable? Does the visitor feel they can communicate with the patient?*

Intervention versus propofol group responses were compared fitting a generalised linear mixed model, with results expressed as Odds Ratios (OR; 95% confidence intervals); ORs >1 indicated greater probability of a 'yes' response.

Results

Nurse responses were available for >90% of trial patients (mean (SD) 12 (12) care periods per patient). Comparing dexmedetomidine versus propofol groups, the OR for a 'yes' response to '*communicate pain*' was 1.38 (1.08 to 1.75), and for clonidine versus propofol was 1.13 (0.89 to 1.43). For '*cooperate with care*' comparing dexmedetomidine versus propofol groups, OR was 1.14 (0.98 to 1.32), and for clonidine versus propofol 0.96 (0.83 to 1.12). Relative responses were available for 32-34% of trial patients across groups (mean (SD) 3 (3) days per patient). For the '*appear awake*' question, the dexmedetomidine versus propofol group OR was 1.48 (1.04 to 2.10), and clonidine versus propofol 1.35 (0.95 to 1.91). For '*appear comfortable*' the dexmedetomidine versus propofol group OR was 0.64 (0.38 to 1.09), and for clonidine versus propofol 0.78 (0.45 to 1.34). For the '*feel they can communicate*' comparison the dexmedetomidine versus propofol group OR was 1.00 (0.68 to 1.47), and for clonidine versus propofol 1.05 (0.71 to 1.54).

Limitations

Interventions were unblinded, with risk of bias; missing data may not have been at random.

Conclusions

Nurses perceived patients receiving dexmedetomidine-based sedation could better communicate pain than with propofol-based sedation, and relatives perceived patients appeared more awake. No differences for the other questions were found, or for the clonidine versus propofol comparisons, although some uncertainty remains due to the wide confidence intervals.

Future work

Additional mixed methods research of sedation quality with different agents from staff and relative perspectives.

Study registration:

NCT03653832

Word count: 498

Funding details - This project was funded by the National Institute for Health and Care Research (NIHR) Health Technology Appraisal programme and will be published in XXX Journal; Vol. XX, No. XX. See the NIHR Journals Library website for further project information

Plain Language summary

Patients receiving intensive care (ICU) usually require sedation to keep them comfortable and pain free. The most widely used sedation is currently a drug called propofol. Two alternatives are the drugs dexmedetomidine and clonidine. Both are called 'alpha2-agonists' based on how they work.

The A2B trial was a large study comparing these three drugs for sedating patients during their ICU stay. The trial found no differences in the time spent on the breathing machine or in the ICU. There were no differences in rates of delirium or the quality of sedation measured using scoring systems, except patients who received the alpha2-agonists appeared to experience more agitation compared with propofol.

As part of the trial, we asked bedside nurses to answer 'yes' or 'no' to questions asking their opinion at the end of their nursing shift on whether patients appeared able to communicate pain and could cooperate with care. We also asked the patients' closest relatives' their 'yes/no' response to questions about whether they thought their relative appeared awake, comfortable, and whether they could communicate with the patient when visiting. We compared responses to each question for both alpha2-agonist groups with the propofol group.

We found that, overall, nurses thought patients sedated with dexmedetomidine were better able to communicate pain than with propofol, but found no difference between clonidine and propofol. There were no differences in the ability to cooperate with care. Relatives thought patients sedated with dexmedetomidine appeared more awake than with propofol. The difference for clonidine was smaller.

Despite finding no differences in clinical outcomes between the sedation approaches used in the A2B trial, both nurses and relatives may perceive differences in some aspects of sedation quality depending on the sedation medicine used.

Word count: 298

Background

Achieving comfort, analgesia and enabling communication with intensive care unit (ICU) patients is a goal of optimum sedation.¹ The views of staff and relatives regarding the overall comfort of patients and their ability to cooperate with care have not previously been studied in trials of ICU sedation. Protocols to guide pain, agitation and delirium management typically use validated rating scales completed by bedside staff, such as the Richmond Agitation Sedation Scale (RASS),² Confusion Agitation Method for ICU (CAM-ICU),³ and Behavioural Pain Scale (BPS).⁴ These categorise patient status in order to prompt changes in treatment rather than reflecting the views of staff and relatives. Many patients recall pain, anxiety and discomfort following ICU discharge, with a high prevalence of frightening memories and long-term psychological morbidity.^{5,6} The need for more holistic approaches to ensuring patient comfort has been highlighted by consensus groups and guidelines.^{1,7}

The A2B trial was a randomised trial comparing ICU sedation based on propofol, dexmedetomidine, or clonidine for patients expected to require more than 48 hours of mechanical ventilation (MV).⁸ The main results of the trial have been reported elsewhere.⁹ Briefly, neither dexmedetomidine- or clonidine-based sedation decreased the primary outcome of time to successful extubation. Among secondary outcomes, neither alpha2-agonist improved rates of delirium, unnecessary deep sedation, or pain behaviours but both were associated with higher rates of patient agitation. Within the trial, we undertook a pre-planned exploration of the views of bedside nurses and relatives visiting patients regarding their perceptions of patient comfort and communication while receiving sedation with the allocated trial interventions. We hypothesised that nurses and relatives would perceive that the patients sedated with dexmedetomidine and/or clonidine were more awake, comfortable, and cooperative than patients receiving only propofol-based sedation. We report here the results of this analysis.

Methods

A2B trial overview:

The A2B trial protocol has been published.⁸ Briefly, eligible patients were receiving MV in the ICU, aged ≥ 18 years, and were within 48 hours of starting MV and sedated with propofol. At randomisation they were expected to require a further 24 hours of MV and a total of ≥ 48 hours. Exclusion criteria included: acute brain injury; neuromuscular paralysis; bradycardia < 50 beats/minute for ≥ 60 minutes; and patients not expected to survive a further 24 hours. Randomisation used a remote web-based system, allocating in a 1:1:1 ratio to the three groups using permuted blocks stratified by centre. Intervention-group patients commenced intravenous infusion

of open-label study drug using a weight-based dose regimen within two hours post-randomisation. Bedside nurses used group-specific algorithms to up-titrate alpha2-agonist and down-titrate propofol to transition patients to receive the allocated alpha2-agonist, with the aim of alpha2-agonist based sedation, supplemented with propofol if required. Usual care was propofol-based sedation without specific dose-guidance. For all groups bedside algorithms indicated a sedation target RASS score of -2 to +1 (range: -5 (unresponsive) to +4 (combative)), unless responsible clinicians requested deeper sedation for therapeutic reasons. The choice and dosing of opioid for analgesia was determined by the clinical team according to usual care and clinical judgement. For the majority of patients this was either fentanyl or alfentanil according to local prescribing policies. Other sedatives were discouraged and recorded as 'rescue medications' if required. Interventions continued until the patient was successfully extubated, died during MV, was transferred before extubation to a non-participating ICU, or until the end of 28 days of MV.

Ethical approval:

The A2B trial received ethical approval from the Scotland A REC (Reference 18/SS/0085) on 21/08/2018. This exploratory analysis was part of the main protocol, and the relevant outcomes were included in the pre-specified Statistical Analysis Plan.¹⁰ Ethical approval allowed any clinical nurse providing care during the trial to participate without additional written consent. For relatives, ethical approval was provided for the Personal Legal Representative (PerLegR) who had provided written consent for participation in the trial, usually the next-of-kin, to be approached to provide responses when they visited the patient. Assessments of relatives' views were therefore always the same individual for each patient. As the A2B trial included a deferred consent model when relatives were unavailable two hours after confirming eligibility, the PerLegR might have provided consent after starting the intervention.

Assessment and data collection:

All patients were assessed while receiving their allocated sedation intervention from randomisation until either primary outcome (successful extubation), death prior to primary outcome, transfer to another ICU before extubation, or 28 days post-randomisation without achieving the primary outcome. Caring nurse data collection was based on 12-hours nursing shifts. The nurse was asked to complete a 'nursing shift' form that included: whether a clinical request for deep sedation was made; regular RASS scores (suggested four hourly); a CAM-ICU score (once per shift); and assessments of two behavioural pain ratings based on ventilator compliance and upper limb movement. These outcomes contributed to outcomes reported for the main trial publication. At the end of each nursing care period ('day shift' and 'night shift') nurses were also asked to respond to two binary 'yes/no' questions about their view of the patient's comfort and cooperation during that

shift, namely: *Is the patient able to communicate pain?* And, *Is the patient able to cooperate with care?*

For relative assessments, when the PerLegR visited and agreed to provide an opinion, they were asked to provide the caring nurse with a binary 'yes/no' response to three questions, namely: *does the patient appear awake to the visitor?* *Does the patient appear comfortable to the visitor?* And, *does the visitor feel they can communicate with the patient?* The bedside nurse recorded responses on the daily shift forms. Data recorded on daily shift forms were entered onto the trial database by local research staff.

Patient and Public Involvement (PPI)

The decision to include the views of visiting relatives was strongly supported by a PPI group who helped develop the trial funding proposal and subsequently contributed to protocol design. The choice and wording of questions put to relatives was informed by a group of ICU survivors and their relatives. A lay co-applicant, Mr Bob Glen, was part of the Trial Management Group throughout the trial and reviewed information materials. An independent PPI representative was a member of the Trial Steering Committee. Mr Glen reviewed and approved the final manuscript, and is a co-author.

Equality, Diversity and Inclusion (EDI)

The trial inclusion/exclusion criteria had no limitations based on gender, LGBTQ preference, ethnicity, social status or geographical location. Children were excluded, but are usually managed in different ICUs from adults, experience a different spectrum of illness, and are typically studied in paediatric-centric ICU trials.

Blinding

Clinical staff were not blinded to group allocation, as they were managing sedation and titrating the allocated sedatives. Relatives were not formally blinded as part of the trial. We did not record whether relatives had asked or been told which group the patient was allocated to.

Trial registration

The A2B trial registration is NCT03653832.

Analysis

Analysis population

The analysis population comprised all randomised patients in whom nursing and/or relative responses were recorded at least once during the intervention period (and who were not excluded for another reason from the pre-defined overall trial analysis population).⁹ We described the

numbers of patients in whom data were available and the mean (Standard Deviation, SD) number of days on which responses were available from nurses and from relatives.

Statistical analyses

Baseline characteristics for patients included in the trial have been published previously.⁹ For the present analysis, we summarised relevant baseline data for those patients included in the analysis of nursing responses and visiting relative responses. We calculated the mean (SD) numbers of days with available data for each question, and mean (SD) proportion of care periods with 'yes' responses to the questions.

We analysed outcomes by fitting a generalised linear mixed model with a logit link function, using all available data. Site was included as a random effect in the model and treatment group as a fixed effect. For these outcomes, which were measured in multiple care periods, a random effect for participant (nested within site) was also included. Results were expressed as the odds ratio (OR) for each of dexmedetomidine and clonidine versus propofol-based sedation, with corresponding 95% confidence interval (CI). A higher OR indicated a greater probability of a 'yes' response to the question analysed.

We recognised that these assessments were most relevant when the patient was either not comatose due to their illness, or concurrent sedation level during the nursing shift was at RASS -3 or higher, indicating patients who were not comatose. In sensitivity analysis, we therefore summarised data when the highest RASS score during the concurrent nursing shift was -3 or higher, and repeated the same generalised linear mixed model analysis restricted to responses for nursing shifts in which a RASS score of -3 or above was recorded.

All analyses were undertaken using SAS version 9.4.

Results

From December 2018 to October 2023, we randomized 1438 patients in 41 ICUs in the United Kingdom. One patient was randomised twice in error. The pre-defined overall trial analysis population comprised 1404 patients allocated to receive propofol (N = 471), dexmedetomidine (N = 457), or clonidine (N = 476) as primary sedative. The numbers of patients and proportions of the overall analysis population in whom data were available for this analysis are summarised in table 1.

A summary of baseline characteristics for the populations analysed is shown in table 2. Patients included in both the bedside nurse assessments and visiting relative assessments were similar to the published overall trial population. There were no clinically relevant differences between the three

groups, or between the sub-group for whom visiting relative assessments were available compared with bedside nursing assessments.

Bedside nurse response to questions

A summary of the bedside nurse responses to the questions is shown in table 3.

'Is the patient able to communicate pain?' The mean proportion of days with 'yes' response was: dexmedetomidine 39%; clonidine 35%; and propofol 33%. Comparing the dexmedetomidine with propofol group the OR for a 'yes' response was: 1.38 (95% confidence interval 1.08 to 1.75); for the clonidine with propofol group comparison the OR for a 'yes' response was 1.13 (95% CI 0.89 to 1.43). Restricting the analysis to days with RASS -3 or above the mean proportion of days was: dexmedetomidine 44%; clonidine 40%; and propofol 37%. Comparing the dexmedetomidine with propofol group the OR for a 'yes' response was: 1.37 (95% confidence interval (CI) 1.07 to 1.74); for the clonidine with propofol group comparison the OR for a 'yes' response was 1.09 (0.86 to 1.38).

'Is the patient able to cooperate with care?' The mean proportion of days was: dexmedetomidine 42%; clonidine 38%; and propofol 38%. Comparing the dexmedetomidine with propofol group the OR for a 'yes' response was: 1.14 (95% CI 0.98 to 1.32); for the clonidine with propofol group comparison the OR for a 'yes' response was 0.96 (95% CI 0.83 to 1.12). Restricting the analysis to days with RASS -3 or above the mean proportion of days was: dexmedetomidine 46%; clonidine 42%; and propofol 42%. Comparing the dexmedetomidine with propofol group the OR for a 'yes' response was: 1.11 (0.95 to 1.30); for the clonidine with propofol group comparison the OR for a 'yes' response was 0.94 (0.80 to 1.09).

Visiting relative responses to questions

A summary of the visiting relative responses to the questions is shown in table 4.

'Does the patient appear awake to the visitor?' The mean proportion of days was: dexmedetomidine 53%; clonidine 50%; and propofol 44%. Comparing the dexmedetomidine with propofol group the OR for a 'yes' response was: 1.48 (95% CI 1.04 to 2.10).; for the clonidine with propofol group comparison the OR for a 'yes' response was 1.35 (0.95 to 1.91). Restricting the analysis to days with RASS -3 or above the mean proportion of days was: dexmedetomidine 56%; clonidine 54%; and propofol 49%. Comparing the dexmedetomidine with propofol group the OR for a 'yes' response was: 1.31 (95% CI 0.93 to 1.85); for the clonidine with propofol group comparison the OR for a 'yes' response was 1.25 (0.89 to 1.77).

'Does the patient appear comfortable to the visitor?' The mean proportion of days was: dexmedetomidine 88%; clonidine 90%; and propofol 93%. Comparing the dexmedetomidine with

propofol group the OR for a 'yes' response was: 0.64 (0.38 to 1.09); for the clonidine with propofol group comparison the OR for a 'yes' response was 0.78 (0.45 to 1.34). Restricting the analysis to days with RASS -3 or above the mean proportion of days was: dexmedetomidine 89%; clonidine 91%; and propofol 93%. Comparing the dexmedetomidine with propofol group the OR for a 'yes' response was: 0.58 (0.33 to 1.02); for the clonidine with propofol group comparison the OR for a 'yes' response was 0.70 (0.39 to 1.25).

'Does the visitor feel they can communicate with the patient?' The mean proportion of days was: dexmedetomidine 54%; clonidine 52%; and propofol 52%. Comparing the dexmedetomidine with propofol group the OR for a 'yes' response was: 1.00 (0.68 to 1.47); for the clonidine with propofol group comparison the OR for a 'yes' response was 1.05 (0.71 to 1.54). Restricting the analysis to days with RASS -3 or above the mean proportion of days was: dexmedetomidine 56%; clonidine 54%; and propofol 57%. Comparing the dexmedetomidine with propofol group the OR for a 'yes' response was: 0.87 (0.59 to 1.30); for the clonidine with propofol group comparison the OR for a 'yes' response was 0.96 (0.64 to 1.42).

Table 1: Proportion of patients in the overall trial analysis population contributing data to each of the bedside nurse and visiting relative questions.

Question	Proportion of analysis population (Number)		
	Propofol	Dexmedetomidine	Clonidine
Bedside Nurse			
<i>Is the patient able to communicate pain?</i>	97% (457)	94% (430)	95% (452)
<i>Is the patient able to cooperate with care</i>	93% (438)	90% (413),	92% (438).
Visiting Relative			
<i>Does the patient appear awake to the visitor?</i>	34% (162)	32% (148)	34% (164)
<i>Does the patient appear comfortable to the visitor?</i>	33% (156)	32% (147)	34% (160)
<i>Does the visitor feel they can communicate with the patient?'</i>	33% (157)	33% (150)	33% (158)

Table 2: Comparison of patients' baseline characteristics for the patients in whom: bedside nurse assessments of patients' comfort and communication were recorded, and visiting relative assessments of comfort and ability to communicate were recorded.

	Bedside nurse assessments			
Variable	Dexmedetomidine group (N = 456)	Clonidine group (N = 476)	Propofol (N = 471)	Overall cohort (N = 1403)
Age mean (SD))	59 (15)	60 (15)	60 (15)	60 (15)
Male number (%)	283 (66)	297 (66)	295 (65)	875 (65)
Admission APACHE II score mean (SD)	20 (8)	20 (8)	21 (9)	20 (8)
Admission FCI mean (SD)	1.6 (1.5)	1.7 (1.4)	1.6 (1.5)	1.6 (1.4)
Sepsis at baseline number (%)	282 (66)	296 (66)	300 (66)	878 (66)
Diagnostic category number (%)				
Medical	268 (62)	270 (60)	285 (62)	823 (62)
Surgical	130 (30)	140 (31)	141 (31)	411 (31)
Unavailable	32 (7)	42 (9)	31 (7)	105 (8)
	Visiting relative assessments			
	Dexmedetomidine group (N = 153)	Clonidine group (N = 168)	Propofol (N = 164)	Overall cohort (N = 485)
Age mean (SD)	60 (14)	61 (14)	60 (15)	60 (15)
Male Number (%)	95 (62)	100 (60)	108 (66)	303 (63)

Admission APACHE II score mean (SD)	20 (9)	20 (7)	21 (8)	20 (8)
Admission FCI mean (SD)	1.7 (1.5)	1.9 (1.4)	1.6 (1.5)	1.7 (1.5)
Sepsis at baseline number (%)	104 (68)	111 (66)	114 (70)	329 (68)
Diagnostic category number (%)				
Medical	103 (67)	105 (63)	110 (67)	318 (66)
Surgical	41 (27)	52 (31)	43 (26)	136 (28)
Unavailable	9 (6)	11 (7)	11 (7)	31 (6)

FCI, Functional Co-morbidity index

Table 3: Bedside nurses’ assessments of ability to communicate pain and cooperate with care.

Outcome	All available care periods included				Care periods restricted to RASS score -3 or greater			
	Dexmedetomidine (N = 430)	Clonidine (N = 452)	Propofol (N = 457)	Overall (N = 1339)	Dexmedetomidine (N = 418)	Clonidine (N = 434)	Propofol (N = 439)	Overall (N = 1291)
Number of care periods for which patient was able to communicate pain Number (percentage)	319 (74)	315 (70)	313 (69)	947 (71)	316 (76)	313 (72)	309 (70)	938 (73)
Number of care periods per patient during follow-up for which communication of pain data were available Mean (SD)	12 (12)	12 (11)	12 (12)	12 (12)	10 (10)	11 (11)	11 (11)	10 (11)
Number of care periods per patient during follow-up for which patient able to communicate pain Mean (SD)	5 (8)	4 (7)	4 (8)	5 (7)	5 (8)	4 (7)	5 (8)	5 (7)
Percentage of care periods per patient during follow-up for which patient was able to communicate pain Mean (SD)	39% (35)	35% (33)	33% (33)	36% (34)	44% (36)	40% (35)	37% (35)	40% (35)

Ability to cooperate with care	Dexmedetomidine (N = 413)	Clonidine (N = 438)	Propofol (N = 438)	Overall (N = 1289)		Dexmedetomidine (N = 401)	Clonidine (N = 419)	Propofol (N = 420)	Overall (N = 1240)
Number of care periods for which patient was able to cooperate with care Number (percentage)	329 (80)	325 (74)	329 (75)	983 (76)		319 (80)	319 (76)	319 (76)	957 (77)
Number of care periods per patient during follow-up for which cooperation with care data were available Mean (SD)	11 (10)	11 (11)	11 (11)	11 (10)		9 (9)	10 (10)	9 (10)	9 (10)
Number of care periods per patient during follow-up for which patient able to cooperate with care Mean (SD)	5 (7)	4 (7)	5 (7)	5 (7)		5 (6)	4 (7)	5 (7)	5 (7)
Percentage of care periods per patient during follow-up for which patient was able to cooperate with care Mean (SD)	42% (33)	38% (33)	38% (33)	39% (33)		46% (35)	42% (34)	42% (34)	43% (34)

Table 4: Visiting relative assessments of whether patient appears awake, seems comfortable, and whether they feel able to communicate with patient.

Outcome	All available days included				Days restricted to RASS score -3 or greater			
	Dexmedetomidine (N = 148)	Clonidine (N = 164)	Propofol (N = 162)	Overall (N = 474)	Dexmedetomidine (N = 144)	Clonidine (N = 158)	Propofol (N = 151)	Overall (N = 453)
Patient appears awake to the visiting relative								
Number of days recorded on which patient appeared awake to the visitor Number (percentage)	109 (74)	108 (66)	100 (62)	317 (67)	107 (74)	108 (68)	99 (66)	314 (69)
Number of days during follow-up for which data available regarding whether patient appears awake to the visitor visiting relative Mean (SD)	3 (3)	3 (3)	3 (3)	3 (3)	3 (3)	3 (3)	3 (3)	3 (3)
Number of days during follow-up on which patient appears awake to the visitor visiting relative Mean (SD)	2 (2)	2 (2)	1 (2)	2 (2)	2 (2)	2 (2)	2 (2)	2 (2)
Percentage of days during follow-up for which patient appears awake to the visiting relative Mean (SD)	53% (40)	50% (42)	44% (41)	49% (41)	56% (40)	54% (42)	49% (41)	53% (41)

Patient seems comfortable to the visitor	Dexmedetomidine (N = 147)	Clonidine (N = 160)	Propofol (N = 156)	Overall (N = 463)		Dexmedetomidine (N = 143)	Clonidine (N = 154)	Propofol (N = 145)	Overall (N = 442)
Number of days recorded on which patient appeared comfortable to the visitor Number (percentage)	140 (95)	153 (96)	152 (97)	445 (96)		137 (96)	148 (96)	141 (97)	426 (96)
Number of days during follow-up for which data available regarding whether patient appears comfortable to the visitor visiting relative Mean (SD)	3 (3)	3 (3)	3 (3)	3 (3)		3 (3)	3 (3)	3 (3)	3 (3)
Number of days during follow-up on which patient appeared comfortable to the visitor Mean (SD)	3 (2)	3 (3)	3 (3)	3 (3)		3 (2)	2 (3)	3 (3)	3 (3)
Percentage of days during follow-up on which patient appeared comfortable to the visitor Mean (SD)	88% (26)	90% (24)	93% (19)	90% (23)		89% (25)	91% (23)	93% (20)	91% (23)
Visiting relative feels they can communicate with the patient	Dexmedetomidine (N = 150)	Clonidine (N = 158)	Propofol (N = 157)	Overall (N = 465)		Dexmedetomidine (N = 146)	Clonidine (N = 151)	Propofol (N = 147)	Overall (N = 444)
Number of days for which the visitor feels they can communicate	107 (71)	102 (65)	104 (66)	313 (67)		107 (73)	100 (66)	102 (69)	309 (70)

with the patient Number (percentage)									
Number of days during follow-up for which data available regarding whether relative feels able to communicate with patient Mean (SD)	3 (3)	3 (3)	3 (3)	3 (3)		3 (2)	3 (3)	3 (3)	3 (3)
Number of days during follow-up for which the visitor feels they can communicate with the patient Mean (SD)	2 (2)	2 (2)	2 (3)	2 (2)		2 (2)	2 (2)	2 (3)	2 (2)
Percentage of days during follow- up for which the visitor feels they can communicate with the patient Mean (SD)	54% (41)	52% (44)	52% (42)	52% (42)		56% (41)	54% (44)	57% (43)	56% (42)

Discussion

In this pre-planned analysis of the A2B trial⁹ we found that, overall, bedside nurses felt patients were able to communicate pain in a mean 36% of care periods. Comparing each alpha2-agonist to propofol-based sedation, dexmedetomidine was associated with greater probability of being able to communicate pain; the probability for clonidine was similar to propofol-based sedation. Overall, bedside nurses felt patients were able to cooperate with care for a mean 39% of care periods, with no significant differences between either dexmedetomidine or clonidine and propofol. Overall, visiting relatives felt patients appeared awake for a mean of 49% of days they visited. Comparison of dexmedetomidine and clonidine to propofol-based sedation suggested effects that favoured both alpha2-agonists compared with propofol, but the confidence interval only excluded a null effect for the dexmedetomidine to propofol comparison. Relatives felt patients appeared comfortable for a mean of 90% of days they visited. There were no marked differences between groups, but the observed effects and confidence intervals indicated a trend towards greater perceived comfort with propofol. Finally, relatives felt they could communicate with patients on a mean of 52% of days they visited, with no apparent differences between the groups.

To our knowledge, this is the first exploration of nurses' and relatives' perceptions of patient comfort and ability to communicate during ICU sedation. Embedding this sub-study within a randomised trial enabled direct comparison between sedation with the three strategies. The proportion of trial patients (>90%) and numbers of care periods with data (mean 11-12 care periods, equating to 6 days) was high for the bedside nurse responses. Given the study occurred in 41 ICUs over 5 years, with data recorded by nurses providing clinical care, it is likely that several thousand different nurses contributed opinions. The study therefore has high validity for representing nurse opinions, based on the questions asked. Approximately a third of patients had opinions from visiting relatives, and for a smaller number of days per patient (mean 3 days). The smaller sample resulted from several factors. First, the ethics committee approval only allowed the relative who had provided consent for each participant in the trial to provide views, which restricted data to days on which they visited. Second, the COVID19 pandemic had an extended effect on relative visiting behaviours during much of the trial. Despite this, we obtained views from 474 different relatives distributed evenly across the three groups. The patients in whom relative responses were available had similar baseline characteristics to the overall trial cohort, suggesting inclusion bias was unlikely.

Guidelines recommend the clinical assessment of sedation state, pain status, and cognition using validated scales designed to have high discriminative ability and inter-rater consistency.⁷ As such they do not reflect nurses' personal views and nurse preferences. These can be discordant with guidelines, for example in relation to performing sedation breaks and/or maintaining wakefulness,

and because of concerns about patient comfort, distress and safety.¹¹⁻¹³ Factors such as personal beliefs and previous experiences, for example adverse events, are also potentially important,¹¹⁻¹⁴ and practices such as increasing overnight sedation to promote sleep and safety are common.¹⁵ Our approach sought nurse and relative views based on their personal opinion, providing novel insights into sedation quality from these perspectives. This is relevant to understanding sedation practice and clinician preferences, given the complex interplay of factors involved in sedation delivery.^{11 14}

Dexmedetomidine aims to achieve light sedation where patients are readily roused, more cooperative, and interactive when stimulated.⁷ This may be mediated by clearer cognition and reduced delirium.¹⁶⁻¹⁸ Analgesic properties also potentially contribute to patient comfort. These benefits underpinned our hypothesis that nurse and relative views would demonstrate superiority compared with propofol. In the A2B trial, patients first achieved the target RASS of -2 after a median 24 hours in all three groups, and on around 75% of intervention days, demonstrating that attempts were made to maintain light sedation.⁹ This was also reflected in the high proportion of care periods available for analyses restricted to RASS score -3 or greater. Our findings are consistent with the proposed benefits of dexmedetomidine in relation to nurse communication and relative perceptions of wakefulness, although confidence intervals were wide. The lesser effect with clonidine is consistent with its much lower alpha2-receptor selectivity. In the trial, we found no differences between the groups in rates of unnecessary deep sedation, delirium or pain behaviours, but agitation occurred at a 50% higher rate in both alpha2-agonist groups compared with propofol.⁹ This might explain why nurses reported no significant differences in the ability to cooperate with care, and relatives reported no significant differences in perceived ability to communicate. Relative assessment of comfort may have had a ceiling effect, as mean proportions were 88-93% across the groups. The trend to greater perception of comfort with propofol compared to both alpha2-agonists was unexpected, but might be explained by the greater agitation rates with alpha2-agonists. For all questions the confidence intervals from the modelling included a wide range of potentially important differences between the groups, perhaps reflecting the sample size available for analysis, diversity of views from nurses and relatives, and differences between individual patients.

Strengths of this sub-study include the randomised design, large sample size especially for the nurse data, and pragmatic real-world context. Patients were also closely involved throughout the design, data collection, analysis, and interpretation of the data. Our analytic approach included modelling that maximised use of available data with adjustment for site and multiple observations within participants. However, our study has limitations. Data were not available on every day, with only a third of patients contributing relative data. The populations with data were similar to the overall trial, but we cannot exclude some inclusion or response bias. Group allocation was not blinded,

which might have influenced subjective responses, especially among bedside nurses. Although sedation was randomised, most patients in the alpha2-agonist groups also received some propofol, at around a third of usual care dose, which meant the responses did not reflect sedation with alpha2-agonists alone in most cases. The majority of patients also received opioid infusions at the discretion of clinical teams.

Future research should further investigate the differences in perceived sedation quality from carer and relative perspective with different drugs and sedation strategies, potentially using opinion-based tools as in this study and/or qualitative methods. To investigate whether differences translate into improvements in clinical or patient-centred outcomes, the relationship between improved communication or wakefulness and changes in care decisions (e.g. earlier weaning or mobilization), or family satisfaction could be explored.

In conclusion, although the A2B trial found no significant differences in objective measures of sedation status (RASS score), delirium (CAM-ICU scores), or pain behaviours this sub-study suggested that nurses may perceive patients receiving dexmedetomidine as better able to communicate pain compared with propofol. There did not appear to be perceived differences in ability to cooperate with care for either alpha2-agonist versus propofol. Visiting relatives perceived patients appeared more awake with dexmedetomidine compared with propofol. Perceptions of comfort and ability to communicate were not notably different between groups, although confidence intervals did not rule out large effect sizes. Given the primary and other key clinical outcomes in the A2B trial were not superior with dexmedetomidine or clonidine compared to propofol, the implications of these findings for optimising ICU sedation are uncertain and reflect the complexity of ICU sedation practice.

Acknowledgements

The A2B trial investigators, who all contributed to the data presented in this analysis, are listed in Appendix 1

CRedit Statement

Timothy Walsh: conceptualisation (lead), formal analysis (supporting), funding acquisition (lead), investigation (equal), methodology (equal), administration (equal), supervision (lead), visualisation (equal), writing – original draft (lead), reviewing and editing (equal)

Richard Parker: Data curation (equal), formal analysis (lead), investigation (supporting), methodology (equal), software (lead), validation (lead), visualisation (equal), reviewing and editing (equal)

Leanne Aitken: conceptualisation (supporting), funding acquisition (supporting), investigation (supporting), methodology (supporting), administration (supporting), supervision (supporting), visualisation (equal), reviewing and editing (equal)

Cathrine McKenzie: conceptualisation (supporting), funding acquisition (supporting), investigation (supporting), supervision (supporting), visualisation (equal), reviewing and editing (equal)

Robert Glen: conceptualisation (supporting), funding acquisition (supporting), investigation (supporting), visualisation (equal), reviewing and editing (equal)

Christopher Weir: conceptualisation (supporting), Data curation (supporting), formal analysis (lead), funding acquisition (supporting), investigation (supporting), methodology (lead), administration (supporting), software (supporting), supervision (supporting), validation (lead), reviewing and editing (equal).

Disclosure of interest statement

Timothy Walsh declares receiving a grant from NIHR EME Board as a Co-applicant and a grant from NIHR Programme Board as a Co-applicant. Neither grant related to this work, and funding was to the University of Edinburgh. Christopher Weir declares membership of several boards: HS&DR Commissioned - Board Member (31/07/2013 to 01/03/2016); HS&DR Commissioned R&R (Bird) Sub Board (31/07/2013 to 01/05/2016); HS&DR Funding Committee Member (31/07/2013 to 31/01/2018); and EME - Funding Committee Member (01/07/2018 to 01/07/2022). Richard Parker declares membership of the HS&DR Board (31/07/2013 to 31/05/2018). Cathrine McKenzie declares being in receipt of an NIHR Senior Clinical Practitioner Research Award, and an NIHR Wessex Applied Research Collaborative Research Enhancement Award.

Data-sharing statement

De-identified participant data will be made available to researchers if their proposal for use is within the agreed uses for which participants provided consent, the proposal is approved by the trial team, and any agreements are in place for data-sharing. A data dictionary for the trial data will be made available. All data requests should be submitted to the corresponding author for consideration. Access to anonymised data may be granted following review. #datasaveslives You can find out more about the background to this citation here: <https://understandingpatientdata.org.uk/data-citation>

Patient Data Statement

This work uses data provided by patients and collected by the NHS as part of their care and support. Using patient data is vital to improve health and care for everyone. There is huge potential to make better use of information from people's patient records, to understand more about disease, develop new treatments, monitor safety, and plan NHS services. Patient data should be kept safe and secure, to protect everyone's privacy, and it's important that there are safeguards to make sure that it is stored and used responsibly. Everyone should be able to find out about how patient data is used.

Ethics statement

The A2B trial received ethical approval from the Scotland A REC (18/SS/0085) on 21/08/2018.

Information Governance statement

The University of Edinburgh and NHS Lothian (Co-sponsors) are committed to handling all personal information in line with the UK Data Protection Act (2018) and the General Data Protection Regulation (EU GDPR) 2016/679.

Under the Data Protection legislation, the University of Edinburgh is the Data Controller, and you can find out more about how we handle personal data, including how to exercise your individual rights and the contact details for our Data Protection Officer here (<https://data-protection.ed.ac.uk/contact>). The University of Edinburgh is also the Data Processor

References

1. Vincent JL, Shehabi Y, Walsh TS, et al. Comfort and patient-centred care without excessive sedation: the eCASH concept. *Intensive Care Med* 2016;42(6):962-71. doi: 10.1007/s00134-016-4297-4 [published Online First: 20160413]
2. Ely EW, Truman B, Shintani A, et al. Monitoring sedation status over time in ICU patients: reliability and validity of the Richmond Agitation-Sedation Scale (RASS). *Jama* 2003;289(22):2983-91. doi: 10.1001/jama.289.22.2983
3. Ely EW, Inouye SK, Bernard GR, et al. Delirium in mechanically ventilated patients: validity and reliability of the confusion assessment method for the intensive care unit (CAM-ICU). *Jama* 2001;286(21):2703-10. doi: 10.1001/jama.286.21.2703
4. Payen JF, Bru O, Bosson JL, et al. Assessing pain in critically ill sedated patients by using a behavioral pain scale. *Crit Care Med* 2001;29(12):2258-63. doi: 10.1097/00003246-200112000-00004
5. Train S, Kydonaki K, Rattray J, et al. Frightening and Traumatic Memories Early after Intensive Care Discharge. *Am J Respir Crit Care Med* 2019;199(1):120-23. doi: 10.1164/rccm.201804-0699LE
6. Righy C, Rosa RG, da Silva RTA, et al. Prevalence of post-traumatic stress disorder symptoms in adult critical care survivors: a systematic review and meta-analysis. *Crit Care* 2019;23(1):213. doi: 10.1186/s13054-019-2489-3 [published Online First: 20190611]
7. Devlin JW, Skrobik Y, Gélinas C, et al. Clinical Practice Guidelines for the Prevention and Management of Pain, Agitation/Sedation, Delirium, Immobility, and Sleep Disruption in Adult Patients in the ICU. *Crit Care Med* 2018;46(9):e825-e73. doi: 10.1097/ccm.0000000000003299
8. Walsh TS, Aitken LM, McKenzie CA, et al. Alpha 2 agonists for sedation to produce better outcomes from critical illness (A2B Trial): protocol for a multicentre phase 3 pragmatic clinical and cost-effectiveness randomised trial in the UK. *BMJ Open* 2023;13(12):e078645. doi: 10.1136/bmjopen-2023-078645 [published Online First: 20231210]
9. Walsh TS, Parker RA, Aitken LM, et al. Dexmedetomidine- or Clonidine-Based Sedation Compared With Propofol in Critically Ill Patients: The A2B Randomized Clinical Trial. *Jama* 2025 doi: 10.1001/jama.2025.7200 [published Online First: 20250519]
10. Edinburgh Clinical Trials Unit. Available from: <https://usher.ed.ac.uk/edinburgh-clinical-trials/our-studies/ukcrc-studies/a2b>. Last accessed 23/7/2025
11. Kydonaki K, Hanley J, Huby G, et al. Challenges and barriers to optimising sedation in intensive care: a qualitative study in eight Scottish intensive care units. *BMJ Open* 2019;9(5):e024549. doi: 10.1136/bmjopen-2018-024549 [published Online First: 20190524]
12. Macpherson D, Hutchinson A, Bloomer MJ. Factors that influence critical care nurses' management of sedation for ventilated patients in critical care: A qualitative study. *Intensive Crit Care Nurs* 2024;83:103685. doi: 10.1016/j.iccn.2024.103685 [published Online First: 20240316]
13. Everingham K, Fawcett T, Walsh T. 'Targeting' sedation: the lived experience of the intensive care nurse. *J Clin Nurs* 2014;23(5-6):694-703. doi: 10.1111/jocn.12058 [published Online First: 20130121]
14. Varga S, Ryan T, Moore T, Seymour J. What are the perceptions of intensive care staff about their sedation practices when caring for a mechanically ventilated patient?: A systematic mixed-methods review. *Int J Nurs Stud Adv* 2022;4:100060. doi: 10.1016/j.ijnsa.2021.100060 [published Online First: 20220102]
15. Aitken LM, Emerson LM, Kydonaki K, et al. Alpha 2 agonists for sedation to produce better outcomes from critical illness (A2B trial): protocol for a mixed-methods process evaluation of a randomised controlled trial. *BMJ Open* 2024;14(4):e081637. doi: 10.1136/bmjopen-2023-081637 [published Online First: 20240405]

16. Mirski MA, Lewin JJ, 3rd, Ledroux S, et al. Cognitive improvement during continuous sedation in critically ill, awake and responsive patients: the Acute Neurological ICU Sedation Trial (ANIST). *Intensive Care Med* 2010;36(9):1505-13. doi: 10.1007/s00134-010-1874-9 [published Online First: 20100408]
17. Goodwin HE, Gill RS, Murakami PN, et al. Dexmedetomidine preserves attention/calculation when used for cooperative and short-term intensive care unit sedation. *J Crit Care* 2013;28(6):1113.e7-13.e10. doi: 10.1016/j.jcrc.2013.07.062 [published Online First: 20131018]
18. Maagaard M, Barbateskovic M, Andersen-Ranberg NC, et al. Dexmedetomidine for the prevention of delirium in adults admitted to the intensive care unit or post-operative care unit: A systematic review of randomised clinical trials with meta-analysis and Trial Sequential Analysis. *Acta Anaesthesiol Scand* 2023;67(4):382-411. doi: 10.1111/aas.14208 [published Online First: 20230207]

Appendix 1: The A2B trial investigators

Natalie Pattison
Barry Williams
Louise Rose
Paul Mouncey
John Prowle
David Wellsted
Stephen Brett
Julian Bion
Graeme McLennan
Matt Stevenson
Alistair Nichol
Timothy Walsh
Maria Amamio
Lucy Barclay
Sophie Birch
Kate Briton
Sarah Clark
Jessica Crossan
Katherine Doverman
David Hope
Lucy Macdonald
Corrienne McCulloch
Nicola Rae
Scott Simpson
Jo Singleton
Maggie Wishart
Ruth Thompson
Neill Aitken
Rachel Fairlie
Nabeel Salim
Sam Talbot
Sarah Ackroyd
Valeria Alicino
Euan Allan
Thomas Anderson
Rosemary Andrew
Andrew Baigey
Kenneth Baillie
Calum Barnettson
Ruth Begbie
Richie Biggers
Michael Blaney
Richard Broom
David Birrell
Will Calkin
Keegan Chuavilong

Rebecca Cowden
Thomas Cox
Coral Darjee
Simon Davies
Annemarie Docherty
Luke Dornan
Mark Dunn
Stuart Edwardson
Ross Gillespie
Jane Greenwood
David Griffith
Alasdair Hay
Amy Hu
Ali Hunter
Karen Jones
Helen Jordan
Ancy Joseph
Kallirroï Kefala
Stephanie Kelly
Laura Kemp
Bara Kubanova
Victoria Leng
John Livesey
Nazir Lone
James Lyon
Olivia Mansfield
Dean McAvoy
Aaron McClatchey
Jonathan Miller
Ananda Mirchandani
Peter Moffitt
Steven Morrison
Alexandra Muir
Kieran Nunn
John Ochiltree
Emily Ogden
Matthew Parks
Marc Pass
Rachael Penrose
Harry Putnam
Thomas Quinn
Jonathan Rhodes
Alexander Rollings
Stephen Ross
Ralph Shackleton
Manu Shankar
Sunil Sharma
Iain Slessor

Zack Slevin
Duncan Stickle
Louise Symons
Fiona Walker
Luke Walls
Ian Whiteford
Sue Yin Yong
Neil Young
Kevin Rooney
Michael Kinsella
Brian Digby
Michael Brett
Paul McConnell
Mark Henderson
Radha Sundaram
Lisa Gemmell
Fiona Christie
Philip Henderson
Fiona MacGregor
Steven Henderson
Natalie Rodden
Kirsty Fallon
Lynn Abel
Barbara McLaren
Emma Hughes
Deborah McGlynn
Nicola Thomson
Lauren Walker
Susan Currie
Natasha Parker
Donna Gillan
Farooq Brohi
Sarah Purvis
Michele Clark
Pam Race
Lynne Williams
Ahmed Shahin
Eusebius Nworah
Jonathan Gui
Li-Chin Cheng
Katelyn Stewart
Rebecca Cusack
Mark Tomlin
Clare Bolger
Rachel Burnish
Sue Jackson
Alice Baker
Jonathan Biss

Karen Salmon
Michael Carter
Catherine McKenzie
Razaz Elsheikh
Missy (Anne) Harrison
Charlotte Thomas
James Ward
Andrew Cumpstey
Ahilanandan Dushianthan
Ivan Kemp
Valerie Page
Xiaobei Zhao
Nazril Nordin
Ahmed Hegazy
Elvira Hoxha
Owen Hardaker
Chimenime Ede
Nailia Kotrikova
Acharya Devaraja
Thomas Stambach
Prasun Mukherjee
Mark Louie Guanco
Matthew P Wise
Jade Cole
Helen Hill
Jenny Brooks
Michelle Davies
Rhys Davies
Emma Thomas
Angharad Williams
Lauren Lodhi
Matt PG Morgan
Simon Ridler
Christopher Smith
Maria Faulkner
Alison Ivison
Laura McKay
Helen Jeffrey
Jude Price
Lucy Slater
Angela Davies
Edward Hughes.
Matt Thomas
Dominic Janssen
Ian Thomas
Kate Crewdson
Christopher Newell
Robert Hirst

Stephen West
Agnieszka Skorko
Emma Gendall
Ruth Worner
Beverley Faulkner
Borislava Borislavova
Kati Hayes
Andrew Parsons
Elizabeth Goff
John Sowersby
Annie Wood
Kieran Oglesby
Idrisu Sanusi
Charlie Pope
Andrew Baird
Hayley Blackmore
Robert Healey
Philip Hopkins
Eleanor Corcoran
Gillian Selman
Clare Finney
Evita Pappa
John Smith
Emma Clarey
Maeve Cockrell
Sian Saha
Harriet Noble
Kevin O'Reilly
Maria Depante
Anna Broderick
Marianette Anne
Axalan Burt
Vergara Reena Mehta
Henrik Reschrieter
Sarah Patch
Julie Camsooksai
Sarah Jenkins
Madga Pomichowska
Ken Power
Spike Briggs
Elizabeth Woodward
Christopher Loew
James Bromilow
James Keegan
Matthew Taylor
Emma Langridge
Dinesh Kulandhaisamy
Saah Savage

Yasmin de'Ath
Charlotte Humphrey
Sue Roffe
Matthew Bayliss
Leanne Bartlett
Richard Gordon-Williams
Kate Tatham
Sam Smith
Isabel Noris
Sharjeel Tahir E
mma Yates S
hivali Patel
Tanith Westerman
Sekina Bakare
Hugh Furness
Emma Hunt
Reyhaneh Sadegh Zadeh
Maria Khan
William Sherwood
Claudio Addari
Roshni Manex
Nicole Whitehead
Fred Wilson
Luke Edwards
Kshiteeja Nalk
Sophie Biddle
Suzannah Lant
Francesca Holden
Shree Voralia
Nicola Ocean
Arun Sahni
Prakhar Srilastava
Sultan Iqbal
Shamil Tana
Vishal Venkat Raman
Zoszka Webb
Luke Parker
Arnold Dela Rosa
Miran Kadr
Eleanor Harvey
Ryan Howle
Aatif Husain
Olivia Morley
Sarah Loftus
Jenna Hutchinson
Shaman Jhanji
Ethel Black
David Parkinson

Ravishankar Raobaikady
Mark Borthwick
Christie James
Grace Polley
Neil Davidson
Sally Beer
Paula Hutton
Archana Bashyal
Jean Wilson
Soyamol Mathew
Jung Ryu
Jason Cupitt
Gareth Hardy
Leonie Benham
Robert Downes
Neil Flint
Michael Little
Ravindra Pochiraju
Prematie Andreou
Dawn Hales
Jessica Hailstone
Megha Mathews
Martin Huntley
Lorraine Stephenson
Jacqui Hussey
Hao-Ern Tan
Simon Holbrook
Hayley Kemp David Earl
Richard Innes
Benjamin Plumb
Patricia Doble
Rebecca Purnell
Ashly Thomas
Muhammad Hamza Noor
Waqas Khaliq
Micheal Jennings
Bernd Oliver Rose
Rosaleeta Reece-Anthony
Sagira Khatun
Samantha Dickinson
Jayson Clarke
Charlie Cox
Adam Longley
Tariq Ali
Babita Gurung
Mohamed Moubarak
Alan Williams
Jonathan Ball

Susannah Leaver
Sarah Farnell-Ward
Maria Thanasi
Shreeja Dangol
Vince Ventura
Massimiliano Valcher
Christine Sicat
Nikki Yun
Rebecca Kanu
Maria Maiz Cordoba
Ha Trinh
Karen Lloyd
Romina Pepermans Saluzzio
Lijun Ding
Helen Farrah
Edna Fernandes
Chris Nutt
Jon Silversides
Danny McAuley
Peter McGuigan
Emmet Major
Elliott Lonsdale
Nerielle Fundano
Kathryn Ward
Christine Turley
Aisling O'Neill
Stephanie Finn
Jackie Green
Erin Collins
Julie McAuley
Jeanette Mills
Chris Wright
Michelle Growcott
Iain McCullagh
Stephen Wright
Ian Clement
Jonathan Shelton
Matthew Faulds
Thomas Hellyer
Harriet Morton
Christopher Pollard
Christopher White
Leigh Dunn
Verity Calder
Susan Taylor
Pamela Garcia
Benjamin Brown
James Savage

Maite Babio-Galan
Kimberley Webster
Tessa Wilkinson
Arti Gulati
Tara Shrestha
Carole Hays
Lauren Butler
Fatima Simoes
Margaret McNeil
Ian Storey
Simon Whiteley
Elizabeth Wilby
Susan Trott
Sarah Watts
Shailamma Mathew
Sheila Salada
Adam Neep
Nora Youngs
Clare Howcroft
Matthew Powell
Michael Adlam
Elankumaran Paramasivam
Zoe Friar
David Antcliffe
Stephen Brett
Anthony Gordon
Dorota Banach
Roceld Rojo
Sonia Sousa Arias
Ziortza Fernandez de Pinedo Artaraz
Phoebe Coghlan
Amal Mohammed
Eleanor Jepson
Jenny Wong
Anita Tamang Gurung
Caoimhe O'Dwyer
Sara Perez Guillotin
Maie Templeton
James Hanison
Jonathan Bannard-Smith
Daniel Conway
Shoneen Abbas
Mohamad Aly
Stephen Benington
Teh Eng Hean
Daniel Hayley
Ellen McGuckin
Andrew Martin

Thomas Morris
William Musselbrook
Bhaskar Narayan
Thomas Wright
Chris Wheeler
Melanie Barker
Richard Clark
Emma Connaughton
Rose Jama
Deborah Paripoorani
Rachael Quayle
Anila Sukumaran
Charlotte Taylor
Megan Balmer
Saejohn Lingeswaran
Lauren Edmunds
Katharine Wylie
Andrew Owen
Gavin Perkins
Sean Munnelly
Daniel Park
Jo Gresty
Ellie Reeves
Celina Maliaykal
Teresa Melody
Jacobus Preller
Petra Polgarova
Cristina Bravoelvira
Sofia Teixeira
James Varley
Sapna Sharma Hajela
Kay Elston
Siobhan Campbell
Meike Keil
Muhammad Elbehery
Jocelyn Marshall
Susan Stevenson
Andrew Conway Morris
Prasad Gogineni Venkateskara
Michael Reay
Karen Reid
Rebecca Brown
Chinenyenwa Amareihe
Elliot Yates
Jia Luen Goh
Edward Jones
Aamer Mughal
David Brealey

Niall MacCallum
Samuel Clark
Deborah Smyth
Georgia Bercades
Ingrid Hass
Gladys Martir
Jung Ryu
Anna Reyes
Maria Alexandra Zapata Martinez
Laura Gallagher
Chi Yee Chung
Graeme Sanders
Vipal Chawla
Namrata Maheshwari
Tessa Glazebrook
Hollie Angel
Rebecca Squires
Hayley Dolan
Christopher Donnelly
Lucy Mires
Robert Musalagani
Suzanne Williams
Robin Heij
Peter Young
Mark Blunt
Gayathri Wijewardena
John Gibson
Aricsa Mariya Joshy
Jeremy Bewley
Kieron Rooney
Katie Sweet
Kim Wright
Lisa Grimmer
Denise Webster
Casandra Bazan Lacerot
Rachel Shiel
Eva Maria Hernandez Morano
Christina Coleman
Eleanor Daniel
Oluwatosin Komolafe
Josephine Bonnici
Linda Pipira
Rebekah Johnson
Anna Chillingworth
Ya-Hui Liang
Georgia Efford
Angeliki Kolovou
George Davies

Zoe Garland
Bethany Gumbrell
Ivan Collin
Matthew Gibbins
Thomas Brougham
Agnieszka Skorko
Dan Harvey
William Phipps
Kathryn Harrold
Nick Plummer
Ben Lowe
Paul James
Sara Ahmed
Rukmini Ghosh
Omer Mohamed
Tanushree Santra
James Shilston
Andrew Russell
Viresh Patel
Upasana Topiwala
Habideen Bello
Julia Sampson
Lucy Ryan
Cecilia Peters
Megan Meredith
Louise Conner (Now Hughes)
Lucy Morris
Amy Clark
Alice Baddeley
Lisa McLoughlin
Cate Walton
Treesa Joseph
Anju Thomas
Sophie Lubbock
David Ford
Alexandra McCoy
Tony N'Dungu
Ingeborg Welters
Vinoth Sankar
Alicia Waite
Brian Johnston
David Shaw
Vicki Waugh
Karen Williams
Maria Lopez Martinez
Maria Norris
Maria Arra Carlota Mahiya
Jamie Fernandez Roman

Jin-Xi. Yuan
Silvia Manes
Caitlin Lythgoe
Ibrahim Almafrefji
Josh Colfar
Laura Medhurst
Stephanie Beresford
Sofia Farina
Lema Imam
Syamlam Ali Zachary
Thomas Francesca Bold
Edward Hughes
Katherine Hodson A
leem Morenikeji
Daniel Watkin
Tamas Szakmany
Amy Cardwell
Anne Frawley
Marlies Ostermann
Gillian Radcliffe
Nicholas Barrett
Simon Sparkes
Adam Woodman-Bailey
Eirini Kosifidou
Aneta Bociek
Ellie Hendrie R
osario Lim
Fabiola D'Amato
Sarah Fordyce
Benjie Cendreda
Kyma Morera Vas
Jacqueline Pan
Christopher Meddings
Vladimir Milic
Mike Barker
Jennifer Owusu-Afriyie
Carolin Engelhard
Malcolm Sim
Richard Appleton
Maximilian Ralston
Andrew Arnott
Steven Henderson
Izabela Orlikowska
Sophie Kennedy-Hay
Christopher Murray
Matthew Devine#
Padraig Headley
John McCaffrey

Daniel Donnelly
Richard Young
Samantha Hagan
Victoria Adell
Elizabeth Murphy
Alasdair Hay
Jian Que
Stephen Wilson
Catherine Jardine
Mark Forrest
Emma Collins
Miqdad Ibrahim
Mark Wheeley
Mostafa Kodous
Mathew Blake
Victoria Lacey
Michael Eager
Robin Jootun
Janine Birch